NET WT, 20 LBS.

## PLUS

KILLS WEEDS AS IT FEEDS YOUR LAWN

#### **DIRECTIONS**

Apply anytime during growing season-Spring, Summer or Fall. Always use a mechanical spreader.

SPREADER SETTINGS-On standard models-Set all makes at about 1/5 of total opening. Supergreen Models 1622 E, 1622 F...... Central, Greenfield, Cross Country ...... 51/2 Scotts-Standard Models Only .....4 These settings are furnished as a guide ONLY -- Age, condition of spreader, speed of operation and roughness of terrain can cause variations in rates applied.

#### **HIGHLY EFFECTIVE AGAINST**

WILD CARROT **CURLED DOCK** RAGWEED CHICKWEED

WILD HONEYSUCKLE (MORNING GLORY) **WILD MUSTARD** 

PLANTAIN DANDELION

WILD ONION and many other broadleaf weeds.

The plant food supplies the quick pick-up needed to fill out bare spots left by dead weeds and help prevent re-establishment of weeds by encouraging growth and formation of thick, resistant turf.

#### BE CAREFUL

- 1) Select a time when rain is not expected. (Rain washes weed killer off weeds - reducing kill.)
- 2) For best results apply when grass is damp dew or after watering. 3) Apply evenly - avoid drift to and on plants, flowers and shrubs -
- it can kill them. 4) Chickweeds - Best kill when temperature is low (spring and fall). A repeat application may be required in two weeks.
- 5) Water 2-3 days after application—this will start the feeding.
- Weed killer has started to kill weeds the first day. 6) New lawns should have 2-3 months growth before using this product - DO NOT USE ON BENT GRASS
- 7) Will not burn when applied as recommended.
- 8) To avoid possible injury to St. Augustine Grass, Dichondra, Clover Use at 1/2 recommended rate.
- 9) Do not store near fertilizers, seeds, insecticides, or fungicides.

#### **CAUTION**

May be harmful if swallowed. Avoid contact with skin or eyes. Avoid breathing dust. Avoid contamination of foodstuffs or feeds. Wash thoroughly with soap and water after using.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not reuse empty bag. Dispose of through regular refuse collection system.

ANTIDOTE-EXTERNAL - In case of spillage on skin, wash promptly with soap and water.

MITERNAL - Emetic of 1 tablespoon mustard in

## PLUS

KILLS WEEDS AS IT FEEDS YOUR LAWN

#### DIRECTIONS

Apply anytime during growing season- Spring, Summer or Fall. Always use a mechanical spreader.

#### 

#### HIGHLY EFFECTIVE AGAINST

WILD CARROT BINDWEED
CURLED DOCK PLANTAIN
RAGWEED WILD ONION
CHICKWEED DANDELION

BINDWEED WILD HONEYSUCKLE (MORNING GLORY)
PLANTAIN WILD MUSTARD

WILD ONION and many other broadleaf weeds.

DANDELION

The plant food supplies the quick pick-up needed to fill out bare spots left by dead weeds and help prevent re-establishment of weeds by encouraging growth and formation of thick, resistant turf.

#### **BE CAREFUL**

- 1) Select a time when rain is not expected. (Rain washes weed killer off weeds reducing kill.)
- 2) For best results apply when grass is damp dew or after watering.
- 3) Apply evenly—avoid drift to and on plants, flowers and shrubs—it can kill them.
- 4) Chickweeds—Best kill when temperature is low (spring and fall).

  A repeat application may be required in two weeks
- 5) Water 2-3 days after application—this will start the feeding. Weed killer has started to kill viseds the first day.
- 6) New lawns should have 2-3 months growth before using this product DO NOT USE ON BENT GRASS
- 7) Will not burn when applied as recommended.
- 8) To avoid possible injury to St. Augustine Grass, Dichondra, Clover Use at ½ recommended rate.
- 9) Do not store near fertilizers, seeds, insecticides, or fungicides.

#### CAUTION

May be harmful if swallowed. Avoid contact with skin or eyes.

Avoid breathing dust. Avoid contamination of foodstuffs or

feeds. Wash thoroughly with soap and water after using.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not reuse empty bag. Dispose of through regular refuse collection system.

ANTIDOTE—EXTERNAL—In case of spillage on skin, wash promptly with soap and water.

INTERNAL—Emetic of 1 tablespoon mustard in

tumbler of water. CALL A PHYSICIAN.

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the face hereof. All statements concerning this product apply only when used as directed.

Manufactured for SUPERFILEN PRODUCTS

By DOLD FEED CO., INC.

BUFFALO, N. Y. 14210

U.S. PAT. NO. 3,281,00

POST-MERGER may be used on most established lawn grasses.

Although somewhat slower acting than some other selective weed killers, POST-MERGER is preferable, because of this very fact, so as to minimize injuring the turf.

It has been demonstrated that POST-MERGER will control the following weeds:

Black Mustard
Bedstraw
Common Chickweed
Field Pennycress
Ground Ivy
Knotweed
Lambsquarter
Mouse-ear Chickweed

Pigweed
Plantain
Ragweed
Red Clover
Shepherds Purse
Stitchwort
Wild Radish

White Clover

### CAUTION

Avoid spray drift to susceptible plants as this product may injure tomatoes, beans, peas, ornamentals, etc. (Coarse sprays are less likely to drift.) Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes (or do not use this same equipment for other purposes.) Do not transport or store near seeds, fertilizers, insecticides or fungicides. Do not use in or near greenhouses. Avoid contamination of water intended for irrigation of crops or domestic purposes. Protect from freezing. If allowed to freeze, thaw and re-mix before using.

E. P. A. REG. NO. 11515-15



DO NOT STORE WITH FOODSTUFFS
SEE LEFT PANEL FOR ADDITIONAL CAUTIONS.

WEED KILLER

#### CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid contact with skin, eyes or clothing. Avoid inhaling spray mist. Harmful if swallowed: induce vomiting and call Physician immediately. After using, wash hands and other parts of the body contaminated with concentrate spray or washings.

#### DO NOT REUSE EMPTY DRUM

Return To Drum Reconditioner or Destroy By Perforating or Crushing and Burying in a Safe Place.

#### **IMPORTANT**

Mow lawn first. Do not apply if rain is forecast within 24 hours, and do not sprinkle for 24 hours following application. Apply when weeds are young, as in Spring or Fall. Do not apply during drought or very hot weather conditions. Do not use on newly seeded areas, or if you want clover in the lawn. Reseed, or replace bare spots 3 to weeks after treatment.

#### CAUTION

Do not use POST-MERGER on St. Augustine, Carpet buffalo, dichondra, and bent grasses. These grasses may show sensitivity.

#### **DIRECTIONS**

Established Grasses: Mix 3 oz. POST-MERGER in 4 gals. water to treat 1,000 sq. ft. Use 7 pints POST-MERGER in 40 gals. water to treat 1 acre.

#### CAREFULLY CLEAN SPRAYERS!

Immediately after use flush thoroughly with water. Fill tank with a solution of 1 qt. household ammonia for every 25 gals. of water volume. Spray a small amount through the hose and nozzle — keep the remainder in sprayer 18 to 24 hours. Now, drain tank and flush several times with clear water for tank, hose and nozzle.

**NET CONTENTS** 

GALS.

AN EXCLUSIVE PRODUCT MANUFACTURED BY

SOUTHFIELD

# MODERN RESEARCH CORPORATION

MICHIGAN

NEW YORK . CHICAGO . ATLANTA . ST. LOUIS . FORT WAYNE . NASHVILLE . DETROIT . DENVER . CLEVELAND