

Germicide and Air Refreshner

Red Cap Refresh-R is recommended as an air refreshner and as a disinfectant for such surfaces as bathroom for tures, garbage cans, bed frames, wastebaskets, toilet seats, telephones and other non-porous surfaces

TO DISINFECT: Hold container 6-8 inches away from surface and spray until covered with a fine mist


TO REFRESHEN AIR: Point sprayer upward toward center of room and spray 2 to 5 seconds. Repeat when necessary TO CONTROL MILDEW: Clean surface thoroughly. Hold container 6.8 inches away from surface and spray until covered with a fine mist. Repeat cleaning and spraying process when new growth appears.

89.47%

ACTIVE INGREDIENTS
Ethyl Alcohol 84 55%, Perfume 2 98%; Oil of Lavender 0.73%, n.Alkyl (50%, C.14, 40%, C.12, 10%, C.16) dimethyl benzyl ammonium chlorides 0.58%, Menthol 0.44%, Oil of Eucalyptus 0.19%, INERT INGREDIENTS
Water

100 00%

Rinse For pty Container thoroughly with water and discard it EPA Reg. No. 114502



ACCEPTED WITH COMMENTS

NET CONTENTS 16 HL. 02.

Cold Disinfection

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1½ ounce SAN-FECT per gallon of water for 10 minutes.

Restrurant and Bar Rinse

Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent; rinse thoroughly and immerse in a sanitizing solution containing 1 ounce SANFECT to 4 gallons of water. No Terminal Potable Water Rinse Required.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield, and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

Rinse empty container thoroughly with water and discard.

NET CONTENTS: _____LBS





ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides	
ethylbenzyl ammonium chlorides 5%	
INERT INGREDIENTS:	
100%	

DANGER KEEP OUT OF REACH OF CHILDREN

See other precautionary statements on left side panel.

USE DIRECTIONS:

Household

For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage pails, Sink Tops, Corridors, Classrooms, Offices and shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce SANFECT to 4 gallons of water. Disinfect with 3 ounces SANFECT to 5 gallons of water. Kills Staphylococcus aureus and Salmonella Choleraesuis.

Hospitals

For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use **SANFECT** at 3½ ounces per 5 gallons of water. At this level **SANFECT** will be effective against Pseudomonas aeruginosa PRD-10.

Food Processing Equipment

For sanitization of food processing equipment and food utensils dilute 1 ounce SAN-FECT per 4 gallons of water to provide 200 ppm of active quaternary. At this level, No Potable Water Rinse is required.

Dairies

To sanitize dairy equipment, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce SAN-FECT to 4 gallons of water (200 ppm). At this level, No Potable Water Rinse is required. Follow recommendations of local Health Board.

E.P.A. R.E.G. No. 11515 3 1

SOUTHFIELD - THE PROPERTY OF ST. LOWIS -- LEORIPHYMYN -- NOT WITH A PROPERTY -- DENVER- CHEVELAND