

FOR PROFESSIONAL USE ONLY

CAUTION

Avoid contact with skin or prolonged breathing of Spray mist. Avoid contamination of feed or foodstuffs. Keep away from Domestic Animals. Keep livestock off treated areas. Wash hands after using. Do not contaminate water used for domestic purposes. Harmful if swallowed.

ANTIDOTE

If taken internally induce vomiting and call a Doctor at once.

NOTE

Do not apply to dry turf. Restore soil moisture by watering thoroughly before application of CRAB-GONE.

Keep the area mowed to about 1" to 1-1/2" during treatment. Good grasses may be temporarily affected, especially in hot dry weather. Watering will revive them.

On new lawns or areas do not treat until after three mowings.

DO NOT USE ON ST. AUGUSTINE OR CENTIPEDE GRASS.

On turf that is predominantly bent or fescue use the lower rate only.

E. P. A. 11515-47



Active Ingredients.	
Disodium Methanearsonate Anhydrous	18.9%
Inert Ingredients	81.1%
	100.00%
Total Arsenic, expressed as elemental, all in water soluble form 7.65%	
Equivalent to 3 pounds DISODIUM METHANEARSONATE HEXA-	
HYDRATE per gallon	

CAUTION: KEEP OUT OF REACH OF CHILDREN

See left panel for additional cautions

DIRECTIONS:

For Young Crab Grass (Two and Three Leaf Stage) — Apply at a rate of 1 pint in 10 to 20 gallons of water per 4000 square feet. Two or three applications are necessary at seven-day intervals.

Branched Crab Grass — Apply at a rate of 1 pint in 10 to 20 gallons of water per 3000 square feet. Two to three applications are necessary at seven-day intervals. NOTE: At temperatures over 85° F apply at a rate of 1 pint in 10 to 20 gallons of water per 5000 square feet.

On Bent or Fescue: Apply at a rate of 1 pint in 10 to 20 gallons of water per 4000 square feet. Two to three applications at seven-day intervals.

For Dallis Grass: Apply at a rate of 2 gallons of CRAB-GONE in 100 gallons of water over one acre. Repeat 2 times at intervals of 5 to 7 days. Observe the same precautions as for Crab Grass Control.

DO NOT REUSE EMPTY DRUM
Return to Drum Reconditioner or Destroy by Perforating or Crushing And Burying In a Safe Place.

NET CONTENTS GALS.

AN EXCLUSIVE PRODUCT SOLD BY

SOUTHFIELD MODERN RESEARCH CORPORATION MICHIGAN

ST. LOUIS FORT WAYNE NASHVILLE DETROIT DENVER CLEVELAND

WHEN TO APPLY

MOD-RID is best applied during Spring or Fall.

Do not re-use shipping containers: **destroy** by crushing and burying where possible. Do not transport or store near insecticides, fungicides, fertilizers, or seeds. Avoid contamination of water **intended** for irrigation of crops or domestic purposes.

CAUTION- This product is toxic to fish. **Keep** out of lakes, streams or ponds. **Apply** this product only as specified on this label.

KEEP FROM FREEZING


CAUTION: Avoid contact with skin, eyes or clothing. Avoid inhaling spray mist.

Avoid spray drift to susceptible plants as **this** product may injure tomatoes, beans, peas, other vegetables, legumes, ornamentals, etc (Coarse sprays are less likely to drift). Do not use in or near greenhouses. Do not use same spray equipment for other purposes if traces of this product in such equipment are likely to be harmful to sprayed vegetation.

ANTIDOTE: If taken internally induce vomiting and call a Physician at once.

NET CONTENT _____ GALS

July 31, 1973



ACTIVE INGREDIENT Potassium Salt of 2,4-Dimethyl-4-chlorophenoxypropionic acid (MCPPE) 1.75%
INERT INGREDIENTS 98.25%
2,4-Dimethyl-4-chlorophenoxypropionic acid (MCPPE) 1.75%
Total Weight 100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid contact with skin, eyes or clothing. Avoid inhaling spray mist. Harmful if swallowed, induce vomiting and call Physician immediately. After using, wash hands and other parts of the body contaminated with concentrate, spray or washings.

DO NOT REUSE EMPTY DRUM

Return To Drum Reconditioner or Destroy By Perforating or Crushing and Burying in a Safe Place.

See Left Panel for Additional Cautions

For the control of Chickweed (Common and Mouse Ear Chickweed) Knotweed, Clover and Common Plantain in Bent-grasses, Bluegrasses, Fescues and their mixtures.

DIRECTIONS

Make sure of soil moisture before applications are made, if a drought occurs, water area well before treating. Do not apply on newly seeded turf until after the **second** mowing. Avoid cutting or mowing turf for at least 24 hours before and after spraying. Do not apply during periods of hot, humid weather or if a heavy rainfall is expected.

For best control, applications should be made during Spring or Fall when the good grasses are vigorous and the unwanted weeds are young.

APPLICATION RATES

GREENS: Two quarts of **MOD-RID** diluted in 100 to 200 gallons of water treats one acre. (1 1/2 oz. in 5 gal. water per 1000 sq. ft.)

DIRWAYS & LAWNS: Three quarts of **MOD-RID** diluted in 40 to 60 gallons of water will treat one acre.

Cameraman's Note
Poor Copy

EPA REG. NO. 11515-48

SOUTHFIELD MODERN RESEARCH CORPORATION MICHIGAN
NEW YORK • CHICAGO • ATLANTA • ST. LOUIS • CINCINNATI • NASHVILLE • DETROIT • DENVER • CLEVELAND