


FOR PROFESSIONAL USE ONLY

ABC CHEMICAL CORPORATION

ACCEPTED
 AUG 20 1975
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, this pesticide
 registered under
 EPA Reg. No. 11515-59
Comments



CAUTION: Do not use on newly seeded lawns until after third mowing. Do not reseed lawns until 3 weeks after last application, and weed plants are dead. Berms, Fescues and Clovers may be temporarily discolored. Do not use on St. Augustine, Carpet, Dichondra or Centipede grasses or on Bent Golf Greens.

WARNING: Harmful if swallowed. Do not get in eyes, on skin or on clothing. In case of contact, wash immediately with warm soap and water. For eyes, flush with clear water and get medical attention. Avoid inhalation of spray mist.

Do not contaminate water used for irrigation or domestic purposes. Avoid contamination of feed, foodstuffs, streams or other water sources. Do not spray or allow spray drift on desirable flowers, trees, shrubs or vegetables. Do not store near fertilizers, seeds, insecticides or fungicides. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Keep out of reach of children and domestic animals.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Licensed for use in the control of crabgrass according to the method of Pat. No. 2,678,265.

ACTIVE INGREDIENTS:	By Weight
Dodecyl Ammonium Methanearsonate	8.00%
Octyl Ammonium Methanearsonate	8.00%
* Octyl Ammonium Salt of 2,4 Dichlorophenoxyacetic Acid	5.44%
INERT INGREDIENTS	78.56%
TOTAL	100.00%
Total Arsenic Metallic all in water soluble form	4.00%
* Equivalent to 2,4 Dichlorophenoxyacetic Acid	3.43%

CAUTION: KEEP OUT OF REACH OF CHILDREN
 See side panel for additional CAUTIONS.
 AN EXCLUSIVE PRODUCT SOLD BY

SUPER-KILL Plus 2,4-D contains a combination of herbicides that will kill CRABGRASS and many BROAD-LEAF LAWN WEEDS in the same application. The broadleaf weeds include Dandelion, Knotweed, Plantain, Pennywort, Purslane and Spotted Spurge.

DIRECTIONS: For best results, apply SUPER-KILL Plus 2,4-D in the late Spring or early Summer when the plant growth is vigorous and before the seed heads have formed. The soil should be moist and the lawn in good growing condition before and three days after treatment. If necessary, irrigate the day before treatment, then again two days after the day of application. Late afternoon is the preferred time for treatment.

Measure the area to be treated and apply the correct amount of chemical evenly over the entire lawn area. Use a hose and sprayer or any applicator that will deliver a coarse spray preferably in a "fan" pattern.

Two or more treatments at seven day intervals may be necessary until weed plants are killed. Applications may be necessary later in the season as new seedlings germinate. The amount of chemical to be applied is dependent on the temperature.

FOR TEMPERATURES 80° AND ABOVE: Dilute 4 oz. (1/2 cupful) in 2.5 gallons of water for 500 sq. ft.

FOR TEMPERATURES BELOW 80°: Dilute 8 oz. (1 cupful) in 2.5 gallons of water for 500 sq. ft. of lawn.

E.P.A. REG. NO. 11515-59
 E.P.A. EST. 11515-MI-1
 NET CONTENTS GALS.

ACCEPTED WITH COMMENTS

ABC CHEMICAL CORPORATION

14288 MEYERS RD DETROIT, MICHIGAN 482 (313) 931-3010

ACCEPTED

OTHER WEEDS CONTROLLED

BEDSTRAW, BLACK MEDIC, BUCKHORN, CHICORY, DOCK, HEAL ALL, LAMBS QUARTER, LESPEDEZA, MALLOW, MORNING GLORY, PEPPERGRASS, PIGWEED, POISON OAK, PURSLANE, SHEPHERD'S PURSE, SPEEDWELL, WILD CARROT, WILD LETTUCE, YARROW.

DIRECTIONS FOR USE

Do not allow spray to get on vegetables or ornamental plants since injury may result. Since injury can be expected when used on Centipede or St. Augustine grass lawns, "Spot-Spray" weeds only. Clover present will be damaged if sprayed. Repeat if necessary for hard to kill weeds.

WHEN TO USE: Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall or in the spring, as needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot Spray" weed treatment, however, may be made at any time with a small pressure sprayer, spraying only the weeds at ground level. Wait 3 weeks before putting new grass seed or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results mow before treating. Do not water lawn within 24 hours after treatment.

HOW MUCH TO USE - ENTIRE LAWN

HOSE-END SPRAYER: Use a hose-end sprayer designed for lawns. Put recommended amount of chemical (Column 1) in bottle and add water to indicated level (Column 2), and spray on area indicated. For example—to spray 625 sq. feet, measure 10 ozs. (20 tbsp.) in jar and add water to the 10 gallons level and mix. Turn on hose and spray as evenly as possible over 625 sq. feet.

Col. 1	Col. 2	Col. 3
Amt. SELECT-1000 to use (Tbsp.) (Fl. Oz.)	Fill Bottle to (Gallons)	Area to Treat (Sq. Feet)
10	5	312
20	10	625
30	15	938
40	20	1250

NOTE: Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage ornamental or garden plants.

SELECT-1000

SELECTIVE WEED KILLER

FOR
LAWNS, VACANT LOTS, PLAYGROUNDS

Controls many surface creeping and lawn broadleaf weeds — without injury to most ornamental grasses — Blue Grass, Bermuda or Zoysia.

Contains: *TRIMEC Herbicide; 2, 4-D, MCPP, DICAMBA

Also effective against chickweed, clover, dandelion, spurge, knotweed, plantains, henbit, poison ivy, wild onion, ragweed.

1 PINT COVERS 1000 SQ. FT.

* Trimec® is a registered trademark of P.B.I. — Gordon Corp. U.S. Patent #3,284,186

CAUTION:
KEEP OUT OF REACH OF CHILDREN
See side panel for complete caution statement

KEEP FROM FREEZING

Sold By
CONTROL CHEMICAL CORPORATION
2090 ROUTE 110
FARMINGDALE, N.Y. 11735

THREE COLUMN USE

PRESSURE SPRAYER: Use a pressure sprayer designed for lawns. Put recommended amount of chemical (Column 1) in bottle and add water to indicated level (Column 2), and spray on area indicated. For example—to spray 625 sq. feet, measure 10 ozs. (20 tbsp.) in jar and add water to the 10 gallons level and mix. Turn on hose and spray as evenly as possible over 625 sq. feet.

Col. 1	Col. 2	Col. 3
Amt. SELECT 1000 to use (Tbsp.) (Fl. Oz.)	Fill Bottle to (Gallons)	Area to Treat (Sq. Feet)
10	5	312
20	10	625
30	15	938
40	20	1250

HOW MUCH TO USE: SPOT SPRAY

To spray small areas or individual weeds, do not use hose-end sprayers. Mix 1 oz. (2 tbsp.) in quart water in pressure sprayer and spray as required being certain to "wet" each weed.

CAUTION:

Harmful if swallowed. Avoid skin, eye and clothing contact. In case of contact, flush with water. Avoid contamination of feed or foodstuffs.

CAUTION:

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solution near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable bentgrass or clovers are present. When container is empty, wash thoroughly and destroy. Never re-use. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass and centipede grass.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

Active Ingredients:	By Weight
*Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	0.92%
**Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	2.02%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	0.21%
Dimethylamine salts of related compounds	0.03%
Inert Ingredients:	96.82%
Total	100.00%
*2 (2-methyl-4-chlorophenoxy) propionic acid equivalent 0.755% or 0.065 pounds per gallon	
**2, 4-dichlorophenoxyacetic acid equivalent 1.68% or 0.144 pounds per gallon	
***3, 6-dichloro-o-anisic acid equivalent 0.172% or 0.015 pounds per gallon	