

USE DIRECTIONS:

DISINFECTION

Use at 2 ounces per gallon. This product will be effective against *Pseudomonas aeruginosa* PRD-10 at this level. This has been substantiated by the AOAC Use Dilution Confirmation Test. At this level, the product is fungicidal against pathogenic fungi.

For best results, use Q-TROL with a cloth or mop and apply to walls, floors and other hard surfaces.

For heavy soil or organic matter, a pre-cleaning step is recommended.

SANITIZATION

If used for sanitization of previously cleaned food equipment or food contact items, limit the level of active quaternaries to 200 ppm (2.8 ounces per 5 gallons of water.) Rinse with potable water prior to reuse.

This product when used on environmental, inanimate, hard surfaces at 2 ounces per gallon of water, is effective against Influenza A₂, Herpes Simplex, Adenovirus type 2, and Vaccinia viruses.

PHENOL COEFFICIENTS:

Guaranteed	Minimum
<i>S. aureus</i>	44
<i>S. typhosa</i>	28

Rinse empty container thoroughly with water and discard.

WARRANTY

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its mis-use as such, or in combination with other materials. Statements concerning the use of these products are not to be construed as recommending the infringement of any patent, and no liability or infringement arising out of any such use is assumed.



Q-TROL

**CLEANER
DEODORIZER**

**DISINFECTANT
FUNGICIDE**

Designed for Hospital and Institutional Use.

ACTIVE INGREDIENTS:

- n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
dimethyl benzyl ammonium chlorides 2.25%
- n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl
ethylbenzyl ammonium chlorides 2.25%
- Sodium Carbonate 3.00%
- Tetrasodium ethylenediamine tetraacetate 1.00%

INERT INGREDIENTS: 91.50%

100.00%

WARNING

KEEP OUT OF REACH OF CHILDREN

See right side panel for additional
precautionary and first aid statements.

Sold by
CONTROL CHEMICAL CORP.
2090 ROUTE 110
FARMINGDALE, N.Y. 11735

EPA Reg. No. 11524-6
EPA Est. No. 11524-NY-1

Net Contents: gals.

Q-
tic
ho
cre
ass
flo
an
mc
ag

Wh
eq
pp

Ca
clo

In
for
col
wh
qu



Q-TROL

**CLEANER
DEODORIZER**

**DISINFECTANT
FUNGICIDE**

Designed for Hospital and Institutional Use.

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	
dimethyl benzyl ammonium chlorides	2.25%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl	
ethylbenzyl ammonium chlorides	2.25%
Sodium Carbonate	3.00%
Tetrasodium ethylenediamine tetraacetate	1.00%

INERT INGREDIENTS:	91.50%
	<hr/>
	100.00%

WARNING

KEEP OUT OF REACH OF CHILDREN

See right side panel for additional
precautionary and first aid statements.

Sold by

CONTROL CHEMICAL CORP.

2090 ROUTE 110

FARMINGDALE, N.Y. 11735

EPA Reg. No. 11524-6

EPA Est. No. 11524-NY-1

Net Contents: gals.

Q-TROL is a cleaner, disinfectant and sanitizer, designed for institutional use such as Hospitals, Nursing Homes and Institutions where housekeeping will be of prime importance in controlling the hazard of cross-infection from environmental surfaces. Q-TROL can provide the assurance you need in this area, when used as directed. For use on floors, walls, sink tops, garbage pails, telephones, rest rooms, bed pans and operating rooms. Effective against Staphylococcus aureus, Salmonella choleraesuis and Pseudomonas aeruginosa. Also effective against pathogenic fungi, Trichophyton interdigitale.

When used for sanitization of previously cleaned food processing equipment and food contact items, limit active quaternaries to 200 ppm. and rinse with potable water prior to reuse.

WARNING

KEEP OUT OF REACH OF CHILDREN.

Causes eye and skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Call a physician.