

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION**

Avoid contact with eyes, skin or clothing. In case of contact immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Harmful if swallowed. Do not induce vomiting. Call physician immediately.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 120°F may cause bursting. Never throw container into fire or incinerator.

DIRECTIONS FOR USE

Shake well before each use. Remove protective cap. Spray opening away from person. Push button to spray. Do not use in edible product areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. IN THE HOME — Remove or cover exposed foods. Avoid contamination of cooking utensils and food preparation surfaces. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

STORAGE — Store at temperatures below 120°F. Keep away from heat or open flame.

DISPOSAL — Do not reuse empty container. Wrap container and put in trash collection.

USE WITH CAUTION on young tender foliage. Avoid heavy wetting. Plants such as flowering quince, ash and camellias may be subject to burn under certain conditions.

For adult whiteflies, two spotted mites and mealybugs be sure to direct spray to contact these pests. Thoroughly cover the infested area. Use several repeat applications at two day intervals if necessary.

PROTECT HOUSE PLANTS: Spray infested African violets, philodendron, geraniums, begonias, coleus for listed pests. Leave the sprayed area when done indoors, and ventilate area after treatment is completed.

NUISANCE PESTS-ON OR ENTERING BUILDINGS: Wasps: Applications should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary. Spray directly on stray wasps that enter buildings.

Spiders: Direct short burst of spray on spiders and repeat as necessary.

Earwigs: Spray directly on earwigs in crevices and other hiding places on outdoor porches, patio floor and doorway entrances.

P/P INSECTICIDE NO. 35

KILLS: Houseflies, mosquitoes, gnats, wasps, roaches, earwigs, two spotted mites, sowbugs, spiders, ants, deerflies, horse flies, aphids (plant lice), leafhoppers, exposed thrips, adult whiteflies, and fleas.

ACTIVE INGREDIENTS

Resmethrin [5-(phenylmethyl)-3-furanyl] methyl 2, 2-dimethyl-3-(2-methyl-1-propenyl) cyclopropanecarboxylate	0.200%
--	--------

Related compounds	0.027%
-------------------	--------

d-trans Allethrin (allyl homolog of Cinerin I)	0.125%
--	--------

Related compounds	0.009%
-------------------	--------

Aromatic petroleum hydrocarbons	0.265%
---------------------------------	--------

INERT INGREDIENTS	99.374%
--------------------------	----------------

TOTAL	100.000%
--------------	-----------------

*Cis/trans isomers ratio: max. 30% (±) cis and min. 70% (±) trans.

†U.S. Patent No. 3,465,007

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Do not induce vomiting. Call physician immediately.

If On Skin: Wash with plenty of soap and water.

If In Eyes: Flush eyes with plenty of water for at least 15 min.

SEE SIDE PANEL FOR PRECAUTIONARY STATEMENTS

EPA Reg. No. 11525-36

EPA Est. No. 11525-IL-1

NET CONTENTS

ACCEPTED

JUN 18 1973

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 11525-36

HOUSEHOLD PESTS: Kill houseflies, mosquitoes and gnats. Close all doors and windows. Point nozzle upward. Do not spray mist to all parts of the room, especially windows and other light resources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

KILLS CRAWLING INSECTS: Roaches, earwigs, sowbugs and spiders. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects where possible.

ANTS: Spray trails, nests and points of entry. Spray on ants when possible. Repeat as necessary.

ANIMAL PESTS: Fleas on Dogs, Cats: Apply directly to animals along back, under forelegs and on other areas. Part hair to contact skin and insure good coverage. Hold nozzle about one foot away while spraying. Apply until hair is thoroughly moist. Do not directly spray in ears, eyes, scrotum, anus or mouth of animal. Repeat within 24 hours of first treatment if necessary. To reduce reinfestation, thoroughly spray the animal's bedding and other resting areas. Treat these areas at the rate of two seconds per linear foot while spraying at a distance of 12 to 18". Also spray entire inside surface of dog houses. Repeat as necessary.

OUTDOORS-TO REDUCE ANNOYANCE FROM FLYING INSECTS: Reduce mosquitoes, gnats (midges), houseflies, deerflies, horse flies and flying moths (millers) by spraying their resting areas. Follow precautions listed above. Hold container 18 to 24" from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard of space, while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

OUTDOORS-TO PROTECT ORNAMENTAL PLANTS: To control aphids, exposed thrips, adult whiteflies, leafhoppers and two spotted mites on smaller rhododendrons, pines, spruce, hemlock, spirea, dogwood, palm, mimosa, toyan, viburnum, euonymus oaks, maples, locust, forsythia and althea, spray directly on insects. Hold container 18 inches from the infested plants. Spray in short spray bursts of one to three seconds or spray at the rate of one linear foot of growth per second. Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control. Spray peonies, iris, zinnia, snapdragons, ageratum, chrysanthemums, nasturtiums, daisy, geraniums, marigolds, daphne, fatshedera, ivy, tobira, queen sage, and broad leaf acacia for control of aphids, exposed thrips adult whiteflies and leafhoppers. Spray as directed above.