### Precautionary Statements Hazards to Humans and Domestic Animals

## **CAUTION:**

Harmful if swanowed. Avoid inhalation of mist. This formula is an effective insecticidal aerosol for use in timed dispensing units during the "after hours" when food handling establishments, restaurants, stores, host itals, food processing plants and other institutions where flies, mosquitoes and gnats are an annoyance. Do not use in nurseries or rooms where infants, ill or aged persons are confined. Cover fishbowls while spraying. Remove birds and other pets from area to be sprayed.

### STATEMENT OF PRACTICAL TREATMENT

If swallowed: Do not induce vomiting. Call physician immediately.

If on Skin: Wash with penty of soap and water. If in Eyes:, Flush eyes with plenty of water for at least 15 minutes.

### PHYSICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

### **STORAGE AND DISPOSAL**

Storage: Store in a cool area away from heat or open flame.

Disposal: Do not reuse empty container. Wrap container and put in trash collection.

#### Made by

# PETERSON/PURITAN, Inc.

anville, illinois 61832

## P/P INSECTICIDE FOR TIMED DISPENSER FOR USE IN AUTOMATIC TIMED DISPENSER DURING "AFTER HOURS"

This formula is an effective insecticidal aerosol for use in timed dispensing units during the "after hours" when food handling establishments, restaurants, stores, hospitals, food processing plants and other institutions where flies, mosquitoes and gnats are an annoyance would not be in operation. All food processing surfaces should be covered or thoroughly cleaned before using. Avoid contamination of feed and foodstuffs.

## **ACTIVE INGREDIENTS:**

TRESMETHRIN	*[5-(Phanyimethyl)-3-furanyi] methyl 2, 2-dimethyl-3-(2 methyl-1-propenyi) cyclopro	
	panecarboxylate	1.00% w/w
Kerosene	* * * * * * * * * * * * * * * * * * * *	18.96%
INERT INGREDIENTS:		80.04%
2	4) 1	100.00%

\*Cis/trans isomers ratio max. 11% (±) cis and min. 89% (±) trans. 1U.Ş. Patent No. 3,465,007

## CAUTION KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL

NET-WT., 12 OZ.

## DIAECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.


This aerosol is designed for use in adjustable automatic activators set to dispense a 100 mg. dose every 15 minutes for 8 hours while the plant or institution is not in operation.

Use one aerosol in an automatic activator for each 6,000 cubic feet of room space to be treated. Install the activator on a wall approximately 7 feet from the floor. Do not install directly over any food handling or dispensing area. Never locate automatic activators in drafty hall ways or where air currents are excessive. Select areas where the air currents are least noticeable. The rate of air change, shape of space to be treated, humidity and temperature affect insect activity and insect control. One unit per 6,000 cubic feet shoud be considered as an average ratio. Adjust the ratio depending upon the conditions under which the automatic activator is used.

When properly located, the effectiveness of the SBP-1382 formula will be noticeable within one hour after the activator is started. After one hour of operation, enough insecticide is suspended in the air to kill a substantial proportion of the flies, mosquitoes and gnats present.

This 12-ounce aerosol contains enough material to operate 825 hours, or 103 days on an 8-hour basis. It is suggested the aerosol be changed once every 3 months on an 8-hour use basis. Also the activator should be periodically checked to assure it is operating only on the off hours period.

As with any method of insect control, improvedperformance is achieved when good sanitary practices and other supplementary fly control methods are used.



EPA Reg. No. 11525-48

EPA Est. No. 11525-

40



-----

.