# MICRO-GEN BP300 INSECTICIDE

**ACTIVE INGREDIENTS:** 

Pyrethrins

\*Piperonyl butoxide, technical

N-octyl bicycloheptene dicarboximide Petroleum distillate

81.00% \*Consists of 4.80% (butylcarbityl) (6-propylpiperonyl) ether and 1.20% related compounds. \*\*MGK 264 Insecticide Synergist

Space Spray - Use undiluted to kill accessible stages of insects\* infesting dried fruit products in storage or being processed:

Close room or warehouse and shut off all ventilating systems. Use a spraying device similar to the Micro-Gen Corp. equipment to dispense this products as a fine mist in the air above the trays and shelves. Do not apply to fruit directly. Use a dosage not to exceed 1/2 gallon of this spray per 50,000 cubic feet of room space. Repeat application at this dosage in any given month. Leave the room closed for one hour. Do not remain in treated areas and ventilate area before re-entry.

> \*Indian Meal Moth, Dried Fruit Beetle, Confused Flour Beetle, Saw-toothed Grain Beetle and Roaches.

Space Spray — Use undituted to kill. the adult and accessible forms of insect pests\* of stored food (stored in multiwall paper bags or in cloth bags) in warehouses, storage rooms and similar locations:

Close room or warehouse and shut of all ventilating systems. Use a spraying device similar to the Micro Gen equipment to dispense this product. Use dosage of one fluid curve of this spice. per 1,000 cubic feet of Repeat applications as insect control. Leave for one bour. Do areas

etie,

Space Spray mo**şqjiitdes** <u>al</u>i islate as the vale trate to two parts of at a time when the temperature

(75°F or less) and the wind velocity is approximately five MPH. With equipment similar to the Micro-Gen equipment, spray at a dosage rate of fifteen ounces per minute upwind of the area to be controlled. If the equipment is truck mounted, keep the speed to approximately five MPH. Repeat application as required.

3.00%

6.00%

10.00%

Space and Contact Spray - Use undiluted to kill roaches, silverfish, waterbugs, earwigs, ants, spiders, and beetles in sheltered situations:

With equipment described above, apply the spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Do not allow spray to contact stored food. Use at a dosage rate of approximately one-half fluid ounce per 1,000 cubic feet of room space. Repeat dosage as necessary for control Leave the room closed for one hour. Do not remain in treated areas and ventilate area before re-

### KEER OUT OF REACH OF CHILDREN

vallowed. Do not apply tod. Cover or remove any dod processing equipment application. Do not apply while sing is underway. After spreying in meat packing plants, bakeries and other food processing wash all equipment, benches, etc. where exposed food will andled, with portable water.

drum. Return to destroy by perd burying in a

> efore each use. and store in NÕT OUR OR SPILL

ents: One Gailon

### GEN CORP

porate Offices 4438 Centerview Dr Antonio, Texas 78228

EPA Reg. No. 11540-1

## ACCEPTED

11540-1