For Broadleaf Weed Control in Small Grains, Rice, Flax, Grassland and N

Directions for Use

CONTROLS THESE WEEDS

SUSCEPTIBLE (Most Areas)

LESS SUSCEPTIBLE

IN RICE

SELECTIVE SPRAYING Wheat Oats Barley and Rye App
sprayer or airplane Use
susceptible weeds after crop has
up to boot stage Use up to 3
susceptible weeds after crop has
stage Do not spray from boot to

and legumes see the following

SMALL GRAINS
Alfalfa Birdsfoot Trefoil
Field Corn For emergency control
weeds yellow rocket and other suscep-
Make application after cereal is
and before reach boot stage
should provide a
canopy
gallons
at low
Do not
damage to the
with vetch or sweet clover
some risk to other
by a canopy

RICE (California and Other Areas where Recommended): For
plantain and red stem, apply
airplane in 10 gallons of water per
after sedge, nutgrass and bul-
established
than 55 and no later than
stem begins to elongate
deep at spikelet
Do not spray when temperature is over 90 degrees F Do
not spray after rice is in boot stage

FLAX
HAYNES & MORGAN MCPA AMINE
by airplane in sufficient water
Apply only when weed
reaches height and before it comes
may result in severe
present it may be necessary to go
to prevent seed head production
by mower

CAUTION: Avoid contact with skin eyes or clothing. In case of eye contact, flush eyes with plenty of water. Wash exposed skin with plenty of soap and water. Remove and wash contaminated clothing and shoes before re-use.

Avoid spray drifts onto susceptible plants such as vegetables, flowers, grapes, tomatoes, cotton, beans and other legumes, etc. Use coarse sprays as these are less likely to drift. Do not spray at all in the near vicinity of susceptible plants. Do not contaminate water intended for irrigation or domestic use. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes, or do not use same spray equipment for other purposes. Do not transport with or store near fertilizers, seeds, insecticides or fungicides. Be sure before using that the use of this product conforms to all applicable regulations. Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

TOXIC TO FISH: Do not contaminate any body of water by cleaning of equipment and disposing of wastes and containers.

KEEP out of any body of water.

DESTROY empty containers by piercing top and bottom and burying in non-crop land.

Do NOT Store with Foodstuffs

HAYNES & MORGAN

MCPA AMINE

CAUTION

Keep Out of Reach of Children

EPA Reg. No 11544 14

Ingredients

ACTIVE INGREDIENT: Dimethylamine salt of 2-methyl-4-chlorophenoxy- acetic Acid	52.2%
INERT INGREDIENTS	47.8%
	100.0%

Contains the equivalent of 42.6% 2-methyl-4-chlorophenoxyacetic acid equivalent to 4.0 lbs. per gallon at 68 degrees F.

ACC
April 9, 1973
115444

PRODUCT INFORMATI

HAYNES & MORGAN MCPA-AMINE can b
as easily as the 3-leaf stage of small grains . .
is injurious at this stage.

Better select[illegible] and earlie weed control
higher small grain yields than with 2,4 D
More selective in wheat, oats and barley than
Early control (when weeds are young) is most
and prevents weed competition before it ca
toll No grazing restriction[illegible] after application.
Contains 4 pounds of MCPA acid equivalent p
Applied as a water-mixed spray by airplane o
sprayer.

PRODUCT NO. 304

MANUFACTURED BY HAYNES & MORGAN CHEMICALS, INC. – GREAT FALLS, MONTANA – P.O. BOX 3002 – PHONE 406-453-4631