

GRANULAR WEED CONTROL

ACCEPTED
APR 28 1975
EPA Reg. No. 11547-25

Cermet

UNCONDITIONALLY GUARANTEED
WHEN USED AS DIRECTED

NON-SELECTIVE HERBICIDE FOR CONTROLLING A WIDE RANGE OF ANNUAL AND PERENNIAL WEEDS AND GRASSES. RECOMMENDED ONLY FOR NON-CROPLAND AREAS SUCH AS RAILROAD RIGHTS OF WAY AND INDUSTRIAL AREAS.

ACTIVE INGREDIENTS:

Bromacil (5-bromo-3-sec-butyl 6-methyluracil) 4.0%

INERT INGREDIENTS:

96.0%

100.0%

Net Contents: 50 pounds

E.P.A. Reg. No. 11547-25

FOR INDUSTRIAL USE ONLY

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE OTHER PRECAUTIONS ON SIDE PANEL

GRANULAR WEED CONTROL

GRANULAR WEED CONTROL is a non-selective herbicide for controlling a wide range of annual and perennial weeds and grasses. It is recommended only for noncropland areas such as railroad rights-of-way and industrial areas. The application of 200 to 400 pounds of GRANULAR WEED CONTROL per acre usually results in a non-productive condition of the soil for a period of a year or more.

The duration of non-productivity is dependent upon rainfall, type of soil and other conditions.

DIRECTIONS: Apply GRANULAR WEED CONTROL as furnished using a seed spreader, fertilizer spreader, shaker type applicator or any equipment which will distribute this product uniformly over the area to be treated.

To control annual and most perennial weeds and grasses, apply at the rate of 200 to 400 pounds per acre; on smaller plots, 0.5 to 1.0 pound per 100 square feet. Repeat spot treatment may be required when deep rooted perennial weeds are present. To obtain best results, apply GRANULAR WEED CONTROL to the ground where it will be absorbed by the roots. Applications early in the season have been found to give results superior to applications made later in the season. GRANULAR WEED CONTROL may, however, be applied at any time of the year. Maximum effectiveness in arid regions is obtained when applied just prior to the rainy season.

Users should consult State Agricultural Experimental Stations or Extension Service Weed Specialists for recommendations as to use of GRANULAR WEED CONTROL in their particular area.

IMPORTANT: Do not apply on or near valuable woody herbaceous plants or in areas where their roots may be injured because of possible injury to such plants. Thoroughly clean spreading equipment with a suitable chemical cleaner before using for other purposes (or do not use same spreading equipment for other purposes). Do not store near fertilizers, seeds, insecticides or fungicides. Store away from feed and do not contaminate feed and foodstuffs.

Do not contaminate irrigation ditches or waterways for domestic purposes. Do not use on croplands, or land to be used for subsequent cropping. Do not reuse container.

CAUTION:

MAY CAUSE IRRITATION TO SKIN, EYES, NOSE AND THROAT. Avoid contact with skin, eyes and clothing. Avoid breathing dust.

In case of contact, flush with large amounts of water for at least 15 minutes and get medical attention. Remove and wash contaminated clothing before re use.

Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

KEEP OUT OF REACH OF CHILDREN

FOR INDUSTRIAL USE ONLY

AQUATIC WEED CONTROL

A CONTACT, NON-SELECTIVE VEGETATION
CONTROL OF AQUATIC WEEDS

ACTIVE INGREDIENTS:

Diquat dibromide (6, 7-Dihydrodipyrro
(1, 2-a: 2', 1'-c) pyrazinedium dibromide) 1.85%
INERT INGREDIENTS 98.15%

Contains 0.0837 lb. Diquat Cation per gal. as 0.155 lbs. Salt per gal.

This product is a herbicidal chemical, when used as directed, will kill most all forms of plant growth without damage to the soil. Absorption and herbicidal action is usually quite rapid with effects visible in a few days. This product is inactivated on contact with soil.

CAUTION!

KEEP OUT OF REACH OF CHILDREN

Read entire label. Use strictly in accordance with label cautions, warnings and directions and in conformity with Federal and State Regulations. Wash and destroy container when empty. Never re-use.

NET CONTENTS:

EPA REG. NO. 11547-31

UNCONDITIONALLY GUARANTEED
WHEN USED AS DIRECTED

AQUATIC WEED CONTROL

AQUATIC USE DIRECTIONS: AQUATIC WEEDS (infesting Still

Ponds, Lakes, Ditches, Laterals, Canals.)

Water Treatments	Gals./Surface Acre	PPM**
*Algae (Spirogyra and Pithophora spp.)	1/2 to 1 1/2	
Bladderwort (Utricularia spp.)	20 to 40 gal.	1/4 to 1/2
Coontail (Ceratophyllum demersum)	1 1/2 to 2 1/2	

Water Treatments	Gals./Surface Acre
Elooea (Eloodea spp.)	
Najad (Najas spp.)	20 gal.
*Pondweed (Potamogeto spp.)	40 gal.
Watermilfoil (Myriophyllum spp.)	20 to 40 gal.

WATER TREATMENTS: Make application from a boat or other suitable craft either by injecting this product below the water by pouring it directly from the container into the water while moving slowly over the water surface. *Repeat application may be necessary under conditions of heavy infestations or where reinfestation may readily occur from seed, fragmentation or escapes.

**This product is a dry weight. (See dilution chart)

CAUTION: Treatment of aquatic weeds can result in oxygen loss from decomposition of dead weed. This loss can cause fish suffocation. Therefore, to minimize this hazard treat 1/3 to 1/2 of the water area in a single operation and wait at least 10 to 14 days between treatments. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Consult your State Fish and Game Agency before applying this product to public waters. Apply this product only as specified on this label. (Do not apply to muddy water.)

DILUTION CHART

Depth of Water (Ft.)	Gallons to Use per Acre at Indicated PPM Cation (By Wt.)	Depth of Water (Ft.)	Gallons to Use per Acre at Indicated PPM Cation (By Wt.)
1	1 ppm: 7 gal. 2 ppm: 14 gal.	3	21 gal. 4 ppm: 42 gal.
2	3 ppm: 21 gal. 4 ppm: 28 gal.	4	28 gal. 5 ppm: 56 gal.
	5 ppm: 35 gal. 6 ppm: 42 gal.	5	35 gal. 7 ppm: 70 gal.

FOLIAGE SPRAY TREATMENTS -- AMOUNT/ACRE:

(Salvinia rotundifolia)	10 to 15 gal.
Waterlettuce (Pistia stratiotes)	10 to 15 gal.
Water hyacinth (Eichhornia crassipes)	10 to 15 gal.

FOLIAGE SPRAY TREATMENTS should be applied in sufficient water (150-200 gals. per acre) to thoroughly wet foliage. As progresses and vegetation increases in mass, the higher dosages indicated should be used and more uniform distribution is necessary. Conditions may affect the use of this chemical. Consult Agricultural Experiment Station or Extension Service Weed Specialist for recommendations to dosage and time of application, and for control of other weeds. Check with state authorities for necessary permits.

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes and clothing. Concentrate may cause skin irritation. Wear waterproof gloves and face shield or goggles, when handling concentrate. Avoid breathing spray mist. In case of contact, immediately remove clothing and wash thoroughly and get medical attention. For eyes, wash thoroughly with water. Do not store with or near foodstuffs or drinking water.

During application avoid drift to adjacent food and forage plantings that might be damaged or the crops thereof rendered unusable or consumption. Do not use treated water for human or animal consumption, swimming, spraying or overhead and furred within 10 days after treatment. Do not apply to muddy water and do not agitate water excessively with outboard motor equipment during application.

Wash sprayer immediately after application with detergent and water, then rinse with clean water. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.