Hold container at a distance from the area to be treated that will permit even distribution of the dust. Keep lid closed when container is not in use.

CONDITIONS OF SALE: These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions

are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. Chemagro and the Seller offer, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

IMPORTANT: Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using CO-RAL 1% Shaker Can.

RECOMMENDED APPLICATIONS

ANIMAL	PARASITE	REMARKS
Cattle (Dairy & Beef)	Horn Flies Lice	DIRECT APPLICATION: Apply not more than 2 oz.*/animal. Dust evenly into the hair over the head, neck, shoulders, back and tailhead. Repeat as necessary. No interval is required between treatment and use of meat or milk as food.
Swine	Lice .•	DIRECT APPLICATION: Apply not more than 1 oz.*/animal as a uniform coat to the shoulders and back. Repeat as necessary, but not more than once every 10 days. No interval is required between treatment and use of meat as food.
		BEDDING TREATMENT: Apply 2 oz.* uniformly over each 30 square feet of fresh, dry bedding. Repeat as necessary, but not more than once every 10 days.
		DIRECT APPLICATION and BEDDING TREATMENT: In severe infestations, both individual animals and the bedding may be treated as directed above.

*NOTE: 1 oz. of this material equals approximately three level tablespoonfuls.

RESTRICTIONS

For external, insecticidal use on above specified animals only, Avoid contamination of feed, troughs, water, and water utensils. Provide thorough ventilation while dusting.

CAUTION

Harmful if inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling.

Do not contaminate feed or food. Keep out of reach of children.

If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression

To Physician - Alropine sulfate is antidotal, 2-PAM is also antidotal, and may be administered in conjunction with atropine.

Do not reuse empty container. Destroy it by bury-

ing with waste or burning. Stay away from smoke or tumes.

WARRANTY AND LIMITATION OF DAMAGES

Chemagro warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to In the Directions for Use and Conditions of Sale, subject to the risks referred to therein. CHEMAGRO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANT-ABILITY, and no agent of Chemagro is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

CO-RAL is Reg. TM in U.S. Pat Off. by Farbenfabriken Bayer AG.



CO-RAL

Animal Insecticide 1% SHAKER CAN

For control of horn flies and lice on beef and dairy cattle and lice on swine.



-GUARANTEE-

ACTIVE INGREDIENT:
O,O-Diethyl O-(3-chloro-4-methyl-2-oxo-(2H)1-benzopyran-7-yl) phosphorothioate† 1%
INERT INGREDIENTS: 99%
100%
†U.S. Patent No. 2.748.146

EPA Reg. No. 11556-4 AA

CAUTION

KEEP OUT OF REACH OF CHILDREN (See rear panel for caution statements)

NET WEIGHT 2 POUNDS

USE ONLY AS DIRECTED

14

313-245

MANUFACTURED FOR CHEMAGRO

Division of Baychem Corporation Box 4913, Kansas City, Missouri 64120

ACCEPTED

MAR 3 1 1972

UNDER THE FORCEAL INSECTICIDE FOR ELECTROMIC PCISUN REGISTERTO ATTACHED COMMENTS.