


Reason for change to bring label in
line with current requirements.

U.S. IAN

Date of Draft 1/5/67
Supersedes Draft Dated 12/1/66 & 8/15/66
S. J. Size

S. U.S. 12/20

ANIMAL INSECTICIDE



11556-12

FOR CONTROL OF SPECIFIED ANIMAL PARASITES

ACTIVE INGREDIENT
2,3-Dihydro-2H-1,3-dioxolo[4,5-c]pyrrolidine-2-oxo
-2H-1-benzopyran-7-one, monohydro-
chloride / 0.5

INERT INGREDIENTS 99.5
100.0

Reg. No. 2,740,146
Canadian Pat. No. 523,206

USDA Reg. No. 3125-78

(See rear panel for caution statements)

KEEP OUT OF REACH OF CHILDREN

NET WEIGHT POUNDS

Note: Capital letters in chemical no. should be italicized when printed,
not underlined.

CO-RAL 0.5%
-----STOP - READ THIS LABEL BEFORE USE

DIRECTIONS FOR USE

CO-RAL 0.5% is suitable for direct application by any conventional dusting equipment, such as rotary or hand dusters, power dusters, or by spraying from a can.

RECOMMENDED DIRECTIONS

ITEM	INSECT	POWDER CO-RAL 0.5%	REMARKS
SHEEP	Flies (L.) Lice Maggots (wool maggots)	(See Remarks)	<u>HAIR TREATMENT</u> - Apply 1 to 2 ozs. of dust per head. Work dust well into wool over entire body. <u>MACHINE TREATMENT</u> - Machine setting: 1½ to 4 lbs. of dust per minute. Treatment rate: 50 to 100 sheep per minute. Repeat as necessary but not within 15 days of clipping. <u>HOUSING TREATMENT</u> : Dust birds thoroughly, especially under wings and around vent. Supplementary control can be obtained by also treating litter and housing. Repeat as necessary but not more often than weekly.
Poultry	Northern fowl mite Lice	(See Remarks)	<u>HAIR AND HOUSING TREATMENT</u> : Apply specified dosage per 70 sq. ft. of surface area to walls, ceiling, floor, roosts, nests, and adjacent areas, as well as per 20 sq. ft. of litter area. Force dust into all cracks and crevices. <u>DOGS</u> : Dust dogs thoroughly under coat hair over entire body. Repeat as necessary but not more often than weekly. Dogs' bedding should also be dusted to help prevent reinestation.
D.	3	(See Remarks)	

RESTRICTIONS

No more than 1 oz. per animal puppies. Do not treat poultry within 10 days of vaccination or within 10 days after dipping.

Avoid contact with food, water, and wastes of man or animals.

Remove dust from animal while drying.

CAUTION

Do not allow treatment to be on...
any prolonged inhalation of dust. Wash hands frequently.

CO-RAL 0.5% Dust

Caution
(Continued)

Do not take food or drink. Keep out of reach of children.

If contact occurs, obtain professional medical aid. Prolonged exposure will result in chronic lung depression.

Toxicity of the sulfate is unknown. 2-PAM is also antidotal and may be administered in conjunction with Acropine.

CO-RAL is U.S. and Canadian Pat. Offs. by Farbenfabriken Bayer A.G.,
Ciba Corporation licensee.

MANUFACTURED BY

CIBA CORPORATION

KANSAS CITY, MO.