



# GO-RAL

Brand of coumaphos

## ANIMAL INSECTICIDE

### 1% BULK DUST

CONTROL OF HORN FLIES & LICE ON BEEF & DAIRY CATTLE — REDUCTION OF FACE FLIES ON BEEF AND DAIRY CATTLE — CONTROL OF LICE ON SWINE

USE IN DUST BAG, SHAKER CAN, OR MECHANICAL DUST APPLICATOR

NET WEIGHT 25 POUNDS

BATCH NO.



**CAUTION**  
KEEP OUT OF REACH OF  
CHILDREN

GUARANTEE


ACTIVE INGREDIENTS	
1,2-Dichloro-3-(4-chlorophenyl)-2-methyl-2-propanone	1.0%
2,2,4,4-Tetrachloro-1,2-diphenylethane	1.0%
INERT INGREDIENTS	99.0%
	100.0%

P.A. Reg. No. 1-156-14

USE ONLY AS DIRECTED



Chemagro



ACCEPTED

5-29-73

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC INSECTICIDES  
REGISTERED UNDER NO. 11556-14

DIRECTIONS FOR USE

**IMPORTANT:** Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using CO-RAL 1% Bulk Dust.

**CONDITIONS OF SALE:** These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. Chemistry and the Seller offer, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

CO-RAL 1% Bulk Dust may be used to fill cattle dust bags or it may be applied directly to cattle, swine, or swine bedding by use of a shaker can or a suitable mechanical dust applicator. When used as recommended, CO-RAL 1% Dust controls horn flies and lice on beef and dairy cattle and lice on swine. CO-RAL 1% Dust will also reduce face fly populations on cattle, however, as face flies are able to travel over an area of several miles, best reduction of face fly populations is achieved when CO-RAL 1% Dust is used on an area-wide basis.

RECOMMENDED APPLICATIONS

ANIMAL	PARASITES	REMARKS
Beef and Dairy Cattle	Horn Flies Lice Face Flies	<b>DUST BAG:</b> Place contents of this package in any commercially available dust bag. Suspend bags in areas frequented by cattle or in gateways or lanes through which the animals pass daily for water, feed, or minerals. Bags may also be suspended in loafing sheds or in front of protected mineral feeders. For lactating dairy cows, bags may be suspended in the exit through which cows leave the milking barn. In all cases, the bags should be adjusted so that the bottom of the bag will hang four to six inches below the topline of the cattle. For reduction of face flies, the bags must be located so the animals will be forced to use them daily and hung at a height to insure that the faces of the cattle will be dusted.
	Horn Flies	<b>DIRECT APPLICATION:</b> Dust may be applied by use of a shaker can or by using a suitable mechanical dust applicator. Animals should be thoroughly dusted during a treatment and the level of dusting should be repeated for each animal per treatment. Repeat at recommended intervals.
		<b>DIRECT APPLICATION:</b> Dust may be applied by use of a shaker can or by using a suitable mechanical dust applicator. Animals should be thoroughly dusted during a treatment and the level of dusting should be repeated for each animal per treatment. Repeat at recommended intervals.

