

PM03

11556-14

12/13/99

1 of 7

Reason to Issue: To remove mechanical duster application from swine uses.

Date: 11/24/99
Supersedes: 10/3/97
Page 1 of 7


(Front Panel)

Co-Ral®

(coumaphos)

ANIMAL INSECTICIDE

1% BULK DUST



Control of Horn Flies and Lice on Beef and Dairy Cattle, Reduction of Face Flies on Beef and Dairy Cattle, Control of Lice on Swine

Use in Dust Bag or Shaker Can

Active Ingredient:

0,0-Diethyl 0-(3-chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl) phosphorothioate

1%

Inert Ingredients:

99%
100%

EPA Reg. No. 11556-14

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel for Statements of Practical Treatment and Other Precautionary Statements

NET WEIGHT 12.5 POUNDS

Manufactured for
Bayer Corporation, Agriculture Division, Animal Health
Shawnee Mission, Kansas 66201

Reason to Issue: To remove mechanical duster application from swine uses.

Date: 11/24/99
Supersedes: 10/3/97
Page 2 of 7

(Side Panel)

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dusts. Avoid contact with eyes, skin, and clothing.

STATEMENTS OF PRACTICAL TREATMENT

- If swallowed - Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.
- If inhaled indicated - Remove victim to fresh air. Apply artificial respiration if
Get medical attention if victim displays signs of poisoning.
- If on skin - Remove contaminated clothing and wash affected areas with soap and water.
- If in eyes - Flush eyes with plenty of water. Call a physician immediately.

To Physician: Prolonged exposure will result in cholinesterase depression. Atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Reason to Issue: To remove mechanical duster application from swine uses.

Date: 11/24/99
Supersedes: 10/3/97
Page 3 of 7

(Side Panel)

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, fish and aquatic invertebrates. Coumaphos washed off of wading treated livestock may be hazardous to aquatic organisms. Do not contaminate water when disposing of equipment washwater or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry place.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty the container by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

Reason to Issue: To remove mechanical duster application from swine uses.

487
Date: 11/24/99
Supersedes: 10/3/97
Page 4 of 7

(Back Panel)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

) Co-Ral 1% Bulk Dust may be used to fill cattle dust bags or it may be applied directly to cattle, swine or swine bedding by use of a shaker can. When used as recommended, Co-Ral 1% Dust controls horn flies and lice on beef and dairy cattle and lice on swine. Co-Ral 1% Dust will also reduce face fly populations on cattle; however, as face flies are able to travel over an area of several miles, best reduction of face fly populations is achieved when Co-Ral 1% Dust is used on an area-wide bases.

Premise Precautions: Do not treat areas such as feed bunks, mangers, or troughs where livestock feed or drink. Do not contaminate food, feedstuffs, food or feed handling equipment or milk or meat handling equipment.

INSTALLATION OF DUST BAGS:

) Suspend the bags by inserting rope, chain or wire through the grommets at the top of the bag and attaching to an overhead structure or by hooking the grommets into hooks fastened to a stationary overhead 2 x 4 board at the heights recommended for louse and horn fly or face fly control. In areas of frequent or heavy rainfall Co-Ral 1% Cattle Duster should be protected from the weather.

Install as many Co-Ral 1% Cattle Dusters as necessary to lanes or other openings to assure proper dusting of the animals. After the bags are in place, pull the rip cord to allow the insecticide to fall into the bottom of the bag. When dust bags are provided on a free-choice basis, allow one 12.5-pound Co-Ral 1% Duster per 25 animals.

Reason to Issue: To remove mechanical duster application from swine uses.

Date: 11/24/99
Supersedes: 10/3/97
Page 5 of 7

(Back Panel)

RECOMMENDED APPLICATIONS

ANIMAL	PARASITE	REMARKS
Beef and Dairy Cattle	Horn Flies Lice Face Flies	<u>DUST BAG:</u> Place contents of this package in any commercially available dust bag. Suspend bags in areas frequented by cattle or in gateways or lanes through which the animals pass daily for water, feed or minerals. Bags may also be suspended in loafing sheds or in front of protected mineral feeders. For lactating dairy cows bags may be suspended in the exit through which cows leave the milking barn. In all cases the bags should be adjusted so that the bottom of the bag will hang four to six inches below the topline of the cattle. <u>For reduction of face flies:</u> (1) The bottom of the bags should be 18 to 24 inches from the ground to insure that the faces of the cattle will be dusted. (2) The bags must be located so the animals will be forced to use them daily.
	Horn Flies Lice	<u>DIRECT APPLICATION:</u> Co-Ral 1% Dust may be applied by use of a shaker can. Animals should be thoroughly dusted using not more than 2 oz.* (6 level tablespoonfuls) of dust per animal per treatment. Treat no more than 12 times per year. Do not make applications less than 10 days apart.

(Continued)

Reason to Issue: To remove mechanical duster application from swine uses.

Date: 11/24/99
Supersedes: 10/3/97
Page 6 of 7

(Back Panel)

RECOMMENDED APPLICATIONS

ANIMAL	PARASITE	REMARKS
Swine	Lice	<p><u>DIRECT APPLICATION</u>: Apply not more than 1 oz.* (3 level tablespoonfuls) per head as a uniform coat to the head, shoulders and back by use of a shaker can. Treat no more than 6 times per year. Do not make applications less than 10 days apart.</p> <p><u>BEDDING TREATMENT</u>: Apply 2 oz.* (6 level tablespoonfuls) uniformly over 30 square feet of fresh, dry bedding by use of shaker can. Treat no more than 6 times per year. Do not make applications less than 10 days apart.</p> <p><u>DIRECT APPLICATION AND BEDDING TREATMENT</u>: In severe infestations both individual animals and the bedding may be treated as directed above.</p>
No interval is required between treatment and slaughter or between treatment and use of milk as feed.		

*Note: 1 oz of this material equals approximately three level tablespoonfuls.

USE RESTRICTIONS

For external insecticidal use only on above-specified animals. Do not contaminate feed, troughs, water or water utensils. Provide thorough ventilation while dusting. Do not apply to sick, stressed or convalescent animals.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Reason to Issue: To remove mechanical duster application from swine uses.

Date: 11/24/99
Supersedes: 10/3/97
Page 7 of 7

(Back Panel)

PROTECTIVE CLOTHING

Applicators and other handlers must wear long sleeve shirt, long pants, chemical-resistant gloves such as barrier laminate or butyl rubber \geq 14 mils, shoes plus socks.

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment. If no such instructions for washables, use detergent and hot water. Keep and wash personal protective equipment separately from other laundry.

LIMITED WARRANTY AND
LIMITATION OF DAMAGES

Bayer Corporation, Agriculture Division, Animal Health warrants that this material conforms to the chemical description on the label. BAYER CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, and no agent of Miles Inc. is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

Co-Ral® is a Reg. TM of Bayer AG.