




DISINFECTANT SECTICIDE



REF ID: A6494 1980

RESTD SERIALIZED

- RESTRICTIONS
- Do not apply as a spray at rates above 1½ ounce of **CD-BAL** Emulsion per acre of water-lactating dairy cattle or non-lactating dairy cattle within 14 days of their parturition. If freshening spray **ED-400** within the same day after spraying **CD-BAL**, wait 14 days before feeding human food or balance of 14-day interval.
- Do not apply in case of convalescence or stressed livestock or to animals less than 3 months old.
- Do not spray animals for 10 days before or after shipping or weaning, or after exposure to contagious and infectious diseases.
- Do not spray in a confined, non-ventilated area.
- Do not apply in conjunction with oral drenches or other internal medications such as phenothiazine nor with tranquilizers, synthetic pyrethroids or their synergists or with other organic phosphates.



CUTTER ANIMAL HEALTH

CODE 132-83



CODE 132.8

ANIMAL	PARASITE	FL OZ CO. RA.	REMARKS
	HORSES	2	SPRAY TREATMENTS All animals 100 gallons of water + 1/2 oz. carbamate per acre. See directions for use.
	CATTLE	2	SPRAY TREATMENT(S) Apply 100 gallons of water + 1/2 oz. carbamate + high pressure spray to cattle with a high pressure sprayer. Repeat as necessary.
Beef and Dairy Cattle	Screw Worms	10	SPRAY TREATMENT Apply 1/2 oz. carbamate per acre + 100 gallons of water + high pressure spray to cattle with a high pressure sprayer. Repeat as necessary. See directions for use for cattle spray treatments.
	GOATS	10	SPRAY TREATMENT(S) All animals 100 gallons of water + 1/2 oz. carbamate per acre. See directions for use.
Beef and Dairy Cattle	Horn Flies	10	CATTLE BACKRUBBERS V. Shaved down to skin of all cattle and 1/2 inch from shoulder blades. The skin of the shoulder blades is shaved with a straight razor while the rest of the body is shaved with a fine blade. The shaved portion of the body is painted with Resinate base spray as directed. The paint must adhere well to the skin. The shoulder blades are then rubbed with the backrubber between the shoulder blades.
	Face Flies	10	SPRAY TREATMENT(S) Apply 100 gallons of water + 1/2 oz. carbamate + high pressure spray to cattle with a high pressure sprayer. Repeat as necessary.
Beef and Dairy Cattle	Horn Flies	10	SPRAY TREATMENT(S) Apply 100 gallons of water + 1/2 oz. carbamate + high pressure spray to cattle with a high pressure sprayer. Repeat as necessary.
Horses	Horn Flies	10	SPRAY TREATMENT(S) Apply 100 gallons of water + 1/2 oz. carbamate + high pressure spray to cattle with a high pressure sprayer. Repeat as necessary.
Beef and Dairy Cattle	Breast Worms	10	SPRAY TREATMENT(S) Apply 100 gallons of water + 1/2 oz. carbamate + high pressure spray to cattle with a high pressure sprayer. Repeat as necessary.
Cattle	Leptospirosis	2	

Key - No 11556-23
Phase 105-3
G. W. Knobell (2)

140

10 King St. West - 23
Floor 2 - 13
Furnished




LTA Ref. No 1153623
Page 3 of 3
Top of Con⁶

ATTENTION:

**CONTENTS MUST BE DILUTED
BEFORE USE**

142

BEST DOCUMENT AVAILABLE





INGREDIENTS.

Date . 11.6°
 83.4°
 ... 50°
 100.00%

GUARANTEE:
10-(3-chloro-4-enzopyran-7-yl)
cate per gallon.
No. 11556-KS-1

WARNING
OF CHILDREN
from freezing.
ing statement(s)

IMPORTANT.
affet, including
on of Damages
before usin g.

**S DIRECTED
OZ (10%)
ANIMAL HEALTH**
Laboratories, Inc.
5500 N. 44th Street
Milwaukee, Wisconsin 53214

CUTTER ANIMAL HEALTH

The logo for Cutter, featuring the word "CUTTER" in a bold, sans-serif font inside a dark, irregular hexagonal border.

CUTTER ANIMAL HEALTH
Division of Cutter Laboratories, Inc.
750 Park Avenue • New York, N.Y. 10021

Digitized by Google

CODE 132-8

ACCEPTED
CODE 132-13

RESTRICTIONS 4 1980

Do not apply as a spray at rates above 1½ times of CD-HAL Emulsion live stock insecticide per 4 quarts of water including dairy cattle or lactating dairy cattle in 1½ days after freshening sheep. Do not apply after spraying the animal with human formulation of 14-day insecticide.

Human foot for balance of 14-day interval.
Do not apply to a cow convalescent from a stress or lesion or to an animal less than 3 months old.
Do not spray animals for 10 days before or after shearing or weaning, or after exposure to contagious and infectious diseases.

Do not spray in a confined, non-ventilated area.
Do not apply in conjunction with oral drenches or other internal medications such as phenothiazine nor with natural synthetic pyrethrins or their synergists or with other organic phosphates.

or with other organic phosphate.

卷之三

CUTTER

Digitized by srujanika@gmail.com

CUTTER ANIMAL HEALTH

CUTTER ANIMAL HEALTH
BAYER TOUGHON CUTTER Laboratories, Inc.


Shawnee, Kansas 66201 U.S.A.

• • • •

ANIMAL	PARASITE	FL. OZ. CO. EACH	REMARKS
Dairy Cattle	Horn Flies	2	SPRAY TREATMENTS: Apply 4 gallons of water to a cow per head. Repeat as necessary.
Dairy Cattle	Lice	5	SPRAY TREATMENT(S): Apply 4 gallons of water at high pressure spray to coat the entire body of the animal. Repeat as necessary.
Dairy Cattle	Worms & Eggs	10	SPRAY TREATMENT(S): Apply 4 gallons of water at high pressure spray to coat the entire body of the animal. See Directions For Use of Spraying Equipment.
Dairy Cattle	Gloes	15	CATTLE BACKRUBBERS: Mix 1/2 oz. orange oil, 1/2 oz. 2% sulfur, 1/2 oz. diatomaceous earth, 1/2 oz. Salvarsan, the third portion of the barbed wire mesh, 1/2 oz. zinc oxide and 1/2 oz. linseed oil. Where Diptes and/or ticks are found regularly, add 1/2 oz. pyrethrum powder to backrubber as needed. Note: For the best results, apply oil to the rubber. The rubber should be constructed so as to fit over the animal to rub oil into the skin. Reapply between treatments and as often as necessary.
Dairy Cattle	Horn Flies Face Flies	9	SPRAY TREATMENT(S): Apply 4 gallons of water to a cow per head. Repeat as necessary.
Dairy Cattle	Lice	14	SPRAY TREATMENT(S): Apply 4 gallons of water to a cow per head. Repeat as necessary.
Diseases Insects and Parasites	Horn Flies	2	SPRAY TREATMENT(S): Apply 4 gallons of water to a cow per head. Repeat as necessary.
Diseases Insects and Parasites	Lice	5	SPRAY TREATMENT(S): Apply 4 gallons of water to a cow per head. Repeat as necessary.
Diseases Insects and Parasites	TEAS	6	SPRAY TREATMENT(S): Apply 4 gallons of water to a cow per head. Repeat as necessary.
Diseases Insects and Parasites	Scabies & J. S.	10	SPRAY TREATMENT(S): Apply 4 gallons of water to a cow per head. Repeat as necessary.
Diseases Insects and Parasites	Lice	3	SPRAY TREATMENT(S): Apply 4 gallons of water to a cow per head. Repeat as necessary.

Price 100³

141



111111
12 7 1979
10 1 1980