



11556-34

04/14/1977

1/2

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

APR 14 1977

Bayvet Division
Cutter Laboratories
Box 390
Shawnee, Kansas 66201

Gentlemen:

Subject: Tiguvon Swine Insecticide Pour-On
EPA Reg. No. 11556-34
Your application of March 29, 1977

Thank you for sending us copies of the finished labeling for the subject product. They have been incorporated into the product file for "TIGUVON Swine Insecticide Pour-On", EPA Reg. No. 11556-34.

Sincerely,

Timothy A. Gardner
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (WH-567)

Keep labeling accepted October 19, 1976

CODE 137-05



TIGUVON®

Brand of fenthion

SWINE INSECTICIDE POUR-ON

FOR CONTROL OF LICE ON SWINE

ACTIVE INGREDIENT:

fenthion (O, O-dimethyl O-[3-methyl-4-(methylthio) phenyl] phosphorothioate)†	3%
INERT INGREDIENTS:	97%
	100%

†U.S. Patent No. 3,042,703
 Canadian Patent No. 631,472
 EPA Reg. No. 11556-34
 EPA Est. No. 11556-KS-1

WARNING

KEEP OUT OF REACH OF CHILDREN

USE ONLY AS DIRECTED

(See back panel for warning statements)

NET CONTENTS: 1 QUART

TAKE TIME



OBSERVE LABEL DIRECTIONS

DIRECTIONS FOR USE

TIGUVON (Brand of fenthion) Pour-On Swine Insecticide is a ready-to-use preparation that requires no further dilution. A single application placed on the backline of swine gives good initial kill of lice.

AMOUNT TO USE FOR LOUSE CONTROL ON SWINE: Apply TIGUVON Pour-On at the rate of 1/2 fluid ounce (15 cc) per 100 pounds of body weight. Measure the correct amount to use in relation to the animal's weight. All swine in the herd, including gestating and lactating sows, should be treated at the same time to reduce sources of reinfestation. Treatment of lactating sows will result in an effective reduction of lice on nursing pigs.

HOW TO APPLY: Pour the correct amount of solution uniformly along the animal's back, starting just behind the ears and continuing to the rump. TIGUVON Pour-On may be dipped from any clean bucket or pail. Do not subsequently use dip containers in any connection with feed, food, or drinking water.

CONTRAINDICATIONS AND RESTRICTIONS

Do not treat sick, convalescent or stressed animals.

→ **WARNING: DO NOT SLAUGHTER SWINE WITHIN 14 DAYS OF TREATMENT.** ←

TIGUVON (Brand of fenthion) is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

WARNING

May be poisonous if swallowed. Harmful if absorbed through the skin. Do not get in eyes or on skin. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

Do not contaminate feed or food. Keep out of reach of children.

This material contains a cholinesterase inhibitor. If poisoning should occur, obtain prompt medical aid.

To Physician and Veterinarian - Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

CONTAINER DISPOSAL

DO NOT REUSE THE CONTAINER - Completely empty the contents. Rinse empty container by filling with water and adding 2 tablespoonsful of household lye. Bury rinse solution and any unused chemical at least 18 inches deep in an isolated location away from water supplies. Punch holes in the top and bottom of the can, crush and bury as above.

Do not use container in any connection with feed, food or drinking water.

PROTECT WILDLIFE

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

TIGUVON is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen.

0137050

CUTTER ANIMAL HEALTH
BAYVET Division Cutter Laboratories, Inc.
 Shawnee, Kansas 66201, U.S.A.

CUTTER ANIMAL HEALTH
BAYVET Division Cutter Laboratories, Inc.
 Shawnee, Kansas 66201, U.S.A.

2/2