



CODE 136-65



LYSOFFTM POUR-ON FOR LICE FOR BEEF AND NON-LACTATING DAIRY CATTLE

CONTAINER DISPOSAL
DO NOT REUSE THE CONTAINER - Completely empty the container. Rinse empty container by Non-Hazardous Waste Disposal. Do not use used chemical at least 18 inches deep in an hole in the soil. Do not use in water supply, Pond and bury as above. Do not use in connection with feed, food or drinking water.

PROTECT WILDLIFE
This product is toxic to fish, birds, and other animals. Do not use near water, streams or ponds. Do not contaminate water, streams or ponds. Do not use in water supply. Do not use in water supply. Do not use in water supply. Do not use in water supply.

LIMITED WARRANTY AND LIMITATION OF DAMAGES
BAYVET warrants that this material conforms to the chemical description on the label. BAYVET does not warrant that the material will be effective against lice. BAYVET does not warrant that the material will be effective against lice. BAYVET does not warrant that the material will be effective against lice.

ACTIVE INGREDIENTS:
Fenitron (O,O-Dimethyl O-t-(methylthio)-p-1991) 1.5%
piperonyl butoxide 5.7%
petroleum distillate 92.8%
INERT INGREDIENTS:
100.00%

U.S. Patent No. 3,042,703
EPA Reg. No. 11556-48
EPA Est. No. 509-65-1

WARNING
KEEP OUT OF REACH OF CHILDREN

See side panel for warning statements and bottom of can for batch No.
NET CONTENTS: 64 fl. oz. (1/2 gallon)
Manufactured by

GUTTER ANIMAL HEALTH LABORATORIES
A Division of Bayer Corporation
SHELDON, KENTUCKY 40381

All printing blue on white except as noted.
Actual size.

CONTRAINDICATIONS AND RESTRICTIONS

Do not treat lactating dairy cattle, calves less than 6 months old, sick, convalescent or stressed animals. Do not treat non-lactating dairy cattle within 28 days of freshening. Freshening should occur as normal. Do not use on animals that are pregnant or nursing. Do not use on animals that are pregnant or nursing.

WARNING: DO NOT SLAUGHTER CATTLE WITHIN 14 DAYS OF SECOND APPLICATION. DO NOT SLAUGHTER CATTLE WITHIN 14 DAYS OF SECOND APPLICATION. DO NOT SLAUGHTER CATTLE WITHIN 14 DAYS OF SECOND APPLICATION.

WARNING

Harmful or fatal if swallowed; inhaled or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water. Wash contaminated clothing immediately. Do not eat, drink or use tobacco. Do not contaminate feed or food. Keep out of reach of children. Do not use as a molluscicide, insecticide or acaricide. Do not use as a molluscicide, insecticide or acaricide.

DIRECTIONS FOR USE

IMPORTANT: Read the entire label before use. This product is a restricted use pesticide. It is not to be used on animals that are pregnant or nursing. It is not to be used on animals that are pregnant or nursing. It is not to be used on animals that are pregnant or nursing.

Amount of LYSOFF Pour-On Desired	Mix (Step A above)	Amount of Water to be Added to LYSOFF Pour-On	Final Mixture (Step B above)	Additional Water to be Added to Final Mixture
1 1/2 quarts	1 1/2 pint	1 1/2 quart	3 quarts	7 pints
2 1/2 quarts	2 1/2 pint	2 1/2 quart	5 quarts	11 pints
3 1/2 quarts	3 1/2 pint	3 1/2 quart	7 quarts	15 pints
4 1/2 quarts	4 1/2 pint	4 1/2 quart	9 quarts	19 pints
5 1/2 quarts	5 1/2 pint	5 1/2 quart	11 quarts	23 pints
6 1/2 quarts	6 1/2 pint	6 1/2 quart	13 quarts	27 pints
7 1/2 quarts	7 1/2 pint	7 1/2 quart	15 quarts	31 pints
8 1/2 quarts	8 1/2 pint	8 1/2 quart	17 quarts	35 pints
9 1/2 quarts	9 1/2 pint	9 1/2 quart	19 quarts	39 pints
10 1/2 quarts	10 1/2 pint	10 1/2 quart	21 quarts	43 pints

Manufactured by

GUTTER ANIMAL HEALTH LABORATORIES
A Division of Bayer Corporation
SHELDON, KENTUCKY 40381

11556-48

12/17/1975

1/1