0	2006	•	A	С



CEPTED MAR 1 0 2006

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under EPA Reg. No. 1155-6-116

Once-a-month topical flea treatment for cats and kittens 8 weeks and older and 9 lbs. and under.

READ THE ENTIRE LABEL BEFORE EACH USE.

- For the Prevention and Treatment of Fiea infestations on Cats
 - Available only through licensed practicing veterinarians
 Kills 98-100% of the fleas on cats within 12 hours
- Active Ingredient

One treatment prevents further flea infestation for up to four weeks
 Kills fleas before they lay equs

% By Weight

..<u>90.9%</u> 100.0%

Imidacloprid; I-[(6-Chloro-3-pyridinyl) methyl]-N-nitro-2- imidazolidinimine... Other Ingredients.....

Total

Jo- 116

KEEP OUT OF REACH OF CHILDREN

See below for First Aid and Precautionary Statements.

Precautionary Statements

Hazards to Humans: Causes eye irritation. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

Hazards to Domestic Animals

For external use only. Do not use on kittens under eight weeks of age. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. If signs persist, or become more severe, consult a veterinarian immediately. If your animal is on medication, consult your veterinarian before using this or any other product. For consumer questions call 1-800-255-6826. For medical emergencies call 1-877-258-2280.

First Aid

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.

To Physician: Treat the patient symptomatically.


Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

How to Apply

1. Use only on cats. Do not use on other animals.

2. Remove one applicator tube from the package.



🐏 Bayer HealthCare

Manutactured For Bayer HealthCare LLC, Animal Health Division, Shawnee Mission, Kansas 66201 USA

Bayer, the Bayer Cross and Advantage are trademarks of Bayer

Advantage is protected by one or more of the following U.S. patents: 6,838 482, 6,613,783, 6,495 573, 6,359,374 6,232,328, 5,428,032, 5,298,507, 5,204,380, 4,742,060

Hold applicator tube in an upright position. Pull cap off tube.
 Turn the cap around and place other end of cap back on tube.



6. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube twice to expel the entire contents of the tube directly on the skin. Do not get this product in your pet's eyes or mouth. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product.

7. Discard empty tube as described in Storage and Disposal.

The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with Advantage rapidly kills fleas and may reduce the incidence of this condition.

Advantage kills 98-100% of the fleas on pets within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting for up to four (4) weeks. Pre-existing pupe in the environment may continue to emerge for six (6) weeks or longer depending upon the climatic conditions.

Larval flea stages in the pet's surroundings are killed following contact with an Advantage treated pet.

Advantage remains efficacious following a shampoo treatment, swimming or after exposure to rain or sunlight.

Monthly treatments are required for optimal control and prevention of fleas.

If re-treatment becomes necessary earlier than four weeks, do not re-treat more than once weekly.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry place.

Disposal: If empty: Do not reuse container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-877-258-2280 for disposal instructions. Never place unused product down any indoor or outdoor drain.

Limited Warranty and Limitation of Damages

Bayer HealthCare LLC. Animal Health Division warrants that this material conforms to the chemical description on the label. BAYER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY, and no agent of Bayer is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

C 2005 Baver HealthCare LLC

01405654	00837869-032699/01791668-053699, R.0 EPA Est. 11556-DEU-1
Made in Germany	EPA Reg. No. 11556-116
To receive monthly email app	lication reminders, go to www.nofleas.com.




advant

01405735

For Cats and Kittens

8 Weeks and Older and 9 lbs and Under.





Once-A-Month **Topical Flea Treatment** for Cats and Kittens. 8 Weeks and Older and 9 lbs and Under.

Available only through licensed practicing veterinarians Kills 98-100% of fleas within 12 hours Provents reinfestation for up to 4 weeks Convenient, easy to apply

advantage

85 x 20 x 150

Once-a-month lopical fies treatment for cats and kittens 8 weeks and older and 8 lbs and under

For the Prevention and Treatment of Flea Infestations on Cats

- Available only through licensed practicing vaterimarians
- Killis 98-100% of the flees on cats within 12 hours
- · One treatment prevents further fles infestation for up to four weeks

· Kills fleas before they by eous

Active Ingradient	% By Weight
Imidacioprid; 1-((6-Chioro-3-pyridinyi) methyi)-N-nitro-2-Imidazolidinimine	
Other Ingradients	
Total	100.0%

KEEP OUT OF REACH OF CHILDREN WARNING

See below for First Aid and Precautionary Statements.

For Directions for Use and other labeling, see supplemental labeling inside.

Precautionary Statements

Hazands to Humana: Cause eye inflation, Hamilul if swallowed, De not get in eyes or an clothing. Avoid contact with side. Wash hands thoroughly with some and warm water after handling. Keep out of reach of children. Do not contaminate feed or fond.

Hazards to Domestic Animals; Fer externel use only, Do not use on kittane under eight weeks of age. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or maning animals, individual sensitivities, while rare, may occur after using AHY pesticide product for peta. # signs persist, or become more severe, consult a velocitarian immediately. If your animal is on medication, consult your veterimentan before using this or any other product,

For consumer questions call 1-800-255-8828. For medical emergencies call 1-877-258-2280.

First Aid

Here the product container or label with you when calling a poleon control center or ductor, or going for treatment,

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact fenses. If present, after the first 5 minutes, then continue rinsing eye. Call a polson combol center or doctor for treatment advice.

If evaluated: Call poison control center or doctor immediately for treatment edvice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the polson control center or doctor. Do not give anything by mouth to an unconscious person,

If an akin or glothing: Take off contaminated clothing, Rinse aids immediately with plonty of water for 15-20 minutes, Call poleon control center or doctor for treatment advice.

To Physician: Trout the petient symptomatically, EPA Est. 11998-DEL-1 EPA Reg. No. 11558-118

(Bayer

Six 0.014 fl oz (0.4 ml) Tubes

01701068-053609, FL8



Mars/lackared Por Bayer HealthCare LLC, Made In Germany Animal Health Division, www.nofees.com I © 2005 Bayer HealthCare LLC Animal Health Division, nes Mission, Kanasa 85201 LISA

Bayer, the Bayer Cross and Advantage are tradomeries of Bayer

Lot No.:

N

advantage 9 9.1% imidacloprid 0.014 floz (0.4 ml) EPA Reg. No.11556-116 WARNING Keep out of reach of children Read The Entire Label Before Use 01223505 LOT NO. 1

 \mathcal{C}_{λ}