UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460



NOV 0 4 2008

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Douglas Spilker, Ph.D Bayer HealthCare LLC, Animal Health Division P.O. Box 390 Shawnee Mission, KS 66201-0390

Dear Dr. Spilker:

Subject:

Amendment of label to revise according to EPA's letter of 3/4/08, PR Notice

2007-4 and other minor changes

Advantage® 9 & 10 Topical Solutions

EPA Registration No. 11556-116 & 11556-117 Your Email Re-submissions were received 10/29/08

The labeling referred to above submitted in connection with the Federal Insecticide, Fungicide and Rodenticide, as amended are acceptable.

A stamped copy of the labeling is enclosed for your records. Please submit one final printed copy of the labeling before releasing the product for shipment. If you have any questions regarding this label, please contact Autumn Metzger at (703) 305-5314.

Sincerely.

Venus Eagle

Product Manager 01

Insecticide-Rodenticide Branch

lemes Pagle

Registration Division (7505P)

03/04/2008, and make other minor changes. Supersedes: 02/04/08

Date: 09/05/08

Under the Federal Insecticide, Fungicide,

Note to Reviewer: [Brackets and parentheses indicate alternate language]

Advantage® 9 Topical Solution

Once-A-Month Topical Flea Treatment for Cats and Kittens 8 Weeks and Older and 9 lbs. and Under.

READ THE ENTIRE LABEL BEFORE EACH USE

For the Prevention and Treatment of Flea Infestations on Cats

- One treatment prevents further flea infestation for up to four (4) weeks
- Kills fleas on cats within 12 hours
- Kills fleas before they lay eggs
- Convenient, easy to apply
- Waterproof
- Remains effective after exposure to rain or sunlight
- Fragrance free

	o 11556-DFIL1
Total	100.0%
Other Ingredients	90.9%
Active Ingredient Imidacloprid	. 9.1%
	% By Weight

KEEP OUT OF REACH OF CHILDREN

[For Directions For Use, and Storage and Disposal (instructions), see supplemental labeling inside]

WARNING

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS: Causes eye irritation. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

Page 1 of 5

3/6

Reason To Issue:

Revise according to EPA letter of

03/04/2008, and make other minor changes.

Date: 09/05/08

Supersedes: 02/04/08

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not use on kittens under eight weeks of age. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. If signs persist, or become more severe, consult a veterinarian immediately. If your animal is on medication, consult your veterinarian before using this or any other product. For consumer questions call 1-800-255-6826. For medical emergencies call 1-800-422-9874.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

To Physician: Treat the patient symptomatically.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOW TO APPLY

- 1. Use only on cats. Do not use on other animals.
- 2. Remove one applicator tube from the package.

Reason To Issue:

Revise according to EPA letter of

03/04/2008, and make other minor changes.

Date: 09/05/08 Supersedes: 02/04/08

3. Hold applicator tube in an upright position. Pull cap off tube.

[Visuals Depicting How to Open Applicator Tube]

- 4. Turn the cap around and place other end of cap back on tube.
- 5. Twist cap to break seal, then remove cap from tube.

[Visuals Depicting Application To Animal]

- 6. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents of the tube directly on the skin. Do not get this product in your pet's eyes or mouth. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product.
- 7. Discard empty tube as described in Storage and Disposal.
- 8. In case of severe flea infestation, retreatment may be necessary earlier than four weeks. Do not retreat more often than once every seven days. After flea control is attained, return to a monthly retreatment schedule.

GENERAL INFORMATION

The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of cats with Advantage kills fleas and may reduce the incidence of this condition.

Advantage kills fleas on cats within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting for up to four (4) weeks. Pre-existing pupae in the environment may continue to emerge for six (6) weeks or longer depending upon the climatic conditions.

Larval flea stages in the pet's surroundings are killed following contact with an Advantage treated pet.

5/6

Reason To Issue:

Revise according to EPA letter of

03/04/2008, and make other minor changes.

Date: 09/05/08

Supersedes: 02/04/08

Advantage is waterproof and remains effective following a shampoo treatment, swimming or after exposure to rain or sunlight.

Monthly treatments are required for optimal control and prevention of fleas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry place.

Disposal: Nonrefillable container. If empty: Do not reuse or refill this container. Place in

trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-422-9874 for disposal

instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND LIMITATION OF DAMAGES

Bayer HealthCare LLC, Animal Health Division warrants that this material conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY, and no agent of Bayer is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

Four [or Six] 0.014 fl. oz. (0.4 mL) Tubes

Manufactured For

Bayer HealthCare LLC
Animal Health Division
P.O. BOX 390
Shawnee Mission, Kansas 66201 USA

Made in Germany

Reason To Issue:

Revise according to EPA letter of

03/04/2008, and make other minor changes.

Date: 09/05/08

Supersedes: 02/04/08

[Label on Individual Tube]

Advantage® 9

9.1% Imidacloprid

- 0.014 fl. oz. (0.4 mL)

EPA Reg. No. 11556-116

Warning

Keep Out of Reach of Children

Read The Entire Label Before Use

BAYER

Lot No. 0000000