

PLEASE NOTE

This image contains more than one label approved for this product on this date.

Reason To Issue: Revised according to PR Notice 2001-1.

Date: 01/08/01

Supersedes: 9/13/99

(Front Panel)

Advantage® 10
(imidacloprid) Topical Solution


Once-A-Month Topical Flea Treatment For Dogs and
Puppies 7 Weeks and Older and 10 lbs and Under

READ THE ENTIRE LABEL BEFORE EACH USE

For the Prevention and Treatment of Flea Infestations on Dogs

- Available Only Through Licensed Practicing Veterinarians
- Kills 98-100% of the Fleas on Dogs Within 12 Hours
- One Treatment Prevents Further Flea Infestation For At Least Four Weeks
- Kills Fleas Before They Lay Eggs

Active Ingredient	%
	By Weight
Imidacloprid; 1-[(6-Chloro-3-pyridinyl) methyl]-N-nitro-2-imidazolidinimine ...	9.1%
Inert Ingredients	<u>90.9%</u>
Total	100%



KEEP OUT OF REACH OF CHILDREN

WARNING

See Below for First Aid and Precautionary Statements

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS

Causes eye irritation. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS

For external use only.

Do not use on puppies under seven weeks of age.

As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. If signs persist, or become more severe, consult a veterinarian immediately. If your animal is on medication, consult your veterinarian before using this or any other product. For consumer questions call 1-800-255-6826. For medical emergencies call 1-877-258-2280.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice.

To Physician: Treat the patient symptomatically.

Six 0.4 mL Tubes

EPA Est. 11556-DEU-1
EPA Reg. No. 11556-117

Manufactured For
Bayer Corporation
Agriculture Division
Animal Health
Shawnee Mission, Kansas 66201 USA



(Back Panel)

Advantage® 10


(imidacloprid) Topical Solution

Fast
Effective

Once-A-Month Topical Flea Treatment for Dogs and
Puppies 7 Weeks and Older and 10 lbs and Under

- Available only through licensed practicing veterinarians
- Kills fleas within 12 hours
- Prevents reinfestation for at least 4 weeks
- Convenient, easy to apply

Six 0.4 mL Tubes



(Inside Left Panel)

Advantage® 10

(imidacloprid) Topical Solution

Once-A-Month Topical Flea Treatment for Dogs and
Puppies 7 Weeks and Older and 10 lbs and Under

Read entire label before use.

For the Prevention and Treatment of Flea Infestation on Dogs

- Available Only Through Licensed Practicing Veterinarians
- Kills 98-100% of the Fleas on Dogs Within 12 Hours
- One Treatment Prevents Further Flea Infestation For At Least Four Weeks
- Kills Fleas Before They Lay Eggs

Active Ingredient	%
	By Weight
Imidacloprid; 1-[(6-Chloro-3-pyridinyl) methyl]-N-nitro-2-imidazolidinimine ...	9.1%
Inert Ingredients	<u>90.9%</u>
Total	100%

KEEP OUT OF REACH OF CHILDREN


WARNING

See Below for First Aid and Precautionary Statements

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

Causes eye irritation. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.



HAZARDS TO DOMESTIC ANIMALS

For external use only.

Do not use on puppies under seven weeks of age.

As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. If signs persist, or become more severe, consult a veterinarian immediately. If your animal is on medication, consult your veterinarian before using this or any other product. For consumer questions call 1-800-255-6826. For medical emergencies call 1-877-258-2280.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice.

To Physician: Treat the patient symptomatically.


DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.


HOW TO APPLY

1. Use only on dogs. Do not use on other animals.
2. Remove one applicator tube from the package.






3. Hold applicator tube in an upright position. Pull cap off tube.
4. Turn the cap around and place other end of cap back on tube.
5. Twist cap to break seal, then remove cap from tube.



(Inside Right Panel)

- 6. The dog should be standing for easy application. Part the hair on the dog's back, between the shoulder blades, until the skin is visible. Place the tip of the tube on the skin and squeeze the tube twice to expel the entire contents directly on the skin.

Do not get this product in your pet's eyes or mouth.



- 7. Discard empty tube as described in Storage and Disposal.

The successive feeding activity of fleas on pets frequently elicits a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with Advantage® rapidly kills fleas and reduces the incidence of this condition.

Advantage® kills 98-100% of the existing fleas on pets within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting at least four (4) weeks. Pre-existing pupae in the environment may continue to emerge for six (6) weeks or longer depending upon the climatic conditions.

Larval flea stages in the pet's surroundings are killed following contact with an Advantage® treated pet.

Advantage® remains efficacious following a shampoo treatment, swimming or after exposure to rain or sunlight.

Monthly treatments are required for optimal control and prevention of fleas.

If re-treatment becomes necessary earlier than four weeks, do not re-treat more than once weekly.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.


Storage: Store in a cool, dry place. **Pesticide Disposal:** Securely wrap original container in several layers of newspaper and discard in trash. **Container Disposal:** Do not reuse empty container. Wrap container and put in trash.

LIMITED WARRANTY AND LIMITATION OF DAMAGES

Bayer Corporation, Agriculture Division, Animal Health warrants that this material conforms to the chemical description on the label. BAYER CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY, and no agent of Bayer Corporation is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

EPA Est. 11556-DEU-1

EPA Reg. No. 11556-117



(Label on Individual Tube)

Advantage®

9.1% Imidacloprid

0.4 mL

EPA Reg. No. 11556-117


Warning

Keep Out of Reach of Children

Read The Entire Label Before Use

BAYER

Lot No. 0000000




Reason To Issue: Revised according to PR Notice 2001-1.

Date: 01/08/01

Supersedes: 9/13/99

10/10

(Back of Blister Package)



NEXT

LABEL

11556-117

4-27-2001

1/10

Reason To Issue: Revised according to PR Notice 2001-1

Date: 01/08/01

Supersedes: 9/13/99

(Front Panel)

Advantage® 10
(imidacloprid) Topical Solution


Once-A-Month Topical Flea Treatment For Dogs and Puppies 7 Weeks and Older and 10 lbs and Under

READ THE ENTIRE LABEL BEFORE EACH USE

For the Prevention and Treatment of Flea Infestations on Dogs

- Available Only Through Licensed Practicing Veterinarians
- Kills 98-100% of the Fleas on Dogs Within 12 Hours
- One Treatment Prevents Further Flea Infestation For At Least Four Weeks
- Kills Fleas Before They Lay Eggs

Active Ingredient	%
	By Weight
Imidacloprid; 1-[(6-Chloro-3-pyridinyl) methyl]-N-nitro-2-imidazolidinimine . . .	9.1%
Inert Ingredients	90.9%
Total	100%



KEEP OUT OF REACH OF CHILDREN

WARNING

See Below for First Aid and Precautionary Statements

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS**

Causes eye irritation. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS

For external use only.

Do not use on puppies under seven weeks of age.

As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. If signs persist, or become more severe, consult a veterinarian immediately. If your animal is on medication, consult your veterinarian before using this or any other product. For consumer questions call 1-800-255-6826. For medical emergencies call 1-877-258-2280.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.


If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice.

To Physician: Treat the patient symptomatically.

Four 0.4 mL Tubes

EPA Est. 11556-DEU-1
EPA Reg. No. 11556-117

Manufactured For
Bayer Corporation
Agriculture Division
Animal Health
Shawnee Mission, Kansas 66201 USA



(Back Panel)


Advantage® 10

(imidacloprid) Topical Solution
Fast
Effective

Once-A-Month Topical Flea Treatment for Dogs and
Puppies 7 Weeks and Older and 10 lbs and Under

- Available only through licensed practicing veterinarians
- Kills fleas within 12 hours
- Prevents reinfestation for at least 4 weeks
- Convenient, easy to apply

Four 0.4 mL Tubes



(Inside Left Panel)

Advantage® 10

(imidacloprid) Topical Solution

Once-A-Month Topical Flea Treatment for Dogs and Puppies 7 Weeks and Older and 10 lbs and Under

Read entire label before use.

For the Prevention and Treatment of Flea Infestation on Dogs

- Available Only Through Licensed Practicing Veterinarians
- Kills 98-100% of the Fleas on Dogs Within 12 Hours
- One Treatment Prevents Further Flea Infestation For At Least Four Weeks
- Kills Fleas Before They Lay Eggs

Active Ingredient	%
	By Weight
Imidacloprid; 1-[(6-Chloro-3-pyridinyl) methyl]-N-nitro-2-imidazolidinimine . . .	9.1%
Inert Ingredients	<u>90.9%</u>
Total	100%


KEEP OUT OF REACH OF CHILDREN

WARNING

See Below for First Aid and Precautionary Statements

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS

Causes eye irritation. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.



HAZARDS TO DOMESTIC ANIMALS

For external use only.

Do not use on puppies under seven weeks of age.

As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. If signs persist, or become more severe, consult a veterinarian immediately. If your animal is on medication, consult your veterinarian before using this or any other product. For consumer questions call 1-800-255-6826. For medical emergencies call 1-877-258-2280.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice.


To Physician: Treat the patient symptomatically.


DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.


HOW TO APPLY

1. Use only on dogs. Do not use on other animals.
2. Remove one applicator tube from the package.






3. Hold applicator tube in an upright position. Pull cap off tube.
4. Turn the cap around and place other end of cap back on tube.
5. Twist cap to break seal, then remove cap from tube.



(Inside Right Panel)

6. The dog should be standing for easy application. Part the hair on the dog's back, between the shoulder blades, until the skin is visible. Place the tip of the tube on the skin and squeeze the tube twice to expel the entire contents directly on the skin.

Do not get this product in your pet's eyes or mouth.



7. Discard empty tube as described in Storage and Disposal.

The successive feeding activity of fleas on pets frequently elicits a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with Advantage® rapidly kills fleas and reduces the incidence of this condition.

Advantage® kills 98-100% of the existing fleas on pets within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting at least four (4) weeks. Pre-existing pupae in the environment may continue to emerge for six (6) weeks or longer depending upon the climatic conditions.

Larval flea stages in the pet's surroundings are killed following contact with an Advantage® treated pet.

Advantage® remains efficacious following a shampoo treatment, swimming or after exposure to rain or sunlight.

Monthly treatments are required for optimal control and prevention of fleas.

If re-treatment becomes necessary earlier than four weeks, do not re-treat more than once weekly.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry place. **Pesticide Disposal:** Securely wrap original container in several layers of newspaper and discard in trash. **Container Disposal:** Do not reuse empty container. Wrap container and put in trash.

LIMITED WARRANTY AND LIMITATION OF DAMAGES

Bayer Corporation, Agriculture Division, Animal Health warrants that this material conforms to the chemical description on the label. BAYER CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY, and no agent of Bayer Corporation is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

EPA Est. 11556-DEU-1

EPA Reg. No. 11556-117

9/10

(Label on Individual Tube)

Advantage®

9.1% Imidacloprid

0.4 mL

EPA Reg. No. 11556-117


Warning

Keep Out of Reach of Children

Read The Entire Label Before Use

BAYER

Lot No. 0000000



Reason To Issue: Revised according to PR Notice 2001-1

Date: 01/08/01
Supersedes: 9/13/99

10/10

(Back of Blister Package)

