PLEASE NOTE

This image contains more than one label approved for this product on this date.

Reason To Issue:

Supersedes: 9/13/99

(Front Panel)

Advantage® 18 (imidacloprid) Topical Solution

Once-A-Month Topical Flea Treatment for Cats and Kittens 8 Weeks and Older and Over 9 lbs.

READ THE ENTIRE LABEL BEFORE EACH USE

For the Prevention and Treatment of Flea Infestations

- Available Only Through Licensed Practicing Veterinarians
- Kills 98-100% of the Fleas on Cats Within 12 Hours
- One Treatment Prevents Further Flea Infestation For Up to Four Weeks
- Kills Fleas Before They Lay Eggs

| Active Ingredient | % By Weight |
|---|----------------|
| Imidacloprid; 1-[(6-Chloro-3-pyridinyl) methyl]-N-nitro-2-imidazolidinimine | 9.1% |
| Inert Ingredients | 90.9% |
| Total | 100% |

ACCEPTED

APR 2 7 2001

KEEP OUT OF REACH OF CHILDREN

WARNING

ee Below for First Aid and Precautionary Statements

PRECAUTIONARY STATEMENTS **HAZARDS** TO HUMANS

Causes eye irritation. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not use on kittens under eight weeks of age.

As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. If signs persist, or become more severe, consult a veterinarian immediately. If your animal is on medication, consult your veterinarian before using this or any other product. For consumer questions call 1-800-255-6826. For medical emergencies call 1-877-258-2280.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

To Physician: Treat the patient symptomatically.

Six 0.8 mL Tubes

EPA Est. 11556-DEU-1 EPA Reg. No. 11556-118

> Bayer Corporation Agriculture Division Animal Health Shawnee Mission, Kansas 66201 USA

Manufactured For

(Back Panel)

Advantage® 18

(imidacloprid) Topical Solution
Fast
Effective

Once-A-Month Topical Flea Treatment for Cats and Kittens 8 Weeks and Older and Over 9 lbs.

- Available only through licensed practicing veterinarians
- Kills fleas within 12 hours
- Prevents reinfestation for up to 4 weeks
- Convenient, easy to apply

Six 0.8 mL Tubes

(Inside Left Panel)

Advantage® 18

(imidacloprid) Topical Solution

Once-A-Month Topical Flea Treatment for Cats


For the Prevention and Treatment of Flea Infestation on Cats and Kittens 8 Weeks and Older and Over 9 lbs

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOW TO APPLY

- 1. Use only on cats. Do not use on other animals.
- 2. Remove one applicator tube from the package.




- 3. Hold applicator tube in an upright position. Pull cap off tube.
- 4. Turn the cap around and place other end of cap back on tube.
- 5. Twist cap to break seal, then remove cap from tube.

Reason To Issue: Revised according to PR Notice 2001-1 Date: 01/08/01

Supersedes: 9/13/99

6. Part the hair in the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube twice to expel the entire contents of the tube directly on the skin. Do not get this product in your pet's eyes or mouth. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product.



7. Discard empty tube as described in Storage and Disposal.

(Inside Right Panel)

The successive feeding activity of fleas on pets frequently elicits a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with Advantage® rapidly kills fleas and reduces the incidence of this condition.

Advantage® kills 98-100% of the fleas on pets within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting for up to four (4) weeks. Pre-existing pupae in the environment may continue to emerge for six (6) weeks or longer depending upon the climatic condition.

Larval flea stages in the pet's surroundings are killed following contact with an Advantage® treated pet.

Advantage® remains efficacious following a shampoo treatment, swimming or after exposure to rain or sunlight.

Monthly treatments are required for optimal control and prevention of fleas.

If re-treatment becomes necessary earlier than four weeks, do not re-treat more than once weekly.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry place. **Pesticide Disposal**: Securely wrap original container in several layers of newspaper and discard in trash. Container Disposal: Do not reuse empty container. Wrap container and put in trash.

LIMITED WARRANTY AND LIMITATION OF DAMAGES

Bayer Corporation, Agriculture Division, Animal Health warrants that this material conforms to the chemical description on the label. BAYER CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY, and no agent of Bayer Corporation is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct camages and shall not include consequential commercial damages such as loss of profits or values etc.

EPA Est. 11556-DEU-1

EPA Reg. No. 11556-118

1/8

Reason To Issue:

Revised according to PR Notice 2001-1

Date: 01/08/01

Supersedes: 9/13/99

(Label on Individual Tube)

 $Advantage \\ \\ \mathbb{R}$

9.1% Imidacloprid

 $0.8\ mL$

EPA Reg. No. 11556-118

Warning

Keep Out of Reach of Children

Read The Entire Label Before Use

BAYER

Lot No. 0000000

Reason To Issue:

Revised according to PR Notice 2001-1

Date: 01/08/01 Supersedes: 9/13/99

(Back of Blister Package)

NEXT

LABEL

Supersedes: 9/13/99

Reason To Issue:

Revised according to PR Notice 2001-1

(Front Panel)

Advantage® 18 (imidacloprid) Topical Solution

Once-A-Month Topical Flea Treatment for Cats and Kittens 8 Weeks and Older and Over 9 lbs.

READ THE ENTIRE LABEL BEFORE EACH USE

For the Prevention and Treatment of Flea Infestations

- Available Only Through Licensed Practicing Veterinarians
- Kills 98-100% of the Fleas on Cats Within 12 Hours
- One Treatment Prevents Further Flea Infestation For Up to Four Weeks
- Kills Fleas Before They Lay Eggs

| Active Ingredient | % By Weight |
|---|----------------|
| Imidacloprid; 1-[(6-Chloro-3-pyridinyl) methyl]-N-nitro-2-imidazolidinimine | . 9.1% |
| Inert Ingredients | 90.9% |
| Total | 100% |

ACCEPTED

APR 27 2001

Under the Federal Insecticide, Pengicide, and Rodenticide Act, as amended, for the pesticide Registers No. 177 KEEP OUT OF REACH OF CHILDREN

WARNING

Below for First Aid and Precautionary Statements

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

Causes eye irritation. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS

For external use only.

Do not use on kittens under eight weeks of age.

As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. If signs persist, or become more severe, consult a veterinarian immediately. If your animal is on medication, consult your veterinarian before using this or any other product. For consumer questions call 1-800-255-6826. For medical emergencies call 1-877-258-2280.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

To Physician: Treat the patient symptomatically.

Four 0.8 mL Tubes

EPA Est. 11556-DEU-1 EPA Reg. No. 11556-118

Manufactured For

Bayer Corporation
Agriculture Division
Animal Health
Shawnee Mission, Kansas 66201 USA

(Back Panel)

Advantage® 18

(imidacloprid) Topical Solution Fast Effective

Once-A-Month Topical Flea Treatment for Cats and Kittens 8 Weeks and Older and Over 9 lbs.

- Available only through licensed practicing veterinarians
- Kills fleas within 12 hours
- Prevents reinfestation for up to 4 weeks
- Convenient, easy to apply

Four 0.8 mL Tubes

(Inside Left Panel)

Advantage® 18

(imidacloprid) Topical Solution

Once-A-Month Topical Flea Treatment for Cats


For the Prevention and Treatment of Flea Infestation on Cats and Kittens 8 Weeks and Older and Over 9 lbs

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOW TO APPLY

- 1. Use only on cats. Do not use on other animals.
- 2. Remove one applicator tube from the package.



- 3. Hold applicator tube in an upright position. Pull cap off tube.
- 4. Turn the cap around and place other end of cap back on tube.
- 5. Twist cap to break seal, then remove cap from tube.

6. Part the hair in the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube twice to expel the entire contents of the tube directly on the skin. Do not get this product in your pet's eyes or mouth. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product.



7. Discard empty tube as described in Storage and Disposal.

(Inside Right Panel)

The successive feeding activity of fleas on pets frequently elicits a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with Advantage® rapidly kills fleas and reduces the incidence of this condition.

Advantage kills 98-100% of the fleas on pets within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting for up to four (4) weeks. Pre-existing pupae in the environment may continue to emerge for six (6) weeks or longer depending upon the climatic condition.

Larval flea stages in the pet's surroundings are killed following contact with an Advantage® treated pet.

Advantage ® remains efficacious following a shampoo treatment, swimming or after exposure to rain or sunlight.

Monthly treatments are required for optimal control and prevention of fleas.

If re-treatment becomes necessary earlier than four weeks, do not re-treat more than once weekly.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry place. Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash. Container Disposal: Do not reuse empty container. Wrap container and put in trash.

LIMITED WARRANTY AND LIMITATION OF DAMAGES

Bayer Corporation, Agriculture Division, Animal Health warrants that this material conforms to the chemical description on the label. BAYER CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY, and no agent of Bayer Corporation is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

EPA Est. 11556-DEU-1

EPA Reg. No. 11556-118

7/8

Reason To Issue:

Revised according to PR Notice 2001-1

Date: 01/08/01

Supersedes: 9/13/99

(Label on Individual Tube)

Advantage®

9.1% Imidacloprid

0.8 mL

EPA Reg. No. 11556-118

Warning

Keep Out of Reach of Children

Read The Entire Label Before Use

BAYER

Lot No. 0000000

Reason To Issue: Revised according to PR Notice 2001-1

Date: 01/08/01

Supersedes: 9/13/99

(Back of Blister Package)