

11556-124

11/5/2001

1/9



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

NOV 5 2001

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. F. Terry McNamara
Bayer Corporation
Agriculture Division, Animal Health
P.O. Box 390
Shawnee Mission, KS 66201

Subject: Amendment: Revision of Hazards to Humans to Agree with Parent Product Label
Tempo SC Ultra Premise Spray
EPA Reg. No. 11556-124
Your Submission Dated October 9, 2001

Dear Mr. McNamara:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below. Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records.

1. You must add the following statement to the paragraph under Hazards to Humans and Domestic Animals:

"Avoid breathing spray mist."

[Note: This product is a 100% repack of Bayer's Tempo SC Ultra Insecticide (EPA Reg. No. 3125-498) which bears the above inhalation precaution. Accordingly, this product must also bear this statement.]

-2-

2. You must update the First Aid section in accordance with the Agency's latest guidance in PR Notice 2001-1:

- Add the statement "Do not give anything by mouth to an unconscious person" at the end of the "If swallowed" section.
- Add the following statement at the end of the First Aid statements (above the "Note To Physician"):

"Have the product container or label with you when calling a poison control center or doctor, or going for treatment."

3. In the Tank Mixing section at the top of the second column on page 3, reference is made to the "Extended Cockroach Control with Insect Growth Regulators" section on "page 3". The "Extended Cockroach ..." section is actually located on page 4. You must correct this error.

If you have any questions regarding this action, please contact Susan Stanton of my team at (703) 305-5218.

Sincerely,

George T. LaRocca
Product Manager (13)
Insecticide Branch
Registration Division (7505C)

Enclosure

3/9

Tempo[®] SC Ultra

Premise Spray

For broad-spectrum control of crawling, flying, and wood-infesting insect pests in and around animal housing, warehouses, and processing and packing plants.

ACTIVE INGREDIENT:

Cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropane-carboxylate.....11.8%

INERT INGREDIENTS:.....88.2%
100.0%

Contains 1 lb cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate per gallon.

EPA Reg. No. 11556-124

Six 240-mL Bottles Per Carton, Two Cartons Per Case

EPA Est. No. 10370-TX-1

STOP - Read the label before use.
Keep out of reach of children.

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water

after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

FIRST AID

If **swallowed**: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or a doctor. If **on skin or clothing**: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If **in eyes**: Hold eye open and rinse slowly and gently with water for 15-20

minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center for treatment advice. If **inhaled**: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Note To Physician: No specific antidote is available. Treat the patient symptomatically.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
NOV 5 2001**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
11556-124**

4/4

Tempo SC Ultra Premise Spray

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in

neighboring areas. Do not contaminate water when cleaning equipment or when disposing of equipment wash waters. Apply this product only as specified on this label. This pesticide is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply Tempo SC Ultra Insecticide or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions for Use and Conditions of Sale before using Tempo SC Ultra Insecticide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE

OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

GENERAL INFORMATION FOR STRUCTURAL PEST CONTROL

Tempo SC Ultra Insecticide is a liquid formulation containing 1 lb active ingredient per gallon, which will provide effective knockdown and residual control of the pests listed in this label.

Tempo SC Ultra is intended for use as a general surface, spot, mist, or crack and crevice application in and around buildings and structures and their immediate surroundings. Permitted areas of use include, **but are not limited to**, dairies and dairy product processing plants, grain mills, granaries, greenhouses (structures only), industrial buildings, kitchens, laboratories, livestock housing structures, pet kennels, manufacturing establishments, poultry houses, meat, poultry

and egg processing and packing plants, meat and produce canneries, stores, warehouses, and similar structures.

Tempo SC Ultra is intended to be mixed with water and applied with hand pressurized, power operated sprayers or foam generating equipment. Spray pressure should not exceed 50 PSI at the nozzle tip when used indoors. Add the appropriate amount of product when filling sprayer tank with water, shake or agitate as necessary to mix. Diluted spray mixture can be stored overnight and applied the following day; however, mixture should be agitated prior to application to prevent uneven distribution of product. Before leaving spray mixture unagitated

Tempo SC Ultra Premise Spray

for an extended period of time, release tank pressure and elevate nozzle and hose above tank outlet, activate trigger and drain all dilution from hose back into spray tank. Re-tighten spray tank lid and do not re-pressurize until starting spray again.

Tempo SC Ultra will not stain or cause damage to any painted or varnished household surface, plastic, fabric or other surface where water applied alone causes no damage.

TANK MIXING: The user, at his discretion, can tank mix pesticides currently registered for similar use patterns, unless the product labels specifically prohibit such use. See Extended Cockroach Control with Insect Growth Regulators, page 3. It is always recommended that a small jar compatibility test using proper proportions of chemicals and water be run to check for physical compatibility prior to tank mixing.

GENERAL SURFACE APPLICATION

Use 8 milliliters of Tempo SC Ultra per 1000 sq ft in sufficient water (under conditions of severe pest infestation or when quicker knock-down and/or residual control is needed, use 16 milliliters of Tempo SC Ultra per 1000 sq ft) to adequately cover the area being treated, but which will not allow dripping or run-off to occur (see Note 1). Use a low pressure system with a fan type nozzle to uniformly apply the dilution. Applications can be made to walls, floors, ceilings, in and around cupboards, between, behind and beneath equipment, appliances, around floor drains, window and door frames, and on the underside of

shelves, drawers and in similar areas. Applications may be made to floor surfaces along walls and around air ducts, however, do not treat entire area of floor or floor coverings. All food processing surfaces and utensils should be covered or thoroughly washed following treatment. Cover exposed foodstuffs or remove from area being treated. Re-application can be made at 10-day intervals, if necessary. Do not apply where electrical short circuits could occur. Use a dust or dry bait in these areas. It is recommended to wear safety glasses, goggles or face shield and a respirator when making a general surface treatment to overhead areas.

Note 1: Volume of Water Sprays Applied To A Vertical Surface to Attain Appropriate Coverage

Type of Surface	Quarts/1000 Sq Ft	Gallons/1000 Sq Ft
Ceramic tile (glazed)	0.75	0.2
Stainless steel	0.75	0.2
Latex painted plywood (semigloss)	4.0	1.0
Masonite (smooth)	6.3	1.5
Unpainted plywood	11.0	2.7
Unpainted particle board	11.7	3.0
Cement block (semigloss latex)	12.5	3.2
Cement block (unfinished)	145.5	36.3

SPOT, MIST OR CRACK AND CREVICE APPLICATIONS

Mix 8 milliliters of Tempo SC Ultra in one gallon of water to make a dilution of 0.025% active ingredient concentration (under conditions of severe pest infestation or when quicker knock-down and/or longer residual control is needed use 16 milliliters of Tempo SC Ultra in one gallon of water to make a dilution of 0.05% active ingredient concentration). Use a low pressure system with a pin-point or variable pattern nozzle or appropriate foam generating equipment to apply the suspension or sufficient volume of foam in

specific areas such as: cracks and crevices in or behind baseboards, in floors, walls, expansion joints, areas around water and sewer pipes, wall voids, or voids in table legs and equipment where pests can hide as well as other similar areas. Spot treatments may also be made to areas including but not limited to storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks, stoves and other equipment, the underside of shelves, drawers and similar areas. Tempo SC Ultra can be reapplied at 10-day intervals, if

Tempo SC Ultra Premise Spray

necessary. Do not apply where electrical short circuits could occur. Use a dust or dry bait in

these areas.

EXTENDED COCKROACH CONTROL WITH INSECT GROWTH REGULATORS

Use Tempo SC Ultra in a tank mix combination with an Insect Growth Regulator (IGR) for extended control of American, Asian, German, oriental, brown-banded, smoky brown and other species of cockroaches. Use in food areas of food handling establishments only if all products in tank mix are registered for such use.

Make applications in accordance with those label instructions which are the same for each product. Prepare the tank mix using the recommended rate of Tempo SC Ultra and the IGR of choice by

adding the Tempo SC Ultra first and making sure it's a well mixed dilution, then adding the correct amount of IGR according to its label instructions. Follow all label instructions for use patterns and spray intervals from the IGR label to achieve maximum benefit.

Refer to the appropriate sections of the label for Tempo SC Ultra and the label for the IGR of choice for additional precautions, restrictions, limitations, and application information.

FOOD/FEED HANDLING ESTABLISHMENT APPLICATIONS

Tempo SC Ultra applications are permitted in both food/feed and non-food areas of food/feed handling establishments as a general surface, spot, or crack and crevice treatment. Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food. Serving areas where food is exposed and the facility is in operation are also considered food areas.

General Surface Application: When the facility is in operation or foods are exposed do not apply

as a general surface application in food handling areas. Do not apply directly to food products. Cover or remove all food processing and/or handling equipment during application. After application in food processing plants, bakeries, cafeterias and similar facilities, wash all equipment, benches, shelving and other surfaces which food will contact. Clean food handling or processing equipment and thoroughly rinse with clean, fresh water.

Spot, Crack and Crevice Application: Spot or crack and crevice applications may be made while the facility is in operation; however, food should be covered or removed from area being treated. Do not apply directly to food or food-handling surfaces.

INDOOR PEST CONTROL

CRAWLING INSECT PEST CONTROL

For **general household pests** and **occasional invaders** make applications in a manner as previously described under General Surface, Spot, or Crack and Crevice Application. Particular attention should be given to treating entry points and harborage areas.

For **pantry pest control** of exposed adult and immature stages of insects, make application to cupboards, shelving and storage areas. Particular attention should be given to treating cracks and crevices and other harborage areas. Remove all utensils, uncovered foodstuffs (or any with package opened), shelf paper and other objects before spraying. Allow treated surfaces to dry and cover shelves with clean paper before

replacing any utensils, foodstuff or other items. Any foodstuff accidentally contaminated with spray dilution should be discarded.

For **stored product pest control** of exposed adult and immature stages of insect pests, apply to cracks, crevices, and other surfaces where the pests have been seen or have harborage. Treat warehouses, production facilities, storage areas, rail cars, truck beds and other areas where products are stored before filling with the product. TEMPO SC may be used by pest control operators and grain producers to treat grain storage facilities and other areas noted above for stored product pest control. Cleaning of all areas prior to use of Tempo SC Ultra will increase

Tempo SC Ultra Premise Spray

levels of control. Any foodstuffs infested with pests should be removed and destroyed.

For use by pest control operators or livestock producers in livestock housing structures (including poultry houses) and pet kennels to control crawling and flying insect pests. Apply as a general surface and/or a crack and crevice spray. Control will be enhanced when facilities are cleaned and interior applications are supplemented with exterior perimeter treatments. Do not make interior applications of Tempo SC

Ultra in areas of facility where animals are present. Do not make applications to animal feedstuffs or watering equipment.

FLYING INSECT PEST CONTROL (General Surface Spray)

For control of flying insects indoors make application to surfaces as directed under Directions For Use, where the pests collect or rest. Do not apply as a "space spray" application.

OUTDOOR PEST CONTROL

For control of flying insects outdoors use a 0.05% dilution and make applications to outside surfaces of buildings, porches, patios, garages and other areas where these pests have been seen or found. Use a 0.05% dilution and make pin stream applications directly to the nests of wasps, yellow-jackets and hornets. For perimeter pest control, make a general surface application to outside surfaces of buildings, porches, patios, garages and other adjacent areas where pests have been seen or found. To help prevent infestations of buildings by occasional invaders, treat soil, turf or other substrates adjacent to buildings, the building foundation, walls, around doors and windows, soffit areas where these pests are active and may find entrance or harborage. Apply a minimum of 8 milliliters of Tempo SC Ultra per 1000 sq ft in sufficient water to provide adequate coverage.

For control of the imported red fire ant, use a 0.05% dilution of Tempo SC Ultra and thoroughly drench or inject the entire mound area. Treat all areas of the mound showing activity to provide contact control of the adult fire ants. Tempo SC Ultra will provide contact activity and residual repellency in the active mound.

When using the injection treatment method for control of the imported red fire ant, it is essential that treatment be made to the bottom of the active mound. Apply dilution or sufficient volume of foam throughout the mound using a downward and upward motion of the injector tool. Multiple injection sites may be needed to ensure complete distribution of the suspension throughout the active mound.

Do not use Tempo SC Ultra to treat fire ant mounds in any food crop production area, pasture or area where food crop production may occur.

CONTROL OF WOOD INFESTING PESTS

For control of wood infesting beetles on structure surfaces apply a 0.05% dilution of Tempo SC Ultra. For treatment of small areas, apply by brushing or spraying the suspension evenly on wood surfaces. For large or overhead areas apply as a coarse spray to thoroughly cover the area. Cover all surfaces below the area being sprayed with plastic sheeting or other material which can be disposed of by placing in trash if contamination from dripping occurs.

Sprayed surfaces should be avoided until the treated area is completely dry.

For control of above ground termites and wood-infesting beetles in localized areas, apply a 0.05% dilution or sufficient volume of foam of Tempo SC Ultra to voids and galleries in damaged wood, and in spaces between wooden structural members and between the sill plate and foundation where wood is vulnerable.

5/9

Tempo SC Ultra Premise Spray

PESTS CONTROLLED BY TEMPO*				
I. INDOOR PREMISE PESTS			II. FOOD PROCESSING PESTS	IV. OUTDOOR/ PERIMETER PESTS
1. Crawling Pests	2. Flying Pests	3. Pantry & Stored Product Pests	(Same crawling & flying pests, pantry & stored product pests as for INDOOR PREMISE) and Fruitflies, Phoridflies and Sciaridflies	Ants Bees Boxelder bugs Carpenter ants Cecidflies Centipedes Clusterflies Cockroaches Crickets Elm leaf beetle Fire ants Firebrats Flies Fleas Gnats Ground beetles Hornets Imported red fire ants Midges Millipedes Mosquitoes Moths Scorpions Silverfish Spiders Ticks Wasps Yellow-jackets
Ants (except Pharaoh) Bedbugs Carpet beetles Centipedes Cockroaches American Asian German Oriental brown-banded smoky brown Crickets Darkling beetles Earwigs Firebrats Millipedes Pillbugs Silverfish Sowbugs Spiders	Clothes moths Flies Gnats Midges Mosquitoes Moths Wasps Yellow-jackets	Beetles (exposed adults & immature stages) Cadmelle Cigarette Confused Flour Dermestid Drug-store Granary weevil Hide Larder Leather Lesser grain borer Lesser mealworm Merchant grain Mealworm Red flour Rice weevil Sawtoothed grain Warehouse Moths Indian meal Mediterranean flour	III. WOOD INFESTING PESTS Beetles Ambrosia Deathwatch False powderpost Old house borer Powderpost Carpenter ants Carpenter bees Termites Subterranean Formosan Drywood	

* For GENERAL SURFACE APPLICATION, use 16 milliliters of Tempo SC Ultra per 1000 square feet in sufficient water to adequately cover the area being treated.

* For SPOT, CRACK AND CREVICE, OR FOAM APPLICATIONS, mix 16 milliliters Tempo SC Ultra in one gallon of water to make a 0.05% active ingredient concentration.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original containers and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent

runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Kansas City Emergency Response Telephone No. is 800-414-0244 or contact Chemtrec at 800-424-9300.

9/9

Tempo SC Ultra Premise Spray

Pesticide Disposal: Pesticide wastes are hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Container Disposal: Do not use container in connection with food, feed, or drinking water. Completely empty container into application equipment. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and

Bayer Corporation
Agriculture Division, Animal Health
P.O. Box 390
Shawnee Mission, KS 66201