



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 11556-124	2. EPA Product Manager George LaRocca	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Tempo SC Ultra Premise Spray	PM#	
5. Name and Address of Applicant (Include ZIP Code) Bayer Corporation, Animal Health, Agriculture Division P.O. Box 390 Shawnee Mission, KS 66201 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated March 21, 2002
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

See attached explanation.

NOTIFICATION

JUN 18 2002

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	
				<input type="checkbox"/> Plastic	
				<input type="checkbox"/> Glass	
				<input type="checkbox"/> Paper	
				<input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted					
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Label accompanying product		
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name F. Terry McNamara		Title Director, Preclin Dev. and EPA Reg. Affairs	
		Telephone No. (include Area Code) (913) 268-2580	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature <i>F. Terry McNamara</i>		3. Title Director, Preclinical Development and EPA Reg. Affairs	
4. Typed Name F. Terry McNamara		5. Date June 6, 2002	

Bayer



Agriculture Division

Animal Health

Bayer Corporation
P.O. Box 390
Shawnee Mission, KS 66201-0390
Phone: 913 268-2000

July 18, 2002

Joyce Edwards
7505C
USEPA Headquarters
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, DC 20460

SUBJECT: Certification for Final Printed Labels for Tempo SC Ultra, EPA Reg. No. 11556-124

Dear Ms. Edwards:

As per your request, this letter serves as certification that the only changes made to the labels for the above referenced product are those detailed in the June 6, 2002 submission to the Agency (copy attached). No other changes were made to the label text.

Please call me at 913-268-2588 if you have any questions or need additional information.

Sincerely,

F. T. McNamara
Director
Preclinical Development & EPA Regulatory Affairs

FMT:GGG/lt

Enclosure

NOTIFICATION

JUN 18 2002

3/21

Attachment for Application for Pesticide Registration

Tempo® SC Ultra Premise Spray, EPA Reg. No. 11556-124

Enclosed with this application are two (2) copies of the final printed label for Bayer's Tempo® SC Ultra Premise Spray product (EPA Reg. No. 11556-124).

The final label incorporates all of the changes requested by the Agency (EPA letter dated March 21, 2002). Specifically, these changes are:


- 1) Added "with them" to the new statement in the Tank Mixing Section.
- 2) Added the statement to the directions for "above ground termites" under "Control of Wood Infesting Pests":

"Such applications for termites are not a substitute for mechanical alteration, soil treatment or foundation treatment, but are merely a supplement. For active termite infestations, get a professional inspection."

- 3) Added to a footnote to the table "Pests Controlled by Tempo" clarifying that the product is only for control of above-ground, winged forms of subterranean termites.

In addition, two other minor changes were made, specifically:

- 1) The use rate footnote for General Surface Application in the table "Pests Controlled by Tempo" were changed from "16 milliliters per 1000 square feet" to "8 milliliters per 1000 square feet." This change was made to be consistent with the lowest use rate described in the text under "Directions for Use".
- 2) The use rate footnote for Spot, Crack and Crevice Applications in the table "Pests Controlled by Tempo" were changed from "Mix 16 milliliters... to make a 0.05%..." to "Mix 8 milliliters... to make a 0.025%..." This change was made to be consistent with the lowest use rate described in the text under "Directions for Use".



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or

an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have the product container with you when calling a poison control center or doctor, or going for treatment.

Note To Physician: No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or when disposing of equipment wash waters. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or residues

on crops or weeds. Do not apply Tempo® SC Ultra insecticide or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions for Use and Conditions of Sale before using Tempo® SC Ultra Insecticide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL

NOTIFICATION

JUN 18 2002

4/21

CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

GENERAL INFORMATION FOR STRUCTURAL PEST CONTROL

Tempo® SC Ultra Insecticide is a liquid formulation containing 1 lb active ingredient per gallon, which will provide effective knockdown and residual control of the pests listed in this label.

Tempo® SC Ultra is intended for use as a general surface, spot, mist, or crack and crevice application in and around buildings and structures and their immediate surroundings. Permitted areas of use include, **but are not limited to**, dairies and dairy product processing plants, grain mills, granaries, green-

houses (structures only), industrial buildings, kitchens, laboratories, livestock housing structures, pet kennels, manufacturing establishments, poultry houses, meat, poultry and egg processing and packing plants, meat and produce canneries, stores, warehouses and similar structures.

Tempo® SC Ultra is intended to be mixed with water and applied with hand pressurized, power operated sprayers or foam generating equipment. Spray pressure should not exceed 50 PSI at the nozzle tip when used indoors. Add the appropriate amount of product when filling sprayer tank with water, shake or agitate as necessary to mix. Diluted spray mixture can be stored overnight and applied the following day; however, mixture should be agitated prior to application to prevent uneven distribution of product. Before leaving spray mixture unagitated for an extended period of time, release tank pressure and elevate nozzle and hose above tank outlet, activate trigger and drain all dilution from hose back into spray tank. Retighten spray tank lid and do not repressurize until starting spray again.

Tempo® SC Ultra will not stain or cause damage to any painted or var-

nished household surface, plastic, fabric or other surface where water applied alone causes no damage.

TANK MIXING: The user, at his discretion, can tank mix pesticides currently registered for similar use patterns, unless the product labels specifically prohibit such use. See Extended Cockroach Control with Insect Growth Regulators, page 4. It is always recommended that a small jar compatibility test using proper proportions of chemicals and water be run to check for physical compatibility prior to tank mixing. Tempo® SC Ultra is not compatible with disinfectants and should not be tank mixed with them. Tempo® SC Ultra and disinfectants should be applied separately as per label directions.

GENERAL SURFACE APPLICATION

Use 8 milliliters of Tempo® SC Ultra per 1000 sq ft in sufficient water (under conditions of severe pest infestation or when quicker knock-down and/or residual control is needed, use 16 milliliters of Tempo® SC Ultra per 1000 sq

5/21

ft) to adequately cover the area being treated, but which will not allow dripping or run-off to occur (see Note 1). Use a low pressure system with a fan type nozzle to uniformly apply the dilution. Applications can be made to walls, floors, ceilings, in and around cupboards, between, behind and beneath equipment, appliances, around floor drains, window and door frames, and on the underside of shelves, drawers and in similar areas. Applications may be made to floor surfaces along walls and around air ducts, however, do not treat entire area of floor or floor coverings. All food processing surfaces and utensils should be covered or thoroughly washed following treatment. Cover exposed food-stuffs or remove from area being treated. Re-application can be made at 10-day intervals, if necessary. Do not apply where electrical short circuits could occur. Use a dust or dry bait in these areas. It is recommended to wear safety glasses, goggles or face shield and a respirator when making a general surface treatment to overhead areas.

Note 1: Volume of Water Sprays Applied To A Vertical Surface to Attain Appropriate Coverage

Type of Surface	Quarts/ 1000 Sq Ft	Gallons/ 1000 Sq Ft
Ceramic tile (glazed)	0.75	0.2
Stainless steel	0.75	0.2
Latex painted plywood (semigloss)	4.0	1.0
Masonite (smooth)	6.3	1.5
Unpainted plywood	11.0	2.7
Unpainted particle board	11.7	3.0
Cement block (semigloss latex)	12.5	3.2
Cement block (unfinished)	145.5	36.3

SPOT, MIST OR CRACK AND CREVICE APPLICATIONS

Mix 8 milliliters of Tempo® SC Ultra in one gallon of water to make a dilution of 0.025% active ingredient concentration (under conditions of severe pest infestation or when quicker knock-down and/or longer residual control is needed use 16 milliliters of Tempo® SC Ultra in one gallon of

water to make a dilution of 0.05% active ingredient concentration). Use a low pressure system with a pin-point or variable pattern nozzle or appropriate foam generating equipment to apply the suspension or sufficient volume of foam in specific areas such as: cracks and crevices in or behind baseboards, in floors, walls, expansion joints, areas around water and sewer pipes, wall voids, or voids in table legs and equipment where pests can hide as well as other similar areas. Spot treatments may also be made to areas **including but not limited to** storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks, stoves and other equipment, the underside of shelves, drawers and similar areas. Tempo® SC Ultra can be reapplied at 10-day intervals, if necessary. Do not apply where electrical short circuits could occur. Use a dust or dry bait in these areas.

EXTENDED COCKROACH CONTROL WITH INSECT GROWTH REGULATORS

Use Tempo® SC Ultra in a tank mix combination with an Insect Growth Regulator (IGR) for extended control of American, Asian, German, oriental,

6/21

brown-banded, smoky brown and other species of cockroaches. Use in food areas of food handling establishments only if all products in tank mix are registered for such use.

Make applications in accordance with those label instructions which are the same for each product. Prepare the tank mix using the recommended rate of Tempo® SC Ultra and the IGR of choice by adding the Tempo® SC Ultra first and making sure it's a well mixed dilution, then adding the correct amount of IGR according to its label instructions. Follow all label instructions for use patterns and spray intervals from the IGR label to achieve maximum benefit.

Refer to the appropriate sections of the label for Tempo® SC Ultra and the label for the IGR of choice for additional precautions, restrictions, limitations, and application information.

FOOD/FEED HANDLING ESTABLISHMENT APPLICATIONS

Tempo® SC Ultra applications are permitted in both food/feed and non-food areas of food/feed handling establishments as a general surface, spot, or crack and crevice treatment. Food/feed handling establishments are defined as places other than pri-

vate residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food. Serving areas where food is exposed and the facility is in operation are also considered food areas.

General Surface Application: When the facility is in operation or foods are exposed do not apply as a general surface application in food handling areas. Do not apply directly to food products. Cover or remove all food processing and/or handling equipment during application. After application in food processing plants, bakeries, cafeterias and similar facilities, wash all equipment, benches, shelving and other surfaces which food will contact. Clean food handling or processing equipment and thoroughly rinse with clean, fresh water.

Spot, Crack and Crevice Application: Spot or crack and crevice applications may be made while the facility is in operation; however, food should be covered or removed from area being treated. Do not apply directly to food or food-handling surfaces.

INDOOR PEST CONTROL CRAWLING INSECT PEST CONTROL

For **general household pests** and **occasional invaders** make applications in a manner as previously described under General Surface, Spot, or Crack and Crevice Application. Particular attention should be given to treating entry points and harborage areas.

For **pantry pest control** of exposed adult and immature stages of insects, make application to cupboards, shelving and storage areas. Particular attention should be given to treating cracks and crevices and other harborage areas. Remove all utensils, uncovered foodstuffs (or any with package opened), shelf paper and other objects before spraying. Allow treated surfaces to dry and cover shelves with clean paper before replacing any utensils, foodstuff or other items. Any foodstuff accidentally contaminated with spray dilution should be discarded.

For **stored product pest control** of exposed adult and immature stages of insect pests, apply to cracks, crevices, and other surfaces where the pests have been seen or have harborage. Treat warehouses, production facilities, storage areas, and other areas

DO NOT

Tempo®
SC Ultra Premise Spray

For broad-spectrum control of crawling, flying, and wood-infesting insect pests in and around animal housing, warehouses, and processing and packing plants.

Shake well before using.

ACTIVE INGREDIENT:

Beta-cyfluthrin; Cyano(4-fluoro-3-phenoxyphenyl) methyl 3-(2,2-dichloroethenyl)-2,2-

dimethylcyclopropanecarboxylate 11.8%

OTHER INGREDIENTS 88.2%
100.0%

Contains 1 lb cyano(4-fluoro-3-phenoxyphenyl) methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate per gallon.

EPA Reg. No. 11556-124

EPA Est. No. 432-TX-1

STOP - Read the label before use.

Keep out of reach of children.

CAUTION

08714803-71005340, R.4

08711774-053499

Net Contents 240 mL



Mfg. for: Bayer Corporation, Agriculture Division,
Animal Health, Shawnee Mission, Kansas 66201 U.S.A.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have the product container with you when calling a poison control center or doctor, or going for treatment.

Note To Physician: No specific antidote is available. Treat the patient symptomatically.

See leaflet accompanying this container for complete labeling including Directions for Use, Storage and Disposal, Environmental Hazards and Conditions of Sale.

08714811-71005341, R.2
919874

08711774-053499
053100



10/8

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original containers and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Kansas City Emergency Response Telephone No. is 800-414-0244 or contact Chemtrec at 800-424-9300.

Pesticide Disposal: Pesticide wastes are hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Do not use container in connection with food, feed, or drinking water. Completely empty container into application equipment. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Extended Text™ INTEG™ has the following Patents:
U.K. & Europe 0130053,
U.S.A. 4592572, Canada
1231526.

Tempo® SC Ultra Premise Spray



For broad-spectrum control of crawling, flying, and wood-infesting insect pests in and around animal housing, warehouses, and processing and packing plants. Shake well before using.

ACTIVE INGREDIENT:

Beta-cyfluthrin; Cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-

dimethylcyclopropanecarboxylate 11.8%

OTHER INGREDIENTS 88.2%
100.0%

EPA Reg. No. 11556-124 EPA Est. No. 432-TX-1

Contains 1 lb cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate per gallon.

STOP - Read the label before use. Keep out of reach of children.

CAUTION

240 mL

08711774-053499

Bayer

Mfg. for: Bayer Corporation, Agriculture Division,
Animal Health, Shawnee Mission, Kansas 66201 U.S.A.

10/6

PESTS CONTROLLED BY TEMPO®*

I. INDOOR PREMISE PESTS			II. FOOD PROCESSING PESTS		IV. OUTDOOR/PERIMETER PESTS		
1. Crawling Pests			(Same crawling & flying pests, pantry & stored product pests as for INDOOR PREMISE) and Fruitflies Phoridflies and Sciaridflies		Ants		Ground beetles
2. Flying Pests					Bees		Hornets
Ants (except Pharaoh)	Crickets	Clothes moths			Boxelder bugs	Imported	
Bedbugs	Darkling beetles	Flies			Carpenter ants	red fire ants	
Carpet beetles	Earwigs	Gnats			Cecidflies	Midges	
Centipedes	Firebrats	Midges			Centipedes	Millipedes	
Cockroaches	Millipedes	Mosquitoes			Clusterflies	Mosquitoes	
American	Pillbugs	Moths			Cockroaches	Moths	
Asian	Silverfish	Wasps			Crickets	Scorpions	
German	Sowbugs	Yellow-jackets			Elm leaf beetle	Silverfish	
Oriental	Spiders				Fire ants	Spiders	
brown-banded					Firebrats	Ticks	
smoky brown					Flies	Wasps	
					Fleas	Yellow-jackets	
					Gnats		
3. Pantry & Stored Product Pests			III. WOOD INFESTING PESTS				
Beetles (exposed adults & immature stages)			Beetles				
Cadelle	Larder	Rice weevil	Ambrosia		Carpenter ants		
Cigarette	Leather	Sawtoothed grain	Deathwatch		Carpenter bees		
Confused Flour	Lesser grain borer	Warehouse	False powderpost		Termites		
Dermestid	Lesser mealworm	Moths	Old house borer		Subterranean**		
Drug-store	Merchant grain	Indian meal	Powderpost		Formosan		
Granary weevil	Mealworm	Mediterranean flour			Drywood		
Hide	Red flour						
			*For GENERAL SURFACE APPLICATION , use 8 milliliters of Tempo® SC Ultra per 1000 square feet in sufficient water to adequately cover the area being treated.				
			*For SPOT, CRACK AND CREVICE APPLICATIONS , mix 8 milliliters Tempo® SC Ultra in one gallon of water to make a 0.025% active ingredient concentration.				
			**Only for control of above-ground, winged forms of subterranean termites.				

10/21

where products are stored before filling with the product. Tempo® SC Ultra may be used by pest control operators and grain producers to treat grain storage facilities and other areas noted above for stored product pest control. Cleaning of all areas prior to use of Tempo® SC Ultra will increase levels of control. Any foodstuffs infested with pests should be removed and destroyed.

For use by pest control operators or livestock producers in livestock housing structures (including poultry houses) and pet kennels to control crawling and flying insect pests. Apply as a general surface and/or a crack and crevice spray. Control will be enhanced when facilities are cleaned and interior applications are supplemented with exterior perimeter treatments. Do not make interior applications of Tempo® SC Ultra in areas of facility where animals are present. Do not make applications to animal feedstuffs or watering equipment.

FLYING INSECT PEST CONTROL (General Surface Spray)

For control of flying insects indoors make application to surfaces as directed under Directions For Use,

where the pests collect or rest. Do not apply as a "space spray" application.

OUTDOOR PEST CONTROL

For control of flying insects outdoors use a 0.05% dilution and make applications to outside surfaces of buildings, porches, patios, garages and other areas where these pests have been seen or found. Use a 0.05% dilution and make pin stream applications directly to the nests of wasps, yellow-jackets and hornets. For perimeter pest control, make a general surface application to outside surfaces of buildings, porches, patios, garages and other adjacent areas where pests have been seen or found. To help prevent infestations of buildings by occasional invaders, treat soil, turf or other substrates adjacent to buildings, the building foundation, walls, around doors and windows, soffit areas where these pests are active and may find entrance or harborage. Apply a minimum of 8 milliliters of Tempo® SC Ultra per 1000 sq ft in sufficient water to provide adequate coverage.

For control of the imported red fire ant, use a 0.05% dilution of Tempo® SC Ultra and thoroughly drench or inject the entire mound area. Treat all areas of the mound showing activity to provide

contact control of the adult fire ants. Tempo® SC Ultra will provide contact activity and residual repellency in the active mound.

When using the injection treatment method for control of the imported red fire ant, it is essential that treatment be made to the bottom of the active mound. Apply dilution or sufficient volume of foam throughout the mound using a downward and upward motion of the injector tool. Multiple injection sites may be needed to ensure complete distribution of the suspension throughout the active mound.

Do not use Tempo® SC Ultra to treat fire ant mounds in any food crop production area, pasture or area where food crop production may occur.

CONTROL OF WOOD INFESTING PESTS

For control of wood infesting beetles on structure surfaces, apply a 0.05% dilution of Tempo® SC Ultra. For treatment of small areas, apply by brushing or spraying the suspension evenly on wood surfaces. For large or overhead areas, apply as a coarse spray to thoroughly cover the

area. Cover all surfaces below the area being sprayed with plastic sheeting or other material which can be disposed of by placing in trash if contamination from dripping occurs. Sprayed surfaces should be avoided until the treated area is completely dry.

For control of above ground termites and wood-infesting beetles in localized areas, apply a 0.05% dilution or sufficient volume of foam of Tempo® SC Ultra to voids and galleries in damaged wood, and in spaces between wooden structural members and between the sill plate and foundation where wood is vulnerable.

Such applications for termites are not a substitute for mechanical alteration, soil treatment or foundation treatment, but are merely a supplement. For active termite infestations, get a professional inspection.

000

000000

11/11

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or

an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have the product container with you when calling a poison control center or doctor, or going for treatment.

Note To Physician: No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or when disposing of equipment wash waters. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or residues

on crops or weeds. Do not apply Tempo® SC Ultra insecticide or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions for Use and Conditions of Sale before using Tempo® SC Ultra Insecticide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL

13/21

CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

GENERAL INFORMATION FOR STRUCTURAL PEST CONTROL

Tempo® SC Ultra Insecticide is a liquid formulation containing 1 lb active ingredient per gallon, which will provide effective knockdown and residual control of the pests listed in this label.

Tempo® SC Ultra is intended for use as a general surface, spot, mist, or crack and crevice application in and around buildings and structures and their immediate surroundings. Permitted areas of use include, but are not limited to, dairies and dairy product processing plants, grain mills, granaries, green-

houses (structures only), industrial buildings, kitchens, laboratories, livestock housing structures, pet kennels, manufacturing establishments, poultry houses, meat, poultry and egg processing and packing plants, meat and produce canneries, stores, warehouses and similar structures.

Tempo® SC Ultra is intended to be mixed with water and applied with hand pressurized, power operated sprayers or foam generating equipment. Spray pressure should not exceed 50 PSI at the nozzle tip when used indoors. Add the appropriate amount of product when filling sprayer tank with water, shake or agitate as necessary to mix. Diluted spray mixture can be stored overnight and applied the following day; however, mixture should be agitated prior to application to prevent uneven distribution of product. Before leaving spray mixture unagitated for an extended period of time, release tank pressure and elevate nozzle and hose above tank outlet, activate trigger and drain all dilution from hose back into spray tank. Retighten spray tank lid and do not repressurize until starting spray again.

Tempo® SC Ultra will not stain or cause damage to any painted or var-

nished household surface, plastic, fabric or other surface where water applied alone causes no damage.

TANK MIXING: The user, at his discretion, can tank mix pesticides currently registered for similar use patterns, unless the product labels specifically prohibit such use. See Extended Cockroach Control with Insect Growth Regulators, page 4. It is always recommended that a small jar compatibility test using proper proportions of chemicals and water be run to check for physical compatibility prior to tank mixing. Tempo® SC Ultra is not compatible with disinfectants and should not be tank mixed with them. Tempo® SC Ultra and disinfectants should be applied separately as per label directions.

GENERAL SURFACE APPLICATION

Use 8 milliliters of Tempo® SC Ultra per 1000 sq ft in sufficient water (under conditions of severe pest infestation or when quicker knock-down and/or residual control is needed, use 16 milliliters of Tempo® SC Ultra per 1000 sq

304100

14/21

ft) to adequately cover the area being treated, but which will not allow dripping or run-off to occur (see Note 1). Use a low pressure system with a fan type nozzle to uniformly apply the dilution. Applications can be made to walls, floors, ceilings, in and around cupboards, between, behind and beneath equipment, appliances, around floor drains, window and door frames, and on the underside of shelves, drawers and in similar areas. Applications may be made to floor surfaces along walls and around air ducts, however, do not treat entire area of floor or floor coverings. All food processing surfaces and utensils should be covered or thoroughly washed following treatment. Cover exposed food-stuffs or remove from area being treated. Re-application can be made at 10-day intervals, if necessary. Do not apply where electrical short circuits could occur. Use a dust or dry bait in these areas. It is recommended to wear safety glasses, goggles or face shield and a respirator when making a general surface treatment to overhead areas.

Note 1: Volume of Water Sprays Applied To A Vertical Surface to Attain Appropriate Coverage

Type of Surface	Quarts/ 1000 Sq Ft	Gallons/ 1000 Sq Ft
Ceramic tile (glazed)	0.75	0.2
Stainless steel	0.75	0.2
Latex painted plywood (semigloss)	4.0	1.0
Masonite (smooth)	6.3	1.5
Unpainted plywood	11.0	2.7
Unpainted particle board	11.7	3.0
Cement block (semigloss latex)	12.5	3.2
Cement block (unfinished)	145.5	36.3

SPOT, MIST OR CRACK AND CREVICE APPLICATIONS

Mix 8 milliliters of Tempo® SC Ultra in one gallon of water to make a dilution of 0.025% active ingredient concentration (under conditions of severe pest infestation or when quicker knock-down and/or longer residual control is needed use 16 milliliters of Tempo® SC Ultra in one gallon of

water to make a dilution of 0.05% active ingredient concentration). Use a low pressure system with a pin-point or variable pattern nozzle or appropriate foam generating equipment to apply the suspension or sufficient volume of foam in specific areas such as: cracks and crevices in or behind baseboards, in floors, walls, expansion joints, areas around water and sewer pipes, wall voids, or voids in table legs and equipment where pests can hide as well as other similar areas. Spot treatments may also be made to areas **including but not limited to** storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks, stoves and other equipment, the underside of shelves, drawers and similar areas. Tempo® SC Ultra can be reapplied at 10-day intervals, if necessary. Do not apply where electrical short circuits could occur. Use a dust or dry bait in these areas.

EXTENDED COCKROACH CONTROL WITH INSECT GROWTH REGULATORS

Use Tempo® SC Ultra in a tank mix combination with an Insect Growth Regulator (IGR) for extended control of American, Asian, German, oriental,

20.4.10.00

15/21

brown-banded, smoky brown and other species of cockroaches. Use in food areas of food handling establishments only if all products in tank mix are registered for such use.

Make applications in accordance with those label instructions which are the same for each product. Prepare the tank mix using the recommended rate of Tempo® SC Ultra and the IGR of choice by adding the Tempo® SC Ultra first and making sure it's a well mixed dilution, then adding the correct amount of IGR according to its label instructions. Follow all label instructions for use patterns and spray intervals from the IGR label to achieve maximum benefit.

Refer to the appropriate sections of the label for Tempo® SC Ultra and the label for the IGR of choice for additional precautions, restrictions, limitations, and application information.

FOOD/FEED HANDLING ESTABLISHMENT APPLICATIONS

Tempo® SC Ultra applications are permitted in both food/feed and non-food areas of food/feed handling establishments as a general surface, spot, or crack and crevice treatment. Food/feed handling establishments are defined as places other than pri-

vate residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food. Serving areas where food is exposed and the facility is in operation are also considered food areas.

General Surface Application: When the facility is in operation or foods are exposed do not apply as a general surface application in food handling areas. Do not apply directly to food products. Cover or remove all food processing and/or handling equipment during application. After application in food processing plants, bakeries, cafeterias and similar facilities, wash all equipment, benches, shelving and other surfaces which food will contact. Clean food handling or processing equipment and thoroughly rinse with clean, fresh water.

Spot, Crack and Crevice Application: Spot or crack and crevice applications may be made while the facility is in operation; however, food should be covered or removed from area being treated. Do not apply directly to food or food-handling surfaces.

INDOOR PEST CONTROL CRAWLING INSECT PEST CONTROL

For **general household pests** and **occasional invaders** make applications in a manner as previously described under General Surface, Spot, or Crack and Crevice Application. Particular attention should be given to treating entry points and harborage areas.

For **pantry pest control** of exposed adult and immature stages of insects, make application to cupboards, shelving and storage areas. Particular attention should be given to treating cracks and crevices and other harborage areas. Remove all utensils, uncovered foodstuffs (or any with package opened), shelf paper and other objects before spraying. Allow treated surfaces to dry and cover shelves with clean paper before replacing any utensils, foodstuff or other items. Any foodstuff accidentally contaminated with spray dilution should be discarded.

For **stored product pest control** of exposed adult and immature stages of insect pests, apply to cracks, crevices, and other surfaces where the pests have been seen or have harborage. Treat warehouses, production facilities, storage areas, and other areas

12/9/1

Tempo®
SC Ultra Premise Spray

For broad-spectrum control of crawling, flying, and wood-infesting insect pests in and around animal housing, warehouses, and processing and packing plants.

Shake well before using.

ACTIVE INGREDIENT:

Beta-cyfluthrin; Cyano(4-fluoro-3-phenoxyphenyl) methyl 3-(2,2-dichloroethenyl)-2,2-

dimethylcyclopropanecarboxylate..... 11.8%

OTHER INGREDIENTS 88.2%

100.0%

Contains 1 lb cyano(4-fluoro-3-phenoxyphenyl) methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate per gallon.

EPA Reg. No. 11556-124

EPA Est. No. 432-TX-1

STOP - Read the label before use.

Keep out of reach of children.

CAUTION

08714803-71005340, R.4

08711774-053499

Net Contents 240 mL



Mfg. for: Bayer Corporation, Agriculture Division,
Animal Health, Shawnee Mission, Kansas 66201 U.S.A.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have the product container with you when calling a poison control center or doctor, or going for treatment.

Note To Physician: No specific antidote is available. Treat the patient symptomatically.

See leaflet accompanying this container for complete labeling including Directions for Use, Storage and Disposal, Environmental Hazards and Conditions of Sale.

087114811-71005341, R.2
919874

08711774-053499
053100



17/21

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original containers and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Kansas City Emergency Response Telephone No. is 800-414-0244 or contact Chemtrec at 800-424-9300.

Pesticide Disposal: Pesticide wastes are hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Do not use container in connection with food, feed, or drinking water. Completely empty container into application equipment. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Extended Text™ INTEG™ has the following Patents:
U.K. & Europe 0130053,
U.S.A. 4592572, Canada
1231526.

Tempo® SC Ultra Premise Spray



For broad-spectrum control of crawling, flying, and wood-infesting insect pests in and around animal housing, warehouses, and processing and packing plants.

Shake well before using.

ACTIVE INGREDIENT:

Beta-cyfluthrin, Cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate	11.8%
OTHER INGREDIENTS	88.2%
	100.0%

EPA Reg. No. 11556-124 EPA Est. No. 432-TX-1

Contains 1 lb cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate per gallon.

STOP - Read the label before use. Keep out of reach of children.

CAUTION

240 mL

Bayer

08714803-
71005340, R.4

08711774-053499

Mfg. for: Bayer Corporation, Agriculture Division,
Animal Health, Shawnee Mission, Kansas 66201 U.S.A.

18/81

PESTS CONTROLLED BY TEMPO®*

I. INDOOR PREMISE PESTS			II. FOOD PROCESSING PESTS	IV. OUTDOOR/PERIMETER PESTS
1. Crawling Pests Ants (except Pharaoh) Bedbugs Carpet beetles Centipedes Cockroaches American Asian German Oriental brown-banded smoky brown 2. Flying Pests Crickets Darkling beetles Earwigs Firebrats Millipedes Pillbugs Silverfish Sowbugs Spiders 3. Pantry & Stored Product Pests Beetles (exposed adults & immature stages) Cadelle Cigarette Confused Flour Dermestid Drug-store Granary weevil Hide Larder Leather Lesser grain borer Lesser mealworm Merchant grain Mealworm Red flour Rice weevil Sawtoothed grain Warehouse Moths Indian meal Mediterranean flour 2. Flying Pests Clothes moths Flies Gnats Midges Mosquitoes Moths Wasps Yellow-jackets			(Same crawling & flying pests, pantry & stored product pests as for INDOOR PREMISE) and Fruitflies Phoridflies and Sciariidflies	Ants Bees Boxelder bugs Carpenter ants Cecidflies Centipedes Clusterflies Cockroaches Crickets Elm leaf beetle Fire ants Firebrats Flies Fleas Gnats Ground beetles Hornets Imported red fire ants Midges Millipedes Mosquitoes Moths Scorpions Silverfish Spiders Ticks Wasps Yellow-jackets
			III. WOOD INFESTING PESTS Beetles Ambrosia Deathwatch False powderpost Old house borer Powderpost Carpenter ants Carpenter bees Termites Subterranean** Formosan Drywood	
			*For GENERAL SURFACE APPLICATION , use 8 milliliters of Tempo® SC Ultra per 1000 square feet in sufficient water to adequately cover the area being treated. *For SPOT, CRACK AND CREVICE APPLICATIONS , mix 8 milliliters Tempo® SC Ultra in one gallon of water to make a 0.025% active ingredient concentration. **Only for control of above-ground, winged forms of subterranean termites.	

19/1

where products are stored before filling with the product. Tempo® SC Ultra may be used by pest control operators and grain producers to treat grain storage facilities and other areas noted above for stored product pest control. Cleaning of all areas prior to use of Tempo® SC Ultra will increase levels of control. Any foodstuffs infested with pests should be removed and destroyed.

For use by pest control operators or livestock producers in livestock housing structures (including poultry houses) and pet kennels to control crawling and flying insect pests. Apply as a general surface and/or a crack and crevice spray. Control will be enhanced when facilities are cleaned and interior applications are supplemented with exterior perimeter treatments. Do not make interior applications of Tempo® SC Ultra in areas of facility where animals are present. Do not make applications to animal feedstuffs or watering equipment.

FLYING INSECT PEST CONTROL (General Surface Spray)

For control of flying insects indoors make application to surfaces as directed under Directions For Use,

where the pests collect or rest. **Do not apply as a "space spray" application.**

OUTDOOR PEST CONTROL

For control of flying insects outdoors use a 0.05% dilution and make applications to outside surfaces of buildings, porches, patios, garages and other areas where these pests have been seen or found. Use a 0.05% dilution and make pin stream applications directly to the nests of wasps, yellow-jackets and hornets. **For perimeter pest control**, make a general surface application to outside surfaces of buildings, porches, patios, garages and other adjacent areas where pests have been seen or found. To help prevent infestations of buildings by occasional invaders, treat soil, turf or other substrates adjacent to buildings, the building foundation, walls, around doors and windows, soffit areas where these pests are active and may find entrance or harborage. Apply a minimum of 8 milliliters of Tempo® SC Ultra per 1000 sq ft in sufficient water to provide adequate coverage.

For control of the imported red fire ant, use a 0.05% dilution of Tempo® SC Ultra and thoroughly drench or inject the entire mound area. Treat all areas of the mound showing activity to provide

contact control of the adult fire ants. Tempo® SC Ultra will provide contact activity and residual repellency in the active mound.

When using the injection treatment method for control of the imported red fire ant, it is essential that treatment be made to the bottom of the active mound. Apply dilution or sufficient volume of foam throughout the mound using a downward and upward motion of the injector tool. Multiple injection sites may be needed to ensure complete distribution of the suspension throughout the active mound.

Do not use Tempo® SC Ultra to treat fire ant mounds in any food crop production area, pasture or area where food crop production may occur.

CONTROL OF WOOD INFESTING PESTS

For control of wood infesting beetles on structure surfaces, apply a 0.05% dilution of Tempo® SC Ultra. For treatment of small areas, apply by brushing or spraying the suspension evenly on wood surfaces. For large or overhead areas, apply as a coarse spray to thoroughly cover the

area. Cover all surfaces below the area being sprayed with plastic sheeting or other material which can be disposed of by placing in trash if contamination from dripping occurs. Sprayed surfaces should be avoided until the treated area is completely dry.

For control of above ground termites and wood-infesting beetles in localized areas, apply a 0.05% dilution or sufficient volume of foam of Tempo® SC Ultra to voids and galleries in damaged wood, and in spaces between wooden structural members and between the sill plate and foundation where wood is vulnerable.

Such applications for termites are not a substitute for mechanical alteration, soil treatment or foundation treatment, but are merely a supplement. For active termite infestations, get a professional inspection.

10/20

"The fact that the FBI is not releasing the tapes is a serious problem," says Robert J. Rabinowitz, executive director of the American Civil Liberties Union. "The FBI is not releasing the tapes because it is afraid of the public reaction. The FBI is not releasing the tapes because it is afraid of the public reaction. The FBI is not releasing the tapes because it is afraid of the public reaction."

[illegible][illegible][illegible]

FISH CONTROLLED BY REPORT	
<p>1. Fishery Name</p> <p>2. Fishery Type</p> <p>3. Fishery Location</p> <p>4. Fishery Description</p> <p>5. Fishery Management</p> <p>6. Fishery Status</p> <p>7. Fishery Notes</p>	<p>8. Fishery Type</p> <p>9. Fishery Location</p> <p>10. Fishery Description</p> <p>11. Fishery Management</p> <p>12. Fishery Status</p> <p>13. Fishery Notes</p>
<p>14. Fishery Name</p> <p>15. Fishery Type</p> <p>16. Fishery Location</p> <p>17. Fishery Description</p> <p>18. Fishery Management</p> <p>19. Fishery Status</p> <p>20. Fishery Notes</p>	<p>21. Fishery Type</p> <p>22. Fishery Location</p> <p>23. Fishery Description</p> <p>24. Fishery Management</p> <p>25. Fishery Status</p> <p>26. Fishery Notes</p>

[illegible][illegible]

Category	Topic	Ref
General	General	1
General	General	2
General	General	3
General	General	4
General	General	5
General	General	6
General	General	7
General	General	8
General	General	9
General	General	10
General	General	11
General	General	12
General	General	13
General	General	14
General	General	15
General	General	16
General	General	17
General	General	18
General	General	19
General	General	20
General	General	21
General	General	22
General	General	23
General	General	24
General	General	25
General	General	26
General	General	27
General	General	28
General	General	29
General	General	30
General	General	31
General	General	32
General	General	33
General	General	34
General	General	35
General	General	36
General	General	37
General	General	38
General	General	39
General	General	40
General	General	41
General	General	42
General	General	43
General	General	44
General	General	45
General	General	46
General	General	47
General	General	48
General	General	49
General	General	50
General	General	51
General	General	52
General	General	53
General	General	54
General	General	55
General	General	56
General	General	57
General	General	58
General	General	59
General	General	60
General	General	61
General	General	62
General	General	63
General	General	64
General	General	65
General	General	66
General	General	67
General	General	68
General	General	69
General	General	70
General	General	71
General	General	72
General	General	73
General	General	74
General	General	75
General	General	76
General	General	77
General	General	78
General	General	79
General	General	80
General	General	81
General	General	82
General	General	83
General	General	84
General	General	85
General	General	86
General	General	87
General	General	88
General	General	89
General	General	90
General	General	91
General	General	92
General	General	93
General	General	94
General	General	95
General	General	96
General	General	97
General	General	98
General	General	99
General	General	100

[illegible]

INDIAN TRUSTS CONFUSED
CHIEFS AND MEMBERS ASK QUESTIONS

CHIEFS AND MEMBERS OF THE INDIAN TRUSTS COMMISSION, in a session of the House of Commons, asked questions of the minister of Indian Affairs, Mr. J. G. McLean, on the subject of the Indian Trusts Commission. The minister replied that the commission was not yet organized, and that the members of the commission were not yet appointed. He stated that the commission was created by the Indian Trusts Act, which was passed in 1927. The commission was to be composed of a chairman and four members, one of whom was to be a member of the Senate. The minister stated that the commission was to be responsible for the management of the Indian Trusts, and that it was to be empowered to make recommendations to the government regarding the Indian Trusts. He stated that the commission was to be established as soon as possible, and that the members of the commission were to be appointed as soon as possible. He stated that the commission was to be responsible for the management of the Indian Trusts, and that it was to be empowered to make recommendations to the government regarding the Indian Trusts. He stated that the commission was to be established as soon as possible, and that the members of the commission were to be appointed as soon as possible.

the president's visit, regional leaders in the West and Midwest are working to bring the president to their area. "We're going to be working on that," says a spokesman for the National Governors' Association. "We're going to be working on that."

between Peoples' weekly themes and the kind of products that are sold in the stores. The kind of products that are sold in the stores are the kind of products that are sold in the stores.

power by making a submission of a letter to the appropriate authority. The letter should be signed by the person making the submission and should be addressed to the appropriate authority. The letter should state the name of the person making the submission, the address, and the date. The letter should also state the reason for the submission and the action requested. The letter should be submitted to the appropriate authority in a sealed envelope. The letter should be submitted to the appropriate authority in a sealed envelope. The letter should be submitted to the appropriate authority in a sealed envelope.

Date	Male		Female	
	Number	% ^a	Number	% ^a
April 10	1	0.2	0	0
April 11	0	0	0	0
April 12	0	0	0	0
April 13	0	0	0	0
April 14	0	0	0	0
April 15	0	0	0	0
April 16	0	0	0	0
April 17	0	0	0	0
April 18	0	0	0	0
April 19	0	0	0	0
April 20	0	0	0	0
April 21	0	0	0	0
April 22	0	0	0	0
April 23	0	0	0	0
April 24	0	0	0	0
April 25	0	0	0	0
April 26	0	0	0	0
April 27	0	0	0	0
April 28	0	0	0	0
April 29	0	0	0	0
April 30	0	0	0	0
May 1	0	0	0	0
May 2	0	0	0	0
May 3	0	0	0	0
May 4	0	0	0	0
May 5	0	0	0	0
May 6	0	0	0	0
May 7	0	0	0	0
May 8	0	0	0	0
May 9	0	0	0	0
May 10	0	0	0	0
May 11	0	0	0	0
May 12	0	0	0	0
May 13	0	0	0	0
May 14	0	0	0	0
May 15	0	0	0	0
May 16	0	0	0	0
May 17	0	0	0	0
May 18	0	0	0	0
May 19	0	0	0	0
May 20	0	0	0	0
May 21	0	0	0	0
May 22	0	0	0	0
May 23	0	0	0	0
May 24	0	0	0	0
May 25	0	0	0	0
May 26	0	0	0	0
May 27	0	0	0	0
May 28	0	0	0	0
May 29	0	0	0	0
May 30	0	0	0	0
May 31	0	0	0	0
June 1	0	0	0	0
June 2	0	0	0	0
June 3	0	0	0	0
June 4	0	0	0	0
June 5	0	0	0	0
June 6	0	0	0	0
June 7	0	0	0	0
June 8	0	0	0	0
June 9	0	0	0	0
June 10	0	0	0	0
June 11	0	0	0	0
June 12	0	0	0	0
June 13	0	0	0	0
June 14	0	0	0	0
June 15	0	0	0	0
June 16	0	0	0	0
June 17	0	0	0	0
June 18	0	0	0	0
June 19	0	0	0	0
June 20	0	0	0	0
June 21	0	0	0	0
June 22	0	0	0	0
June 23	0	0	0	0
June 24	0	0	0	0
June 25	0	0	0	0
June 26	0	0	0	0
June 27	0	0	0	0
June 28	0	0	0	0
June 29	0	0	0	0
June 30	0	0	0	0
July 1	0	0	0	0
July 2	0	0	0	0
July 3	0	0	0	0
July 4	0	0	0	0
July 5	0	0	0	0
July 6	0	0	0	0
July 7	0	0	0	0

[illegible]

which is often used to alter the pH of a solution. The pH of a solution is a measure of its acidity or basicity. The pH scale ranges from 0 to 14, with 7 being neutral. A pH of 0 is highly acidic, and a pH of 14 is highly basic. The pH of a solution can be changed by adding an acid or a base. For example, adding hydrochloric acid (HCl) to a solution will lower its pH, while adding sodium hydroxide (NaOH) will raise its pH. The pH of a solution is important in many chemical and biological processes, and it is often used as a measure of the concentration of a particular substance in a solution.

honest disclosure only. Several years ago, the National Multiple Sclerosis Society, a Washington, D.C., organization, was the first to make such a disclosure. The society's annual report, which is available to the public, lists the names of the people who have donated to the society and the amount of the donation. The society also lists the names of the people who have received grants from the society and the amount of the grant. The society's annual report is a valuable source of information for the public and for the media. It is a good idea to read the annual report of the National Multiple Sclerosis Society to learn more about the society and its work.

[illegible][illegible][illegible][illegible]