

11556-124

10-31-2002

1/21

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 11556-124	2. EPA Product Manager George LaRocca	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Tempo SC Ultra Premise Spray	PM#	
5. Name and Address of Applicant (Include ZIP Code) Bayer Corporation, Animal Health, Agriculture Division P.O. Box 390 Shawnee Mission, KS 66201 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____	<b>NOTIFICATION</b>  OCT 31 2002
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

See attached explanation.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal Plastic Glass Paper Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Label accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name F. Terry McNamara	Title Director, Preclin Dev. and EPA Reg. Affairs	Telephone No. (Include Area Code) (913) 268-2588
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>F. Terry McNamara</i>	3. Title Director, Preclinical Development and EPA Reg. Affairs	
4. Typed Name F. Terry McNamara	5. Date October 21, 2002	

2/21

Attachment for Application for Pesticide Registration  
Tempo® SC Ultra Premise Spray, EPA Reg. No. 11556-124


Enclosed with this application are five (5) copies of the final printed labels for Bayer's Tempo® SC Ultra Premise Spray (EPA Reg. No. 11556-124). This label reflects a new package size, specifically, 900 mL per bottle. This new size bottle will be sold concurrently with the old, 240 mL product. The only change made to the enclosed label is the contents statement. No other changes were made and all other text, except for the contents statement, is identical to the current stamped approved label for EPA Reg. No. 11556-124. This change in package size and contents statement is being made by Notification as per PR Notice 98-10, section II-E.

This Notification is consistent with the guidance in PR Notice 98-10 and the requirements of EPA's regulations in 40 CFR 156.10 and 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula for this product. Bayer understands it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statements to EPA. Bayer further understands that if this Notification is not consistent with the guidance of PR Notice 98-10 and the requirements of 40 CFR 156.10 and 40 CRF 152.46, this product may be in violation of FIFRA and Bayer may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**NOTIFICATION**

**OCT 31 2002**


3/24



Label Back

4<sup>13/16</sup>

3<sup>3/4</sup>



NOTIFICATION  
OCT 31 2002

Label Front

PRESS TO RESEAL

3<sup>1/4</sup>  
Booklet

4/21

3 3/4

# Tempo<sup>®</sup> SC Ultra Premise Spray

For broad-spectrum control of crawling, flying, and wood-infesting insect pests in and around animal housing, warehouses, and processing and packing plants.

**SHAKE WELL BEFORE USING.**

**ACTIVE INGREDIENT:**

Beta-cyfluthrin; Cyano(4-fluoro-3-phenoxyphenyl)  
methyl 3-(2,2-dichloroethenyl)-2,2-  
dimethylcyclopropanecarboxylate ..... 11.8%

**OTHER INGREDIENTS** ..... 88.2%  
100.0%

EPA Reg. No. 11556-124

EPA Est. No. 432-TX-1

Contains 1 lb cyano(4-fluoro-3-phenoxyphenyl) methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate per gallon.

**STOP - Read the label before use. Keep out of reach of children.**

## CAUTION

See leaflet accompanying this container for complete labeling including Directions for Use, Storage and Disposal, Conditions of Sale, To Physician, etc.

08731643-066299, R.0

919882

**30.43 fl oz (900 mL)**



Mfg. for: Bayer Corporation, Agriculture Division,  
Animal Health, Shawnee Mission, Kansas 66201 U.S.A.

PRESS TO RESEAL

3 1/4  
Booklet

5/21

■ Black

# Tempo<sup>®</sup> SC Ultra Premise Spray

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

### FIRST AID

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have the product container with you when calling a poison control center or doctor, or going for treatment.

**Note To Physician:** No specific antidote is available. Treat the patient symptomatically.

See leaflet accompanying this container for complete labeling including Directions for Use, Storage and Disposal, Environmental Hazards and Conditions of Sale.

08731643-066299, R.0

909882




7 24089 00662 2

4<sup>13</sup>/<sub>16</sub>

3<sup>3</sup>/<sub>4</sub>

el  
k →





# Tempo<sup>®</sup> SC Ultra Premise Spray



For broad-spectrum control of crawling, flying, and wood-infesting insect pests in and around animal housing, warehouses, and processing and packing plants.

**SHAKE WELL BEFORE USING.**

**ACTIVE INGREDIENT:**

Beta-cyfluthrin; Cyano(4-fluoro-3-phenoxyphenyl)

methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate ... 11.8%

**OTHER INGREDIENTS** ..... 88.2%  
100.0%

EPA Reg. No. 11556-124

EPA Est. No. 432-TX-1

Contains 1 lb cyano(4-fluoro-3-phenoxyphenyl) methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate per gallon.

**STOP - Read the label before use. Keep out of reach of children.**

08731643-066299, R.0

919882

**CAUTION**  
**30.43 fl oz (900 mL)**



Mfg. for: Bayer Corporation, Agriculture Division,  
Animal Health, Shawnee Mission, Kansas 66201 U.S.A.

INSEAL™

FRONT COVER

8 panel Template  
to fit 900ml size  
special shape Inseal™  
Label design.

Available text areas  
are as shown. However,  
text may flow across  
fold lines from panel  
to panel.

- PMS 368
- Cyan
- Black

Center line to view piece backed up

10/2

## **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

### **FIRST AID**

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have the product container with you when calling a poison control center or doctor, or going for treatment.

**Note To Physician:** No specific antidote is available. Treat the patient symptomatically.

**PAGE 2**

## ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or when disposing of equipment wash waters. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply Tempo® SC Ultra insecticide or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**IMPORTANT:** Read these entire Directions for Use and Conditions of Sale before using Tempo® SC Ultra Insecticide.

**CONDITIONS OF SALE:** THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH


INDOORS AND OUTDOORS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

## GENERAL INFORMATION FOR STRUCTURAL PEST CONTROL

Tempo® SC Ultra Insecticide is a liquid formulation containing 1 lb active ingredient per gallon, which will provide effective knockdown and residual control of the pests listed in this label.

Tempo® SC Ultra is intended for use as a general surface, spot, mist, or crack and crevice application in and around buildings and structures and their immediate surroundings. Permitted areas of use include, **but are not limited to**, dairies and dairy product processing plants, grain mills, granaries, greenhouses (structures only), industrial buildings, kitchens, laboratories, livestock housing structures,

11/21




pet kennels, manufacturing establishments, poultry houses, meat, poultry and egg processing and packing plants, meat and produce canneries, stores, warehouses and similar structures.

Tempo® SC Ultra is intended to be mixed with water and applied with hand pressurized, power operated sprayers or foam generating equipment. Spray pressure should not exceed 50 PSI at the nozzle tip when used indoors. Add the appropriate amount of product when filling sprayer tank with water, shake or agitate as necessary to mix. Diluted spray mixture can be stored overnight and applied the following day; however, mixture should be agitated prior to application to prevent uneven distribution of product. Before leaving spray mixture unagitated for an extended period of time, release tank pressure and elevate nozzle and hose above tank outlet, activate trigger and drain all dilution from hose back into spray tank. Re-tighten spray tank lid and do not re-pressurize until starting spray again.

Tempo® SC Ultra will not stain or cause damage to any painted or varnished household surface, plastic, fabric or other surface where water applied alone causes no damage.

**TANK MIXING:** The user, at his discretion, can tank mix pesticides currently registered for similar use patterns, unless the product labels specifically prohibit such use. See Extended Cockroach Control with Insect Growth Regulators, page 4. It is always recommended that a small jar compatibility test using proper proportions of chemicals and water be run to check for physical compatibility prior to tank mixing. Tempo® SC Ultra is not compatible with disinfectants and should not be tank mixed with them. Tempo® SC Ultra and disinfectants should be applied seperately as per label directions.

12/21



### GENERAL SURFACE APPLICATION


Use 8 milliliters of Tempo® SC Ultra per 1000 sq ft in sufficient water (under conditions of severe pest infestation or when quicker knock-down and/or residual control is needed, use 16 milliliters of Tempo® SC Ultra per 1000 sq ft) to adequately cover the area being treated, but which will not allow dripping or run-off to occur (see Note 1). Use a low pressure system with a fan type nozzle to uniformly apply the dilution. Applications can be made to walls, floors, ceilings, in and around cupboards, between, behind and beneath equipment, appliances, around floor drains, window and door frames, and on the underside of shelves, drawers and in similar areas. Applications may be made to floor surfaces along walls and around air ducts, however, do not treat entire area of floor or floor coverings. All food processing surfaces and utensils should be covered or thoroughly washed following treatment. Cover exposed foodstuffs or remove from area being treated. Re-application can be made at 10-day intervals, if necessary. Do not apply where electrical short circuits could occur. Use a dust or dry bait in these areas. It is recommended to wear safety glasses, goggles or face shield and a respirator when making a general surface treatment to overhead areas.

**Note 1: Volume of Water Sprays Applied To A Vertical Surface to Attain Appropriate Coverage**

<b>Type of Surface</b>	<b>Quarts/ 1000 Sq Ft</b>	<b>Gallons/ 1000 Sq Ft</b>
Ceramic tile (glazed)	0.75	0.2
Stainless steel	0.75	0.2
Latex painted plywood (semigloss)	4.0	1.0
Masonite (smooth)	6.3	1.5
Unpainted plywood	11.0	2.7
Unpainted particle board	11.7	3.0
Cement block (semigloss latex)	12.5	3.2
Cement block (unfinished)	145.5	36.3

**SPOT, MIST OR CRACK AND  
CREVICE APPLICATIONS**

**Mix 8 milliliters of Tempo® SC Ultra in one gallon of water to make a dilution of 0.025% active ingredient concentration (under conditions of severe pest infestation or when quicker knock-down and/or longer residual control is needed use 16 milliliters of Tempo® SC Ultra in one gallon of water to make a dilution of 0.05% active ingredient concentration). Use a low pressure system with a pin-point or variable pattern nozzle or appropriate foam generating equipment to apply the suspension or sufficient volume of foam in specific areas such as: cracks and crevices in or behind baseboards, in floors, walls, expansion joints, areas around water**



and sewer pipes, wall voids, or voids in table legs and equipment where pests can hide as well as other similar areas. Spot treatments may also be made to areas **including but not limited to** storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks, stoves and other equipment, the underside of shelves, drawers and similar areas. Tempo® SC Ultra can be reapplied at 10-day intervals, if necessary. Do not apply where electrical short circuits could occur. Use a dust or dry bait in these areas.

### **EXTENDED COCKROACH CONTROL WITH INSECT GROWTH REGULATORS**

Use Tempo® SC Ultra in a tank mix combination with an Insect Growth Regulator (IGR) for extended control of American, Asian, German, oriental, brown-banded, smoky brown and other species of cockroaches. Use in food areas of food handling establishments only if all products in tank mix are registered for such use.

Make applications in accordance with those label instructions which are the same for each product. Prepare the tank mix using the recommended rate of Tempo® SC Ultra and the IGR of choice by adding the Tempo® SC Ultra first and making sure it's a well mixed dilution, then adding the correct amount of IGR according to its label instructions. Follow all label instructions for use patterns and spray intervals from the IGR label to achieve maximum benefit.

Refer to the appropriate sections of the label for Tempo® SC Ultra and the label for the IGR of choice for additional precautions, restrictions, limitations, and application information.

**PAGE 8**

W. P. Carey & Co. Inc.

**FOOD/FEED HANDLING  
ESTABLISHMENT APPLICATIONS**

Tempo® SC Ultra applications are permitted in both food/feed and non-food areas of food/feed handling establishments as a general surface, spot, or crack and crevice treatment. Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food. Serving areas where food is exposed and the facility is in operation are also considered food areas.

**General Surface Application:** When the facility is in operation or foods are exposed do not apply as a general surface application in food handling areas. Do not apply directly to food products. Cover or remove all food processing and/or handling equipment during application. After application in food processing plants, bakeries, cafeterias and similar facilities, wash all equipment, benches, shelving and other surfaces which food will contact. Clean food handling or processing equipment and thoroughly rinse with clean, fresh water.

**Spot, Crack and Crevice Application:** Spot or crack and crevice applications may be made while the facility is in operation; however, food should be covered or removed from area being treated. Do not apply directly to food or food-handling surfaces.

3 1

## INDOOR PEST CONTROL

### CRAWLING INSECT PEST CONTROL

For **general household pests** and **occasional invaders** make applications in a manner as previously described under General Surface, Spot, or Crack and Crevice Application. Particular attention should be given to treating entry points and harborage areas.

For **pantry pest control** of exposed adult and immature stages of insects, make application to cupboards, shelving and storage areas. Particular attention should be given to treating cracks and crevices and other harborage areas. Remove all utensils, uncovered foodstuffs (or any with package opened), shelf paper and other objects before spraying. Allow treated surfaces to dry and cover shelves with clean paper before replacing any utensils, foodstuff or other items. Any foodstuff accidentally contaminated with spray dilution should be discarded.

1

For **stored product pest control** of exposed adult and immature stages of insect pests, apply to cracks, crevices, and other surfaces where the pests have been seen or have harborage. Treat warehouses, production facilities, storage areas, and other areas where products are stored before filling with the product. Tempo® SC Ultra may be used by pest control operators and grain producers to treat grain storage facilities and other areas noted above for stored product pest control. Cleaning of all areas prior to use of Tempo® SC Ultra will increase levels of control. Any foodstuffs infested with pests should be removed and destroyed.

For **use by pest control operators or livestock producers in livestock housing structures (including poultry houses) and pet kennels** to control crawling and flying insect pests. Apply as a general

surface and/or a crack and crevice spray. Control will be enhanced when facilities are cleaned and interior applications are supplemented with exterior perimeter treatments. Do not make interior applications of Tempo® SC Ultra in areas of facility where animals are present. Do not make applications to animal feedstuffs or watering equipment.

### **FLYING INSECT PEST CONTROL (General Surface Spray)**

**For control of flying insects indoors** make application to surfaces as directed under Directions For Use, where the pests collect or rest. **Do not apply as a “space spray” application.**

### **OUTDOOR PEST CONTROL**

**For control of flying insects outdoors** use a 0.05% dilution and make applications to outside surfaces of buildings, porches, patios, garages and other areas where these pests have been seen or found. Use a 0.05% dilution and make pin stream applications directly to the nests of wasps, yellow-jackets and hornets. **For perimeter pest control**, make a general surface application to outside surfaces of buildings, porches, patios, garages and other adjacent areas where pests have been seen or found. To help prevent infestations of buildings by occasional invaders, treat soil, turf or other substrates adjacent to buildings, the building foundation, walls, around doors and windows, soffit areas where these pests are active and may find entrance or harborage. Apply a minimum of 8 milliliters of Tempo® SC Ultra per 1000 sq ft in sufficient water to provide adequate coverage.

For control of the **imported red fire ant**, use a 0.05% dilution of Tempo® SC Ultra and thoroughly drench or inject the entire mound area. Treat all areas of the mound showing activity to provide contact

control of the adult fire ants. Tempo® SC Ultra will provide contact activity and residual repellency in the active mound.

When using the injection treatment method for control of the imported red fire ant, it is essential that treatment be made to the bottom of the active mound. Apply dilution or sufficient volume of foam throughout the mound using a downward and upward motion of the injector tool. Multiple injection sites may be needed to ensure complete distribution of the suspension throughout the active mound.

Do not use Tempo® SC Ultra to treat fire ant mounds in any food crop production area, pasture or area where food crop production may occur.

## CONTROL OF WOOD INFESTING PESTS

For control of **wood infesting beetles** on structure surfaces, apply a 0.05% dilution of Tempo® SC Ultra. For treatment of small areas, apply by brushing or spraying the suspension evenly on wood surfaces. For large or overhead areas, apply as a coarse spray to thoroughly cover the area. Cover all surfaces below the area being sprayed with plastic sheeting or other material which can be disposed of by placing in trash if contamination from dripping occurs. Sprayed surfaces should be avoided until the treated area is completely dry.

For control of **above ground termites and wood-infesting beetles** in localized areas, apply a 0.05% dilution or sufficient volume of foam of Tempo® SC Ultra to voids and galleries in damaged wood, and in spaces between wooden structural members and between the sill plate and foundation where wood is vulnerable.

Such applications for termites are not a substitute for mechanical alteration, soil treatment or foundation treatment, but are merely a supplement. For active termite infestations, get a professional inspection.

← 2 13/16

2 13/16 →

PESTS CONTROLLED BY TEMPO® SC ULTRA*			
<b>I. INDOOR PREMISE PESTS</b>		<b>II. FOOD PROCESSING PESTS</b>	
<b>1. Crawling Pests</b>	<b>2. Flying Pests</b>	(Same crawling & flying pests, pantry & stored product pests as for INDOOR PREMISE) and Fruitflies Phoridflies and Sclaridflies	
Ants (except Pharaoh) Bedbugs Carpet beetles Centipedes Cockroaches American Asian German Oriental brown-banded smoky brown	Crickets Darkling beetles Earwigs Firebrats Millipedes Pillbugs Silverfish Sowbugs Spiders	Clothes moths Flies Gnats Midges Mosquitoes Moths Wasps Yellow-jackets	
<b>3. Pantry &amp; Stored Product Pests</b>		<b>III. WOOD INFESTING PESTS</b>	
Beetles (exposed adults & immature stages) Cadelle Cigarette Confused Flour Dermestid Drug-store Granary weevil Hide Larder Leather		Beetles Ambrosia Deathwatch False powderpost Old house borer Powderpost	
Lesser grain borer Lesser mealworm Merchant grain Mealworm Red flour Rice weevil Sawtoothed grain Warehouse		Carpenter ants Carpenter bees Termites Subterranean** Formosan Drywood	
		<b>IV. OUTDOOR/PERIMETER PESTS</b>	
		Ants Bees Boxelder bugs Carpenter ants Cecidflies Centipedes Clusterflies Cockroaches Crickets Elm leaf beetle Fire ants Firebrats Flies Fleas Gnats	
		Ground beetles Hornets Imported red fire ants Midges Millipedes Mosquitoes Moths Scorpions Silverfish Spiders Ticks Wasps Yellow-jackets	
		*For <b>GENERAL SURFACE APPLICATION</b> , use 8 milliliters of Tempo® SC Ultra per 1000 square feet in sufficient water to adequately cover the area being treated. *For <b>SPOT, CRACK AND CREVICE APPLICATIONS</b> , mix 8 milliliters Tempo® SC Ultra in one gallon of water to make a 0.025% active ingredient concentration. **Only for control of above-ground, winged forms of subterranean termites.	

19/21

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original containers and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Kansas City Emergency Response Telephone No. is 800-414-0244 or contact Chemtrec at 800-424-9300.

**Pesticide Disposal:** Pesticide wastes are hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.



2 7/8



2 1/2

**Container Disposal:** Do not use container in connection with food, feed, or drinking water. Completely empty container into application equipment. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

919882

Extended Text<sup>®</sup> INSEAL<sup>™</sup> has the following Patents: UK 0098092, Europe 0342006, USA 4488922.