

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

July 22, 2011

Douglas A. Spilker, PhD Manager, EPA Regulatory Affairs Bayer HealthCare LLC, Animal Health Division POB 390 Shawnee Mission, KS 66201

Subject:

Revise "Retreatment Statement", per Agency's Request; Add "detailed opening

instructions".

JK9 Advantix Plus 10 (EPA Registration No. 11556-141)

K9 Advantix Plus 20 (EPA Registration No. 11556-142)

K9 Advantix Plus 55 (EPA Registration No. 11556-143)

K9 Advantix Plus 100 (EPA Registration No. 11556-144)

Date of Applications: April 28, 2011

Dear Dr. Spilker:

The above referenced submissions, in connection with registrations under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, are acceptable.

Please note that we have approved the amended labeling for each product, with the understanding that it is consistent with the Agency's previously stamped accepted labels; and other than the revisions stated in your applications (dated April 28, 2011), no other changes have been made to the labeling and/or the Confidential Statement of Formula (CSF).

For your records, see enclosed copy of the stamped label for each product. If you have any questions concerning this action please contact Dr. B.A. Akinlosotu at (703) 605-0653.

Sincerely

Richard Gebken,

Product Manager 10

Insecticide Branch

Registration Division (7505P)

Revise retreatment statement; add additional Date: 04/08/11

opening instructions.

Supersedes: 09/15/10

Page 1 of 9

NOTE TO REVIEWER: [(Brackets and parentheses indicate alternate language)]

K9 Advantix[®] Plus 10 Alternate Brand Name: K9 Advantix II Small Dog

Topical prevention and treatment of fleas, ticks, mosquitoes, biting flies, and lice for monthly use on (small) dogs and puppies 7 weeks of age and older, weighing 10 lbs. and under

READ THE ENTIRE LABEL BEFORE EACH USE

DO NOT USE ON CATS

For the Prevention and Treatment of Fleas, Ticks, Biting Flies, Mosquitoes, and Lice on Dogs

[Selected optional claims bulleted here from page 8 and/or 9]

:	ACCEPTED JUL 2 2 2011 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under:	%		
Active Ingredients	•		By Weight	
Imidacloprid	PA. Reg. No:	8.	.80%	
Permethrin*			.00%	
· · · · · · · · · · · · · · · · · · ·			44%	
Other Ingredients		<u>46.</u>	76%	
	$f_{\mathcal{D}}(\pm)$ cis and min 45% (±) trans	100.	00%	
EPA Reg No. 11556-141	EPA Est. No. 11	t. No. 11556-DEU-1		
		ر ر ر د د د	•	
, I	ZEED OUR OF DEACH OF CHILDREN		3 3 3	
	WARNING)))))))	,,,,,	
For Directions for Use and other labeling; see supplemental labeling inside.			j	

PRECAUTIONARY STATEMENTS

See [below] (or) [back panel] for first aid and precautionary statements.

Revise retreatment statement; add additional opening instructions.

Date: 04/08/11

Supersedes: 09/15/10

Page 2 of 9

HAZARDS TO HUMANS

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

HAZARDS TO DOMESTIC ANIMALS

For external use on dogs only. Do not use on animals other than dogs. Do not use on puppies under seven weeks of age. Do not get this product in dog's eyes or mouth. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. If signs persist, or become more severe, consult a veterinarian immediately. If your animal is on medication, consult your veterinarian before using this or any other product.

DO NOT USE ON CATS



Due to their unique physiology and inability to metabolize certain compounds, this product must not be used on cats. If applied to a cat, or ingested by a cat which actively grooms a recently treated dog, this product may have serious harmful effects. If this occurs contact your veterinarian immediately.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not add directly to water. Do not contaminate water when disposing of product or packaging.

Revise retreatment statement; add additional opening instructions.

Date: 04/08/11 Supersedes: 09/15/10

Page 3 of 9

FIRST AID		
If In Eyes:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 	
If Swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. 	
If On Skin or Clothing:	 Do not give anything to an unconscious person. Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice. 	
HOT LINE NUMBER		
	iner or label with you when calling a poison control center or doctor, For consumer questions call 1-800-255-6826. For medical -422-9874.	
	NOTE TO PHYSICIAN	
Treat the patient sympt	omatically.	

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not contaminate feed or food.



HOW TO OPEN [Optional Language for product in blister packs]

- 1. Being careful not to cut close to the blister cavities, take scissors and cut off one section of the card containing a single tube.
- 2. Take the separated section, and cut into the blister cavity across the small side, close to the cap of the tube.
- 3. Peel off the foil, and take out the tube.
- 4. Repeat steps 1 to 3 for each tube.

Revise retreatment statement; add additional opening instructions.


Date: 04/08/11 Supersedes: 09/15/10

Page 4 of 9


HOW TO APPLY

1. Use only on dogs. DO NOT USE ON CATS or on other animals.

2. Remove applicator tube from the package. Use scissors and peel down foil.



- 3. Hold applicator tube in an upright position. Pull cap off tube.
- 4. Turn the cap around and place other end of cap back on tube.
- 5. Twist cap to break seal, then remove cap from tube.
- 6. The dog should be standing for easy application. The entire contents of the K9 Advantix® Plus tube should be applied evenly to two to three spots on the top of the back from the shoulder to the base of the tail. At each spot, part the hair until the skin is visible. Place the tip of the tube on the skin and gently squeeze to expel a portion of the solution on the skin. Do not apply an excessive amount of solution at any one spot that could cause some of the solution to run off the side of the dog.



7. Discard empty tube as described in Storage and Disposal.

60410

Reason To Issue:

Revise retreatment statement; add additional Date: 04/08/11

opening instructions.

Supersedes: 09/15/10

Page 5 of 9

8. In cases of severe pest infestation, retreatment is allowable earlier than four (4) weeks. Do not retreat more often than once every seven (7) days. After pest control is attained, return to a monthly retreatment schedule.

The successive feeding activity of fleas on dogs may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of dogs with K9 Advantix[®] Plus 10 rapidly kills fleas and may reduce the incidence of this condition.

K9 Advantix[®] Plus 10 kills existing fleas on dogs within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting at least four (4) weeks. Pre-existing pupae in the environment may continue to emerge for six (6) weeks or longer depending upon the climatic conditions.

Larval flea stages in the dog's surroundings are killed following contact with a K9 Advantix® Plus 10 treated dog.

K9 Advantix[®] Plus 10 is waterproof and remains effective following a shampoo treatment, swimming or after exposure to rain or sunlight.

Monthly treatments are required for optimal control and prevention of fleas and ticks.

Under normal conditions, the product is effective for a month. However, in case of severe pest infestation, retreatment may be necessary earlier than four (4) weeks. Do not retreat more often than once every seven (7) days. After pest control is attained, return to a monthly retreatment schedule.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry place. Protect from freezing. Disposal: Nonrefillable container. If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-422-9874 for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND LIMITATION OF DAMAGES!

Bayer HealthCare LLC, Animal Health Division warrants that this material conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR IMPLIED WARRANTY OR IMPLIED W

Revise retreatment statement; add additional Date: 04/08/11 opening instructions.

Supersedes: 09/15/10

Page 6 of 9

NET CONTENTS: XX Tube(s) – each 0.014 fl. oz. (0.4 mL)

[Sample – Not for (Re)Sale]

Manufactured For

Bayer HealthCare LLC P.O. BOX 390 Animal Health Division Shawnee Mission, Kansas 66201 USA

Made in Germany

Revise retreatment statement; add additional Date: 04/08/11 opening instructions.

Supersedes: 09/15/10

Page 7 of 9

(Label on Individual Tube)

K9 Advantix® Plus 10

8.8% imidacloprid 44.0% permethrin 0.44% pyriproxyfen

0.014 fl. oz. (0.4 mL)

EPA Reg. No. 11556-141

WARNING

Keep out of reach of children DO NOT USE ON CATS

Read The Entire Label Before Use

BAYER

Lot No. 0000000

Revise retreatment statement; add additional Date: 04/08/11 opening instructions.

Supersedes: 09/15/10

Page 8 of 9

NOTE TO REVIEWER: [(Brackets and parentheses indicate alternate language)]

OPTIONAL MARKETING CLAIMS

- For use on dogs and puppies 7 weeks of age and older
- K9 Advantix Plus contains [imidacloprid], [permethrin] and [an/the] [insect growth regulator] [IGR] [pyriproxyfen]
- A single topical application remains effective for [(a month)(four (4) weeks)]
- Convenient, easy to apply topical solution
- Once a month topical flea treatment for dogs 7 weeks of age or older
- K9 Advantix Plus is indicated for the prevention and treatment of fleas, ticks, biting flies, mosquitoes, and lice on dogs 7 weeks of age and older
- For the treatment and prevention of flea infestations
- One treatment prevents further flea infestations for [(a month)(four (4) weeks)]
- Kills fleas on dogs within 12 hours and continues to prevent infestations for [(a month)(four (4) weeks)]
- Repels and kills fleas before they lay eggs
- Larval flea stages in the pet's surroundings are killed following contact with a K9 Advantix Plus treated dog
- Kills larval stages of fleas in the dog's environment following contact with K9 Advantix Plus treated dog
- Kills fleas within 12 hours of application
- Controls existing flea infestations by rapidly killing adult fleas
- Prevents reinfestations by killing adult fleas before they lay eggs
- Prevents flea eggs from hatching
- Effectively breaks the flea life cycle
- Flea adulticide, larvicide and ovicide.
- Prevents fleas on treated dogs from infesting (reinfesting) your home.
- Effectively targets all [life] stages of [fleas] and [ticks]
- 3-way flea protection ([kills] [controls] adults, larvae, and eggs)
- Prevents development of fleas, flea eggs, pupae and larvae [for (a month)(four (4) weeks)]
- Prevents development of all flea stages [for (a month)(four (4) weeks)]
- Repels and kills all [life] stages of fleas
- Prevents flea eggs [and flea larvae] from developing into [(biting) (adult)] fleas
- Treatment with K9 Advantix Plus rapidly kills fleas and may reduce the incidence of Flea Allergic Dermatitis [FAD]
- Repels and kills ticks including Deer ticks (vector of Lyme disease). American dog ticks (vector of Rocky Mountain spotted fever), Brown dog ticks (vector of Ehrlichiosis), and Lone Star ticks (vector of Ehrlichiosis) for up to four weeks'
- Repels and kills mosquitoes for up to four weeks
- Repels and kills mosquitoes often before they have a chance to take a blood meal
- [(Prevents blood-feeding by) (Kills and repels)] mosquitoes
- Repels, and inhibits blood-feeding by biting flies
- Repels, and prevents blood-feeding by biting flies

Revise retreatment statement; add additional Date: 04/08/11 opening instructions.

Supersedes: 09/15/10

Page 9 of 9

Repels [(annoying)(bothersome)(nuisance)] biting flies

- Inhibits [(annoying)(bothersome)(nuisance)] biting flies
- [(Prevents)(inhibits)] blood-feeding by biting flies
- Remains effective after bathing and/or swimming
- Remains effective following swimming and/or shampooing
- Kills [(biting) (chewing)] lice
- For treatment and prevention of [(biting) (chewing)] lice [infestations]
- Controls existing [(biting) (chewing)] lice infestations
- 5-way protection (against fleas, ticks, biting flies, mosquitoes, and lice)
- Treats, prevents and controls [(biting) (chewing)] lice [infestations]
- Provides effective control of [(biting) (chewing)] lice [infestations]
- Kills [(biting) (chewing)] lice and prevents further infestations
- For treatment and prevention of infestations with [(biting) (chewing)] lice
- Waterproof
- Remains effective after exposure to rain or sunlight
- Fragrance free