



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

April 23, 2025

Todd Frank
Regulatory Advisor
Elanco Animal Health
2500 Innovation Way
Greenfield, IN 46140

Subject: Approval of Label Amendment; Only Indicated Changes Reviewed – Correct minor typographical errors
Product Name: Advantage II Small Cat
EPA Registration Number: 11556-151
Application Date: 01/23/2025
Case Number: 649571

Dear Mr. Frank:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. However, EPA reviewed only the label changes highlighted, marked, or otherwise indicated on the submitted label. Any other changes to the previously approved label that were not clearly highlighted, marked, or otherwise indicated in your submission were not reviewed and may form the basis of regulatory and/or enforcement action if later discovered by the Agency. Further, submission of a label amendment application with unidentified changes may be considered a knowing submission of false information to the Agency. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

The label submitted with the application has been stamped “Accepted Only Indicated Revisions Reviewed” and is enclosed for your records.

This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 C.F.R. § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently

approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 C.F.R. § 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website contains any false or misleading statement, design, or graphic, the product may be misbranded and unlawful to sell or distribute under FIFRA Sections 2(q)(1)(A) and 12(a)(1)(E). 40 C.F.R. § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on the product label, claims made as part of the product's sale or distribution may not substantially differ from those claims approved through the registration process under FIFRA Section 12(a)(1)(B). Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the product will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Scott Campbell at Campbell.Scott@epa.gov

Sincerely,

A handwritten signature in black ink, appearing to read 'Loren LaPointe', with a stylized, cursive script.

Loren LaPointe, PhD
Acting Product Manager 01
Invertebrate and Vertebrate Branch 3
Registration Division (7505T)
Office of Pesticide Programs

Enclosure

Date: 01/09/2025

Supersedes: 08/06/2024

Advantage™ II Small Cat
EPA Reg No. 11556-151

This label is divided into 3 sub-labels as follows:

Sub-label A: Cat and Ferret

Sub-label B: Cat Only

Sub-label C: Ferret Only

{OPTIONAL MARKETING CLAIMS are listed at the end of this document for Sub-label A, B and C}

{Alternate Brand Names listed at the end of this document.}

A C C E P T E D

ONLY INDICATED

REVISIONS REVIEWED

04/23/2025

Under the Federal Insecticide, Fungicide and
Rodenticide Act as amended, for the pesticide
registered under EPA Reg. No.

11556-151

No label revisions other than those indicated were
reported to the Agency.

SUB-LABEL A: Cat and Ferret Only Product

NOTE TO REVIEWER: Braces “{ }” indicate notes to reviewer. Brackets “[]” indicate optional language

{Front Panel}

For Use ONLY on Cats and Kittens 8 Weeks and Older and Weighing 5-9 lbs. and
Ferrets 10 Weeks and Older and Weighing 1 lb. or Greater

{A large, clear picture of a small cat appropriate weight range for the product will appear on the front panel.
See page 36}

Advantage II Small Cat

{Alternate Brand Names are listed at the end of this document.}

{The “Small Cat” text size will be 40% - 75% of the height of the largest letter in the primary brand name.}

{Selected optional marketing claims bulleted and pictograms from pages 34-36}

ACTIVE INGREDIENTS:	<u>% by Weight</u>
Imidacloprid	9.10%
Pyriproxyfen.	0.46%
OTHER INGREDIENTS:	<u>90.44%</u>
TOTAL	100.00%

Once-A-Month Topical Flea Prevention and Treatment for Cats and Ferrets
For Use ONLY on Cats and Kittens 8 Weeks and Older and Weighing 5-9 lbs. or Ferrets 10
weeks and Older and Weighing 1 lb. or Greater

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for [First Aid,] Precautionary Statements, and Directions for Use. For all information, including
Storage and Disposal see insert [leaflet] inside carton.

EPA Reg. No. 11556-151 EPA Est. No. XXX
{EPA Reg. No. and EPA Est. No. to appear on either front or back panel}

Lot No.
{Lot No. to appear on any panel (front, side, top, or bottom) of outer container}

{Back Panel}

Advantage II Small Cat

{The “Small Cat” text size will be 40% - 75% of the height of the largest letter in the primary brand name.}

Once-A-Month Topical Flea Prevention and Treatment for Cats and Ferrets
For Use ONLY on Cats 8 Weeks and Older and Weighing 5-9 lbs. or
Ferrets 10 weeks and Older and Weighing 1 lb. or Greater

READ THE ENTIRE LABEL BEFORE EACH USE

For the Prevention and Treatment of Flea Infestations on Cats and Ferrets

{Selected optional marketing claims bulleted from pages 34-36}

FIRST AID	
If Swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in Eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing:	• Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. [For medical emergencies or consumer questions call 888-545-5973.] [For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 888-545-5973.] {choice of either or both phone number statements will be used}	
NOTE TO PHYSICIAN	
Treat the patient symptomatically.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Harmful if on skin. Causes moderate eye irritation. Avoid contact with eyes, skin and clothing. Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not apply to cats or kittens under 8 weeks of age or weighing less than 5 lbs. or on kits (young ferrets) under 10 weeks of age or weighing less than 1 lb. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats or ferrets. Individual sensitivities may occur after using ANY pesticide product for cats or ferrets. If signs persist, or become more severe, consult a veterinarian immediately. If your cat or ferret is on medication, consult your veterinarian before using this or any other product.

{Side Effects box to go on [lower] right side of back panel.}

Side Effects (Cats): Monitor your cat after application. Side effects may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 888-545-5973.

Side Effects (Ferrets): Monitor your ferret after application. Side effects may include temporary changes in fecal consistency. If these or other side effects occur, consult your veterinarian or call 888-545-5973.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RESTRICTIONS:

- FOR EXTERNAL USE ONLY.
- Use only on cats or kittens eight [8] weeks and older and weighing 5 – 9 lbs. and only on ferrets ten [10] weeks and older and weighing 1 lb. or greater.
- DO NOT use on other animals.
- Weigh your cat or ferret before applying to ensure you are applying the correct dose for the weight of the animal.
- DO NOT apply more than one [1] tube per treatment.
- DO NOT have contact or allow children to have contact with treated area until completely dry.
- DO NOT allow animal to ingest this product.
- In multi-animal households DO NOT allow cats and/or ferrets to groom each other until the product has dried.
- DO NOT allow children to apply the product.
- Cats or ferrets should be dry prior to application and remain dry for 24 hours after application.

See inside flap for HOW TO OPEN and HOW TO APPLY or insert [leaflet] for full Directions for Use.

Net Contents: 1[2][3][4][5][6][8][10][12] Tube[s] each 0.014 fl. oz. (0.4 mL)

{one of the above Net Content quantities will be used based on the number of tubes in the carton}

Manufactured For:
Elanco US Inc
2500 Innovation Way
Greenfield, IN 46140
Product of Germany

[Sample – NOT for Sale]

{Note to reviewer: cartons containing 1 or 2, individually pouched, tubes are completely labeled (all components) as the 3, 4, 5, 6, 8, 10, and 12 tubes. The “Sample – NOT for Sale” is used as text for the 1 or 2 carton only.}

{Additionally, Inside Flap of Unit Carton for 1, 2, 3, 4, 5, 6, 8, 10 or 12 tubes}

DIRECTIONS FOR USE (continued from back panel)

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
{options of “HOW TO OPEN” and “HOW TO APPLY” will be selected based on packaging}

HOW TO OPEN



{OPTION 1: INSTRUCTIONS FOR BLISTER PACK}

1. Being careful not to cut close to the blister cavities, take scissors and cut off one section of the card containing a single tube.
2. Take the separated section and cut into the blister cavity across the small side, close to the cap of the tube.
3. Peel off the foil and take out the tube.
4. Repeat steps 1 to 3 for each tube.

[OPTION 2: INSTRUCTIONS FOR SHORT NECK TUBE POUCH]

1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.


[OPTION 3: INSTRUCTIONS FOR LONG NECK TUBE POUCH]

1. Be sure the tube is at the bottom of the pouch.
2. Use scissors to cut the pouch across the top, being careful not to cut the tube.
3. Remove the tube from the package.

HOW TO APPLY

{OPTION 1: INSTRUCTIONS FOR BLISTER PACK AND SHORT NECK TUBE POUCH}

1. Use only on cats or kittens 8 weeks and older and weighing 5 – 9 lbs. and only on ferrets 10 weeks and older and weighing 1 lb. or greater. Do not use on other animals.
2. Cats or ferrets should be dry prior to application and remain dry for 24 hours after application [for product to work effectively].
3. Remove one blister [pouch] from the carton. See “HOW TO OPEN” section.
4. Hold tube in an upright position facing away from you and your cat's or ferret's face and eyes. Pull cap off tube.



5. Turn the cap around and place other end of cap back on tube.
6. Twist cap to break seal, then remove cap from tube.
7. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. *Do not get this product in your cat's or ferret's eyes or allow your cat or ferret to ingest this product. The product is bitter tasting and salivation may occur for a short time if the cat or ferret licks the product immediately after treatment.* Treatment at the base of the skull will minimize the opportunity for the cat or ferret to lick the product. Do not allow the product to run off.




8. Discard empty tube as described in Storage and Disposal.
9. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days, not to exceed 4 weekly treatments. On ferrets, do not retreat more often than every 14 days, not to exceed 2 biweekly treatments. After flea control is attained, return to a monthly treatment schedule.

[OPTION 2: INSTRUCTIONS FOR LONG NECK TUBE POUCH]

1. Use only on cats or kittens 8 weeks and older and weighing 5 – 9 lbs. and only on ferrets 10 weeks and older and weighing 1 lb. or greater. Do not use on other animals.
2. Cats or ferrets should be dry prior to application and remain dry for 24 hours after application [for product to work effectively].
3. Remove one pouch from the carton. See “HOW TO OPEN” section.
4. Hold tube in an upright position facing away from you and your cat’s or ferret’s face and eyes. Twist the tip of the applicator to break seal.



5. In order to ensure proper application, part the hair on the neck at the base of the skull. Place the tip of the tube against the cat’s or ferret’s skin.



6. Gently squeeze the tube to expel the entire contents of the tube directly on the skin.
7. *Do not get this product in your cat’s or ferret’s eyes or allow your cat or ferret to ingest this product.* Do not allow the product to run off of the cat or ferret. The product is bitter tasting and salivation may occur for a short time if the cat or ferret licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat or ferret to lick the product.
8. Discard the empty tube as described in the Storage and Disposal section.
9. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days, not to exceed 4 weekly treatments. On ferrets, do not retreat more often than every 14 days, not to exceed 2 biweekly treatments. Once the infestation is under control, return to a monthly treatment schedule.

{Insert/Leaflet}
Advantage II Small Cat

{The “Small Cat” text size will be 40% - 75% of the height of the largest letter in the primary brand name.

Once-A-Month Topical Flea Prevention and Treatment [for Cats and Ferrets]
For Use ONLY on Cats and Kittens 8 Weeks and Older and Weighing 5-9 lbs. or Ferrets 10
Weeks and Older and Weighing 1 lb. or Greater

READ THE ENTIRE LABEL BEFORE EACH USE
For the Prevention and Treatment of Flea Infestations on Cats and Ferrets

{Selected optional marketing claims bulleted here from pages 34-36}

ACTIVE INGREDIENTS:	<u>% by Weight</u>
Imidacloprid	9.10%
Pyriproxyfen.	0.46%
OTHER INGREDIENTS:	<u>90.44%</u>
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID	
If Swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in Eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing:	• Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. [For medical emergencies or consumer questions call 888-545-5973.] [For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 888-545-5973.] {choice of either or both phone number statements will be used}	
NOTE TO PHYSICIAN	
Treat the patient symptomatically.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Harmful if on skin. Causes moderate eye irritation. Avoid contact with eyes, skin and clothing. Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not apply to cats or kittens under 8 weeks of age or weighing less than 5 lbs.

or on kits (young ferrets) under 10 weeks of age or weighing less than 1 lb. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats or ferrets. Individual sensitivities may occur after using ANY pesticide product for cats or ferrets. If signs persist, or become more severe, consult a veterinarian immediately. If your cat or ferret is on medication, consult your veterinarian before using this or any other product.

Side Effects (Cats): Monitor your cat after application. Side effects may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 888-545-5973.

Side Effects (Ferrets): Monitor your ferret after application. Side effects may include temporary changes in fecal consistency. If these or other side effects occur, consult your veterinarian or call 888-545-5973.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RESTRICTIONS:

- FOR EXTERNAL USE ONLY.
- Use only on cats or kittens eight [8] weeks and older and weighing 5 – 9 lbs. and only on ferrets ten [10] weeks and older and weighing 1 lb. or greater.
- DO NOT use on other animals.
- Weigh your cat or ferret before applying to ensure you are applying the correct dose for the weight of the animal.
- DO NOT apply more than one [1] tube per treatment.
- DO NOT have contact or allow children to have contact with treated area until completely dry.
- DO NOT allow animal to ingest this product.
- In multi-animal households DO NOT allow cats and/or ferrets to groom each other until the product has dried.
- DO NOT allow children to apply the product.
- Cats or ferrets should be dry prior to application and remain dry for 24 hours after application.

{options of “HOW TO OPEN” and “HOW TO APPLY” will be selected based on packaging}

HOW TO OPEN



{OPTION 1: INSTRUCTIONS FOR BLISTER PACK}

1. Being careful not to cut close to the blister cavities, take scissors and cut off one section of the card containing a single tube.
2. Take the separated section and cut into the blister cavity across the small side, close to the cap of the tube.
3. Peel off the foil and take out the tube.
4. Repeat steps 1 to 3 for each tube.

[OPTION 2: INSTRUCTIONS FOR SHORT NECK TUBE POUCH]

1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.


[OPTION 3: INSTRUCTIONS FOR LONG NECK TUBE POUCH]

1. Be sure the tube is at the bottom of the pouch.
2. Use scissors to cut the pouch across the top, being careful not to cut the tube.
3. Remove the tube from the package.

HOW TO APPLY

{OPTION 1: INSTRUCTIONS FOR BLISTER PACK AND SHORT NECK TUBE POUCH}

1. Use only on cats or kittens 8 weeks and older and weighing 5 – 9 lbs. and only on ferrets 10 weeks and older and weighing 1 lb. or greater. Do not use on other animals.
2. Cats or ferrets should be dry prior to application and remain dry for 24 hours after application [for product to work effectively].
3. Remove one blister [pouch] from the carton. See “HOW TO OPEN” section.
4. Hold tube in an upright position facing away from you and your cat’s or ferret’s face and eyes. Pull cap off tube.



5. Turn the cap around and place other end of cap back on tube.
6. Twist cap to break seal, then remove cap from tube.
7. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. *Do not get this product in your cat’s or ferret’s eyes or allow your cat or ferret to ingest this product. The product is bitter tasting and salivation may occur for a short time if the cat or ferret licks the product immediately after treatment.* Treatment at the base of the skull will minimize the opportunity for the cat or ferret to lick the product. Do not allow the product to run off.




8. Discard empty tube as described in Storage and Disposal.
9. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days, not to exceed 4 weekly treatments. On ferrets, do not retreat more often than every 14 days, not to exceed 2 biweekly treatments. After flea control is attained, return to a monthly treatment schedule.


[OPTION 2: INSTRUCTIONS FOR LONG NECK TUBE POUCH]

1. Use only on cats or kittens 8 weeks and older and weighing 5 – 9 lbs. and only on ferrets 10 weeks and older and weighing 1 lb. or greater. Do not use on other animals.
2. Cats or ferrets should be dry prior to application and remain dry for 24 hours after application [for product to work effectively].
3. Remove one pouch from the carton. See “HOW TO OPEN” section.

4. Hold tube in an upright position facing away from you and your cat's or ferret's face and eyes. Twist the tip of the applicator to break seal.



5. In order to ensure proper application, part the hair on the neck at the base of the skull. Place the tip of the tube against the cat's or ferret's skin.



6. Gently squeeze the tube to expel the entire contents of the tube directly on the skin.
7. *Do not get this product in your cat's or ferret's eyes or allow your cat or ferret to ingest this product.* Do not allow the product to run off of the cat or ferret. The product is bitter tasting and salivation may occur for a short time if the cat or ferret licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat or ferret to lick the product.
8. Discard the empty tube as described in the Storage and Disposal section.
9. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days, not to exceed 4 weekly treatments. On ferrets, do not retreat more often than every 14 days, not to exceed 2 biweekly treatments. After flea control is attained, return to a monthly treatment schedule.

PRODUCT INFORMATION FOR CATS

[The successive feeding activity of fleas on cats frequently elicits a hypersensitivity skin disorder known as flea allergy dermatitis (FAD) or flea bite hypersensitivity. Treatment of cats with Advantage II Small Cat kills fleas rapidly, within 12 hours, and may reduce the incidence of this condition.]

[Advantage II Small Cat kills the existing fleas on cats within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting for up to four (4) weeks.]

[Advantage II Small Cat provides multi-stage flea control effectively breaking flea life-cycle stages for lasting control (up to four weeks) of flea populations.]

Advantage II Small Cat kills adult fleas on cats quickly, within 12 hours, inhibits the development of immature flea life stages and prevents them from reaching the biting adult stage.

Advantage II Small Cat is waterproof after 24 hours and remains effective following a shampoo treatment or after exposure to rain or sunlight.

Apply monthly treatments for optimal control and prevention of fleas.

To successfully prevent and control fleas, all cats must be treated with their own dose of Advantage II Small Cat.

Apply according to label instructions, ensuring that the product is applied to the skin, not on top of the fur.

If your cat wears a decorative collar, it may be helpful to remove the collar before application, so it doesn't absorb the product.

Remember, your cat needs to be dry at the time Advantage II Small Cat is applied and remains dry 24 hours after application.

Even indoor cats need protection against fleas because you or other pets in the household may bring fleas into the home. Fleas in your cat's environment can re-infest your cat. If you have a flea infestation, consider treating your home and yard with appropriate premise products to protect against fleas.

PRODUCT INFORMATION FOR FERRETS

Advantage II Small Cat kills the existing fleas on ferrets within 24 hours. Reinfesting fleas are killed with protection against further flea infestation lasting for up to four (4) weeks.

Advantage II Small Cat provides multi-stage flea control effectively breaking flea life cycle stages for lasting control (up to four weeks) of flea populations.

Advantage II Small Cat kills adult fleas on ferrets within 24 hours, inhibits the development of immature flea life stages and prevents them from reaching the biting adult stage.

Advantage II Small Cat is waterproof after 24 hours and remains effective following a shampoo treatment or after exposure to rain or sunlight.

Apply monthly treatments for optimal control and prevention of fleas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place inaccessible to children and pets. **Pesticide Disposal and Container Handling:** Nonrefillable container. **If empty:** Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 888-545-5973 for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY, DISCLAIMER, AND LIMITATION OF DAMAGES

By using, buying or handling this product, user and/or buyer accepts the following terms. Subject to the inherent risks referred to above, and to the extent consistent with applicable law, Elanco US Inc.'s ("Elanco") sole and exclusive limited warranty is that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, when used in accordance with the label under normal use conditions. To the extent consistent with applicable law, the above exclusive express limited warranty is void if the product is used contrary to the label. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ELANCO DISCLAIMS ANY IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, AND DISCLAIMS ANY OTHER EXPRESS WARRANTY.** No agent of Elanco is authorized to make any other warranty, except in writing with a specific reference to this warranty. *To the extent consistent with applicable*

law, Elanco shall not be liable whatsoever for special, incidental or consequential damages resulting from the use or handling of this product. To the extent consistent with applicable law, the sole and exclusive remedy of the user and/or buyer for any and all losses, injuries, or damages, regardless of the legal theory, shall not exceed the purchase price paid for the product, or the replacement of product. To the extent the above disclaimer of warranties disclaims implied warranties, it may be limited or ineffective if you are a “consumer” and this product is a “consumer product,” as those terms are defined by the Magnuson Moss Act, 15 U.S.C. § 2301, in which case the duration of any implied warranties is limited to the earlier of either the applicable statute of limitation or repose, and in no case will any implied warranty extend beyond the term of the Exclusive Express Limited Warranty for the product. To the extent consistent with applicable law, some states do not allow the exclusion or limitation of incidental or consequential damages, so the above limitation or exclusion may not apply to you.

Manufactured For:
Elanco US Inc
2500 Innovation Way
Greenfield, IN 46140
[Product of Germany]

EPA Reg. No. 11556-151 [EPA Est. No. XXX]

{OPTION 1: Label on blister pack; one blister pack containing 1, 2, 3, 4, 5, 6, 8, 10 or 12 tubes}
{OPTION 2: Label on pouch containing 1 tube}

Advantage II Small Cat

For external use only on Cats and Kittens 8 weeks and older and weighing 5-9 lbs. or ferrets
10 weeks and older and weighing 1 lb. or greater

See insert [leaflet] for complete product information.

9.10% Imidacloprid
0.46% Pyriproxyfen

1[2][3][4][5][6][8][10][12] Tube[s] - 0.014 fl. oz. (0.4 mL)
{one of the above Net Content quantities will be used based on the number of tubes in the carton}

[THIS UNIT NOT FOR SALE]

{Additional label text for one tube sample package: THIS UNIT NOT FOR SALE}

EPA Reg. No. 11556-151
ELANCO
Lot No.

{Label on Individual Tube - label is colored coded to age and weight grouping}

Advantage II Small Cat
9.10% Imidacloprid
0.46% Pyriproxyfen

0.014 fl. oz. (0.4 mL)

EPA Reg. No. 11556-151

KEEP OUT OF REACH OF CHILDREN
CAUTION

Read The Entire Label Before Use

ELANCO
Lot No.

SUB-LABEL B: Cat Only Product

NOTE TO REVIEWER: Braces “{ }” indicate notes to reviewer. Brackets “[]” indicate optional language

{Front Panel}

For Use ONLY on Cats and Kittens 8 Weeks and Older and Weighing 5-9 lbs.

{A large, clear picture of a small cat appropriate weight range for the product will appear on the front panel.
See page 36}

Advantage II Small Cat

{Alternate Brand Names are listed at the end of this document.}

{The “Small Cat” text size will be 40% - 75% of the height of the largest letter in the primary brand name.}

{Selected optional marketing claims bulleted and pictograms from pages 34-36}

ACTIVE INGREDIENTS:	<u>% by Weight</u>
Imidacloprid	9.10%
Pyriproxyfen.	0.46%
OTHER INGREDIENTS:	<u>90.44%</u>
TOTAL	100.00%

Once-A-Month Topical Flea Prevention and Treatment for Cats
For Use ONLY on Cats and Kittens 8 Weeks and Older and Weighing 5-9 lbs.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for [First Aid,] Precautionary Statements, and Directions for Use. For all information, including Storage and Disposal see insert [leaflet] inside carton.

EPA Reg. No. 11556-151 EPA Est. No. XXX
{EPA Reg. No. and EPA Est. No. to appear on either front or back panel}

Lot No.
{Lot No. to appear on any panel (front, side, top, or bottom) of outer container}

{Back Panel}

Advantage II Small Cat

{The “Small Cat” text size will be 40% - 75% of the height of the largest letter in the primary brand name.}

Once-A-Month Topical Flea Prevention and Treatment for Cats
For Use ONLY on Cats and Kittens 8 Weeks and Older and Weighing 5-9 lbs.

READ THE ENTIRE LABEL BEFORE EACH USE

For the Prevention and Treatment of Flea Infestations on Cats.

{Selected optional marketing claims bulleted from pages 34-36}

FIRST AID	
If Swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in Eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing:	• Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. [For medical emergencies or consumer questions call 888-545-5973.] [For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 888-545-5973.] {choice of either or both phone number statements will be used}	
NOTE TO PHYSICIAN	
Treat the patient symptomatically.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Harmful if on skin. Causes moderate eye irritation. Avoid contact with eyes, skin and clothing. Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not apply to cats or kittens under 8 weeks of age or weighing less than 5 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats. Individual sensitivities may occur after using ANY pesticide product for cats. If signs persist, or become more severe, consult a veterinarian immediately. If your cat is on medication, consult your veterinarian before using this or any other product.

{Side Effects box to go on [lower] right side of back panel.}

Side Effects: Monitor your cat after application. Side effects may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 888-545-5973.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RESTRICTIONS:

- FOR EXTERNAL USE ONLY.
- Use only on cats or kittens eight [8] weeks and older and weighing 5 – 9 lbs.
- DO NOT use on other animals.
- Weigh your cat before applying to ensure you are applying the correct dose for the weight of your cat.
- DO NOT apply more than one [1] tube per treatment.
- DO NOT have contact or allow children to have contact with treated area until completely dry.
- Do NOT allow your cat to ingest this product.
- In multi-cat households DO NOT allow cats to groom each other until the product has dried.
- DO NOT allow children to apply the product.
- Cats or kittens should be dry prior to application and remain dry for 24 hours after application.

See inside flap for HOW TO OPEN and HOW TO APPLY or insert [leaflet] for full Directions for Use.

Net Contents: 1[2][3][4][5][6][8][10][12] Tube[s] each 0.014 fl. oz. (0.4 mL)
{one of the above Net Content quantities will be used based on the number of tubes in the carton}

Manufactured For:
Elanco US Inc
2500 Innovation Way
Greenfield, IN 46140
Product of Germany

[Sample – NOT for Sale]

{Note to reviewer: cartons containing 1 or 2, individually pouched, tubes are completely labeled (all components) as the 3, 4, 5, 6, 8, 10, and 12 tubes. The “Sample – NOT for Sale” is used as text for the 1 or 2 carton only.}

{Additionally, Inside Flap of Unit Carton for 1, 2, 3, 4, 5, 6, 8, 10 or 12 tubes}

DIRECTIONS FOR USE (continued from back panel)

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
{options of “HOW TO OPEN” and “HOW TO APPLY” will be selected based on packaging}

HOW TO OPEN



{OPTION 1: INSTRUCTIONS FOR BLISTER PACK}

1. Being careful not to cut close to the blister cavities, take scissors and cut off one section of the card containing a single tube.

2. Take the separated section and cut into the blister cavity across the small side, close to the cap of the tube.
3. Peel off the foil and take out the tube.
4. Repeat steps 1 to 3 for each tube.

[OPTION 2: INSTRUCTIONS FOR SHORT NECK TUBE POUCH]

1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.


[OPTION 3: INSTRUCTIONS FOR LONG NECK TUBE POUCH]

1. Be sure the tube is at the bottom of the pouch.
2. Use scissors to cut the pouch across the top, being careful not to cut the tube.
3. Remove the tube from the package.

HOW TO APPLY

{OPTION 1: INSTRUCTIONS FOR BLISTER PACK AND SHORT NECK TUBE POUCH}

1. Use only on cats or kittens 8 weeks and older and weighing 5 – 9 lbs. Do not use on other animals.
2. Cats should be dry prior to application and remain dry for 24 hours after application [for product to work effectively].
3. Remove one blister [pouch] from the carton. See “HOW TO OPEN” section.
4. Hold tube in an upright position facing away from you and your cat’s face and eyes. Pull cap off tube.



5. Turn the cap around and place other end of cap back on tube.
6. Twist cap to break seal, then remove cap from tube.
7. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. *Do not get this product in your cat’s eyes or allow your cat to ingest this product. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment.* Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow the product to run off.



8. Discard empty tube as described in Storage and Disposal.
9. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days, not to exceed 4 weekly treatments. After flea control is attained, return to a monthly treatment schedule.

[OPTION 2: INSTRUCTIONS FOR LONG NECK TUBE POUCH]

1. Use only on cats or kittens 8 weeks and older and weighing 5 – 9 lbs. Do not use on other animals.

2. Cats should be dry prior to application and remain dry for 24 hours after application [for product to work effectively].
3. Remove one pouch from the carton. See “HOW TO OPEN” section.
4. Hold tube in an upright position facing away from you and your cat’s face and eyes. Twist the tip of the applicator to break seal.



5. In order to ensure proper application, part the hair on the neck at the base of the skull. Place the tip of the tube against the cat’s skin.



6. Gently squeeze the tube to expel the entire contents of the tube directly on the skin.
7. *Do not get this product in your cat’s eyes or allow your cat to ingest this product.* Do not allow the product to run off of the cat. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product.
8. Discard the empty tube as described in the Storage and Disposal section.
9. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days, not to exceed 4 weekly treatments. Once the infestation is under control, return to a monthly treatment schedule.

{Insert/Leaflet}

Advantage II Small Cat

{The “Small Cat” text size will be 40% - 75% of the height of the largest letter in the primary brand name.}

Once-A-Month Topical Flea Prevention and Treatment **for** Cats
For Use ONLY on Cats and Kittens 8 Weeks and Older and Weighing 5-9 lbs.

READ THE ENTIRE LABEL BEFORE EACH USE

For the Prevention and Treatment of Flea Infestations on Cats

{Selected optional marketing claims bulleted here from pages 34-36}

ACTIVE INGREDIENTS:	<u>% by Weight</u>
Imidacloprid	9.10%
Pyriproxyfen.	0.46%
OTHER INGREDIENTS:	<u>90.44%</u>
TOTAL	100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
If Swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in Eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing:	• Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. [For medical emergencies or consumer questions call 888-545-5973.] [For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 888-545-5973.] {choice of either or both phone number statements will be used}	
NOTE TO PHYSICIAN	
Treat the patient symptomatically.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Harmful if on skin. Causes moderate eye irritation. Avoid contact with eyes, skin and clothing. Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not apply to cats or kittens under 8 weeks of age or weighing less than 5 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats. Individual sensitivities may occur after using ANY pesticide product for cats. If signs persist, or become more severe, consult a veterinarian immediately. If your cat is on medication, consult your veterinarian before using this or any other product.

Side Effects: Monitor your cat after application. Side effects may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 888-545-5973.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RESTRICTIONS:

- FOR EXTERNAL USE ONLY.
- Use only on cats or kittens eight [8] weeks and older and weighing 5 – 9 lbs.
- DO NOT use on other animals.
- Weigh your cat before applying to ensure you are applying the correct dose for the weight of your cat.

- DO NOT apply more than one [1] tube per treatment.
- DO NOT have contact or allow children to have contact with treated area until completely dry.
- DO NOT allow your cat to ingest this product.
- In multi-cat households DO NOT allow cats to groom each other until the product has dried.
- DO NOT allow children to apply the product.
- Cats or kittens should be dry prior to application and remain dry for 24 hours after application.

{options of “HOW TO OPEN” and “HOW TO APPLY” will be selected based on packaging}

HOW TO OPEN



{OPTION 1: INSTRUCTIONS FOR BLISTER PACK}

1. Being careful not to cut close to the blister cavities, take scissors and cut off one section of the card containing a single tube.
2. Take the separated section and cut into the blister cavity across the small side, close to the cap of the tube.
3. Peel off the foil and take out the tube.
4. Repeat steps 1 to 3 for each tube.

[OPTION 2: INSTRUCTIONS FOR SHORT NECK TUBE POUCH]

1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.


[OPTION 3: INSTRUCTIONS FOR LONG NECK TUBE POUCH]

1. Be sure the tube is at the bottom of the pouch.
2. Use scissors to cut the pouch across the top, being careful not to cut the tube.
3. Remove the tube from the package.

HOW TO APPLY

{OPTION 1: INSTRUCTIONS FOR BLISTER PACK AND SHORT NECK TUBE POUCH}

1. Use only on cats or kittens 8 weeks and older and weighing 5 – 9 lbs. Do not use on other animals.
2. Cats should be dry prior to application and remain dry for 24 hours after application [for product to work effectively].
3. Remove one blister [pouch] from the carton. See “HOW TO OPEN” section.
4. Hold tube in an upright position facing away from you and your cat’s face and eyes. Pull cap off tube.



5. Turn the cap around and place other end of cap back on tube.
6. Twist cap to break seal, then remove cap from tube.

7. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. *Do not get this product in your cat's eyes or allow your cat to ingest this product. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment.* Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow the product to run off.




8. Discard empty tube as described in Storage and Disposal.
9. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days, not to exceed 4 weekly treatments. After flea control is attained, return to a monthly treatment schedule.

[OPTION 2: INSTRUCTIONS FOR LONG NECK TUBE POUCH]

1. Use only on cats or kittens 8 weeks and older and weighing 5 – 9 lbs. Do not use on other animals.
2. Cats should be dry prior to application and remain dry for 24 hours after application [for product to work effectively].
3. Remove one pouch from the carton. See “HOW TO OPEN” section.
4. Hold tube in an upright position facing away from you and your cat's face and eyes. Twist the tip of the applicator to break seal.



5. In order to ensure proper application, part the hair on the neck at the base of the skull. Place the tip of the tube against the cat's skin.



6. Gently squeeze the tube to expel the entire contents of the tube directly on the skin.
7. *Do not get this product in your cat's eyes or allow your cat to ingest this product.* Do not allow the product to run off of the cat. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product.
8. Discard the empty tube as described in the Storage and Disposal section.
9. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days, not to exceed 4 weekly treatments. Once the infestation is under control, return to a monthly treatment schedule.

PRODUCT INFORMATION

[The successive feeding activity of fleas on cats frequently elicits a hypersensitivity skin disorder known as flea allergy dermatitis (FAD) or flea bite hypersensitivity. Treatment of cats with Advantage II Small Cat kills fleas and may reduce the incidence of this condition.]

[Advantage II Small Cat kills the existing fleas on cats within 12 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting for up to four (4) weeks.]

[Advantage II Small Cat provides multi-stage flea control effectively breaking flea life-cycle stages for lasting control (up to four weeks) of flea populations.]

Advantage II Small Cat kills adult fleas on cats quickly, within 12 hours, inhibits the development of immature flea life stages and prevents them from reaching the biting adult stage.

Advantage II Small Cat is waterproof after 24 hours and remains effective following a shampoo treatment or after exposure to rain or sunlight.

Apply monthly treatments for optimal control and prevention of fleas.

To successfully prevent and control fleas, all cats must be treated with their own dose of Advantage II Small Cat.

Apply according to label instructions, ensuring that the product is applied to the skin, not on top of the fur.

If your cat wears a decorative collar, it may be helpful to remove the collar before application, so it doesn't absorb the product.

Remember, your cat needs to be dry at the time Advantage II Small Cat is applied and remains dry 24 hours after application.

Even indoor cats need protection against fleas because you or other pets in the household may bring fleas into the home. Fleas in your cat's environment can re-infest your cat. If you have a flea infestation, consider treating your home and yard with appropriate premise products to protect against fleas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place inaccessible to children and pets. **Pesticide Disposal and Container Handling:** Nonrefillable container. **If empty:** Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 888-545-5973 for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY, DISCLAIMER, AND LIMITATION OF DAMAGES

By using, buying or handling this product, user and/or buyer accepts the following terms. Subject to the inherent risks referred to above, and to the extent consistent with applicable law, Elanco US Inc.'s ("Elanco") sole and exclusive limited warranty is that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, when used in accordance with the label under normal use

conditions. To the extent consistent with applicable law, the above exclusive express limited warranty is void if the product is used contrary to the label. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ELANCO DISCLAIMS ANY IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, AND DISCLAIMS ANY OTHER EXPRESS WARRANTY.** No agent of Elanco is authorized to make any other warranty, except in writing with a specific reference to this warranty. ***To the extent consistent with applicable law, Elanco shall not be liable whatsoever for special, incidental or consequential damages resulting from the use or handling of this product. To the extent consistent with applicable law, the sole and exclusive remedy of the user and/or buyer for any and all losses, injuries, or damages, regardless of the legal theory, shall not exceed the purchase price paid for the product, or the replacement of product.*** To the extent the above disclaimer of warranties disclaims implied warranties, it may be limited or ineffective if you are a “consumer” and this product is a “consumer product,” as those terms are defined by the Magnuson Moss Act, 15 U.S.C. § 2301, in which case the duration of any implied warranties is limited to the earlier of either the applicable statute of limitation or repose, and in no case will any implied warranty extend beyond the term of the Exclusive Express Limited Warranty for the product. To the extent consistent with applicable law, some states do not allow the exclusion or limitation of incidental or consequential damages, so the above limitation or exclusion may not apply to you.

Manufactured For:
Elanco US Inc
2500 Innovation Way
Greenfield, IN 46140
[Product of Germany]

EPA Reg. No. 11556-151 [EPA Est. No. XXX]

{OPTION 1: Label on blister pack; one blister pack containing 1, 2, 3, 4, 5, 6, 8, 10 or 12 tubes}

{OPTION 2: Label on pouch containing 1 tube}

Advantage II Small Cat

For external use only on cats and kittens 8 weeks and older and weighing 5-9 lbs.

See insert [leaflet] for complete product information.

9.10% Imidacloprid

0.46% Pyriproxyfen

1[2][3][4][5][6][8][10][12] Tube[s] - 0.014 fl. oz. (0.4 mL)

{one of the above Net Content quantities will be used based on the number of tubes in the carton}

[THIS UNIT NOT FOR SALE]

{Additional label text for one tube sample package: THIS UNIT NOT FOR SALE}

EPA Reg. No. 11556-151

ELANCO

Lot No.

{Label on Individual Tube – label is colored coded to age and weight grouping}

Advantage II Small Cat

9.10% Imidacloprid

0.46% Pyriproxyfen

0.014 fl. oz. (0.4 mL)

EPA Reg. No. 11556-151

KEEP OUT OF REACH OF CHILDREN

CAUTION

Read The Entire Label Before Use

ELANCO

Lot No.

SUB-LABEL C: Ferret Only Product

NOTE TO REVIEWER: Braces “{ }” indicate notes to reviewer. Brackets “[]” indicate optional language.

{Front Panel}

For Use ONLY on Ferrets 10 Weeks and Older and Weighing 1 lb. or Greater
--

{A large, clear picture of a ferret appropriate weight range for the product will appear on the front panel. See page 36}

Advantage II Ferret

{Alternate Brand Names are listed at the end of this document.}

{The “Ferret” text size will be 40% - 75% of the height of the largest letter in the primary brand name.}

{Selected optional marketing claims bulleted and pictograms from pages 34-36}

ACTIVE INGREDIENTS:	<u>% by Weight</u>
Imidacloprid	9.10%
Pyriproxyfen.	0.46%
OTHER INGREDIENTS:	90.44%
TOTAL	100.00%

Once-A-Month Topical Flea Prevention and Treatment [for Ferrets] For Use ONLY on Ferrets 10 weeks and Older and Weighing 1 lb. or Greater
--

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for [First Aid,] Precautionary Statements, and Directions for Use. For all information, including Storage and Disposal see insert [leaflet] inside carton.

EPA Reg. No. 11556-151 EPA Est. No. XXX
{EPA Reg. No. and EPA Est. No. to appear on either front or back panel}

Lot No.

{Lot No. to appear on any panel (front, side, top, or bottom) of outer container}

{Back Panel}

Advantage II Ferret

{The “Ferret” text size will be 40% - 75% of the height of the largest letter in the primary brand name.}

Once-A-Month Topical Flea Prevention and Treatment for Ferrets
For Use ONLY on Ferrets 10 weeks and Older and Weighing 1 lb. or Greater

READ THE ENTIRE LABEL BEFORE EACH USE

For the Prevention and Treatment of Flea Infestations on Ferrets

{Selected optional marketing claims bulleted from pages 34-36}

FIRST AID	
If Swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in Eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing:	• Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. [For medical emergencies or consumer questions call 888-545-5973.] [For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 888-545-5973.] {choice of either or both phone number statements will be used}	
NOTE TO PHYSICIAN	
Treat the patient symptomatically.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Harmful if on skin. Causes moderate eye irritation. Avoid contact with eyes, skin and clothing. Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not apply to kits (young ferrets) under 10 weeks of age or weighing less than 1 lb. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing ferrets. Individual sensitivities may occur after using ANY pesticide product for ferrets. If signs persist, or become more severe, consult a veterinarian immediately. If your ferret is on medication, consult your veterinarian before using this or any other product.

{Side Effects box to go on [lower] right side of back panel.}

Side Effects: Monitor your ferret after application. Side effects may include temporary changes in fecal consistency. If these or other side effects occur, consult your veterinarian or call 888-545-5973.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RESTRICTIONS:

- FOR EXTERNAL USE ONLY.
- Use only on ferrets ten [10] weeks and older and weighing 1 lb. or greater.
- DO NOT use on other animals.
- Weigh your ferret before applying to ensure you are applying the correct dose for the weight of your ferret.
- DO NOT apply more than one [1] tube per treatment.
- DO NOT have contact or allow children to have contact with treated area until completely dry.
- DO NOT allow your ferret to ingest this product.
- In multi-animal households DO NOT allow ferrets to groom each other until the product has dried.
- DO NOT allow children to apply the product.
- Ferrets should be dry prior to application and remain dry for 24 hours after application.

See inside flap for HOW TO OPEN and HOW TO APPLY or insert [leaflet] for full Directions for Use.

Net Contents: 1[2][3][4][5][6][8][10][12] Tube[s] each 0.014 fl. oz. (0.4 mL)
{one of the above Net Content quantities will be used based on the number of tubes in the carton}

Manufactured For:
Elanco US Inc
2500 Innovation Way
Greenfield, IN 46140
Product of Germany

[Sample – NOT for Sale]

{Note to reviewer: cartons containing 1 or 2, individually pouched, tubes are completely labeled (all components) as the 3, 4, 5, 6, 8, 10, and 12 tubes. The “Sample – NOT for Sale” is used as text for the 1 or 2 carton only.}

{Additionally, Inside Flap of Unit Carton for 1, 2, 3, 4, 5, 6, 8, 10 or 12 tubes}

DIRECTIONS FOR USE (continued from back panel)

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
{options of “HOW TO OPEN” and “HOW TO APPLY” will be selected based on packaging}

HOW TO OPEN



{OPTION 1: INSTRUCTIONS FOR BLISTER PACK}

1. Being careful not to cut close to the blister cavities, take scissors and cut off one section of the card containing a single tube.

2. Take the separated section and cut into the blister cavity across the small side, close to the cap of the tube.
3. Peel off the foil and take out the tube.
4. Repeat steps 1 to 3 for each tube.

[OPTION 2: INSTRUCTIONS FOR SHORT NECK TUBE POUCH]

1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.


[OPTION 3: INSTRUCTIONS FOR LONG NECK TUBE POUCH]

1. Be sure the tube is at the bottom of the pouch.
2. Use scissors to cut the pouch across the top, being careful not to cut the tube.
3. Remove the tube from the package.

HOW TO APPLY

{OPTION 1: INSTRUCTIONS FOR BLISTER PACK AND SHORT NECK TUBE POUCH}

1. Use only on ferrets 10 weeks and older and weighing 1 lb. or greater. Do not use on other animals.
2. Ferrets should be dry prior to application and remain dry for 24 hours after application [for product to work effectively].
3. Remove one blister [pouch] from the carton. See “HOW TO OPEN” section.
4. Hold applicator tube in an upright position facing away from you and your ferret’s face and eyes. Pull cap off tube.



5. Turn the cap around and place other end of cap back on tube.
6. Twist cap to break seal, then remove cap from tube.
7. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. *Do not get this product in your ferret’s eyes or allow your ferret to ingest this product. The product is bitter tasting and salivation may occur for a short time if the ferret licks the product immediately after treatment.* Treatment at the base of the skull will minimize the opportunity for the ferret to lick the product. Do not allow the product to run off.



8. Discard empty tube as described in Storage and Disposal.
9. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On ferrets, do not retreat more often than every 14 days, not to exceed 2 biweekly treatments. After flea control is attained, return to a monthly treatment schedule.

[OPTION 2: INSTRUCTIONS FOR LONG NECK TUBE POUCH]


1. Use only on ferrets 10 weeks and older and weighing 1 lb. or greater. Do not use on other

animals.

2. Ferrets should be dry prior to application and remain dry for 24 hours after application [for product to work effectively].
3. Remove one blister [pouch] from the carton. See “HOW TO OPEN” section.
4. Hold tube in an upright position facing away from you and your ferret’s face and eyes. Twist the tip of the applicator to break seal.



5. In order to ensure proper application, part the hair on the neck at the base of the skull. Place the tip of the tube against the ferret’s skin.



6. Gently squeeze the tube to expel the entire contents of the tube directly on the skin.
7. *Do not get this product in your ferret’s eyes or allow your ferret to ingest this product.* Do not allow the product to run off of the ferret. The product is bitter tasting and salivation may occur for a short time if the ferret licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the ferret to lick the product.
8. Discard the empty tube as described in the Storage and Disposal section.
9. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On ferrets, do not retreat more often than every 14 days, not to exceed 2 biweekly treatments. Once the infestation is under control, return to a monthly treatment schedule.

{Insert/Leaflet}

Advantage II Ferret

{The “Ferret” text size will be 40% - 75% of the height of the largest letter in the primary brand name.}

Once-A-Month Topical Flea Prevention and Treatment [for Ferrets]
For Use ONLY on Ferrets 10 weeks and Older and Weighing 1 lb. or Greater

READ THE ENTIRE LABEL BEFORE EACH USE

For the Prevention and Treatment of Flea Infestations on Ferrets

{Selected optional marketing claims bulleted here from pages 34-36}

ACTIVE INGREDIENTS:

% by Weight

Imidacloprid9.10%

Pyriproxyfen0.46%

OTHER INGREDIENTS:90.44%

TOTAL100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
If Swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in Eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing:	• Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. [For medical emergencies or consumer questions call 888-545-5973.] [For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 888-545-5973.] {choice of either or both phone number statements will be used}	
NOTE TO PHYSICIAN	
Treat the patient symptomatically.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Harmful if on skin. Causes moderate eye irritation. Avoid contact with eyes, skin and clothing. Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not apply to kits (young ferrets) under 10 weeks of age or weighing less than 1 lb. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing ferrets. Individual sensitivities may occur after using ANY pesticide product for ferrets. If signs persist, or become more severe, consult a veterinarian immediately. If your ferret is on medication, consult your veterinarian before using this or any other product.

Side Effects: Monitor your ferret after application. Side effects may include temporary changes in fecal consistency. If these or other side effects occur, consult your veterinarian or call 888-545-5973.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RESTRICTIONS:

- FOR EXTERNAL USE ONLY.
- Use only on ferrets ten [10] weeks and older and weighing 1 lb. or greater.
- DO NOT use on other animals.
- Weigh your ferret before applying to ensure you are applying the correct dose for the weight of your ferret.
- DO NOT apply more than one [1] tube per treatment.
- DO NOT have contact or allow children to have contact with treated area until completely dry.

- DO NOT allow your ferret to ingest this product.
- In multi-animal households DO NOT allow ferrets to groom each other until the product has dried.
- DO NOT allow children to apply the product.
- Ferrets should be dry prior to application and remain dry for 24 hours after application.

{options of “HOW TO OPEN” and “HOW TO APPLY” will be selected based on packaging}

HOW TO OPEN



{OPTION 1: INSTRUCTIONS FOR BLISTER PACK}

1. Being careful not to cut close to the blister cavities, take scissors and cut off one section of the card containing a single tube.
2. Take the separated section and cut into the blister cavity across the small side, close to the cap of the tube.
3. Peel off the foil and take out the tube.
4. Repeat steps 1 to 3 for each tube.

[OPTION 2: INSTRUCTIONS FOR SHORT NECK TUBE POUCH]

1. Be sure tube is at bottom of pouch.
2. Using scissors, cut the pouch across the top and remove tube.


[OPTION 3: INSTRUCTIONS FOR LONG NECK TUBE POUCH]

1. Be sure the tube is at the bottom of the pouch.
2. Use scissors to cut the pouch across the top, being careful not to cut the tube.
3. Remove the tube from the package.

HOW TO APPLY

{OPTION 1: INSTRUCTIONS FOR BLISTER PACK AND SHORT NECK TUBE POUCH}

1. Use only on ferrets 10 weeks and older and weighing 1 lb. or greater. Do not use on other animals.
2. Ferrets should be dry prior to application and remain dry for 24 hours after application [for product to work effectively].
3. Remove one blister [pouch] from the carton. See “HOW TO OPEN” section.
4. Hold tube in an upright position facing away from you and your ferret’s face and eyes. Pull cap off tube.



5. Turn the cap around and place other end of cap back on tube.
6. Twist cap to break seal, then remove cap from tube.
7. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. *Do not get this product in your ferret’s eyes or allow your ferret to ingest this product. The product is bitter tasting and salivation may occur for a short time if the ferret licks the product immediately after treatment.* Treatment at the base of the skull will minimize the opportunity for the ferret to lick

the product. Do not allow the product to run off.




8. Discard empty tube as described in Storage and Disposal.
9. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On ferrets, do not retreat more often than every 14 days, not to exceed 2 biweekly treatments. After flea control is attained, return to a monthly treatment schedule.

[OPTION 2: INSTRUCTIONS FOR LONG NECK TUBE POUCH]

1. Use only on ferrets 10 weeks and older and weighing 1 lb. or greater. Do not use on other animals.
2. Ferrets should be dry prior to application and remain dry for 24 hours after application [for product to work effectively].
3. Remove one pouch from the carton. See “HOW TO OPEN” section.
4. Hold tube in an upright position facing away from you and your ferret’s face and eyes. Twist the tip of the applicator to break seal.



5. In order to ensure proper application, part the hair on the neck at the base of the skull. Place the tip of the tube against the ferret’s skin.



6. Gently squeeze the tube to expel the entire contents of the tube directly on the skin.
7. *Do not get this product in your ferret’s eyes or allow your ferret to ingest this product.* Do not allow the product to run off of the ferret. The product is bitter tasting and salivation may occur for a short time if the ferret licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the ferret to lick the product.
8. Discard the empty tube as described in the Storage and Disposal section.
9. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On ferrets, do not retreat more often than every 14 days, not to exceed 2 biweekly treatments. Once the infestation is under control, return to a monthly treatment schedule.

PRODUCT INFORMATION

Advantage II Ferret kills the existing fleas on ferrets within 24 hours. Reinfesting fleas are killed with protection against further flea infestation lasting for up to four (4) weeks.

Advantage II Ferret provides multi-stage flea control effectively breaking flea life cycle stages for lasting

control (up to four weeks) of flea populations.

Advantage II Ferret kills adult fleas within 24 hours, inhibits the development of immature flea life stages and prevents them from reaching the biting adult stage.

Advantage II Ferret is waterproof after 24 hours and remains effective following a shampoo treatment or after exposure to rain or sunlight.

Apply monthly treatments for optimal control and prevention of fleas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place inaccessible to children and pets. **Pesticide Disposal and Container Handling:** Nonrefillable container. **If empty:** Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 888-545-5973 for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY, DISCLAIMER, AND LIMITATION OF DAMAGES

By using, buying or handling this product, user and/or buyer accepts the following terms. Subject to the inherent risks referred to above, and to the extent consistent with applicable law, Elanco US Inc.'s ("Elanco") sole and exclusive limited warranty is that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, when used in accordance with the label under normal use conditions. To the extent consistent with applicable law, the above exclusive express limited warranty is void if the product is used contrary to the label. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ELANCO DISCLAIMS ANY IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, AND DISCLAIMS ANY OTHER EXPRESS WARRANTY.** No agent of Elanco is authorized to make any other warranty, except in writing with a specific reference to this warranty. ***To the extent consistent with applicable law, Elanco shall not be liable whatsoever for special, incidental or consequential damages resulting from the use or handling of this product. To the extent consistent with applicable law, the sole and exclusive remedy of the user and/or buyer for any and all losses, injuries, or damages, regardless of the legal theory, shall not exceed the purchase price paid for the product, or the replacement of product.*** To the extent the above disclaimer of warranties disclaims implied warranties, it may be limited or ineffective if you are a "consumer" and this product is a "consumer product," as those terms are defined by the Magnuson Moss Act, 15 U.S.C. § 2301, in which case the duration of any implied warranties is limited to the earlier of either the applicable statute of limitation or repose, and in no case will any implied warranty extend beyond the term of the Exclusive Express Limited Warranty for the product. To the extent consistent with applicable law, some states do not allow the exclusion or limitation of incidental or consequential damages, so the above limitation or exclusion may not apply to you.

Manufactured For:
Elanco US Inc
2500 Innovation Way
Greenfield, IN 46140
[Product of Germany]

EPA Reg. No. 11556-151 [EPA Est. No. XXX]

{OPTION 1: Label on blister pack; one blister pack containing 1, 2, 3, 4, 5, 6, 8, 10 or 12 tubes}

{OPTION 2: Label on pouch containing 1 tube}

Advantage II Ferret

For external use only **on** ferrets 10 weeks and older and weighing 1 lb. or greater

See insert [leaflet] for complete product information.

9.10% Imidacloprid

0.46% Pyriproxyfen

1[2][3][4][5][6][8][10][12] Tube[s] - 0.014 fl. oz. (0.4 mL)

{one of the above Net Content quantities will be used based on the number of tubes in the carton}

[THIS UNIT NOT FOR SALE]

{Additional label text for one tube sample package: THIS UNIT NOT FOR SALE}

EPA Reg. No. 11556-151

ELANCO

Lot No.

{Label on Individual Tube - label is colored coded to age and weight grouping}

Advantage II Ferret

9.10% Imidacloprid

0.46% Pyriproxyfen

0.014 fl. oz. (0.4 mL)

EPA Reg. No. 11556-151

KEEP OUT OF REACH OF CHILDREN

CAUTION

Read The Entire Label Before Use

ELANCO

Lot No.

{Braces “{ }” indicate notes to reviewer. Brackets “[]” indicate optional language. *All claims apply to all sub-labels (A, B and C) however only language applicable to the species indicated in the sub-label will be used.*}

{Optional marketing claims appearing on any location for Sublabel A, B, or C:}

1. Advantage II [Small Cat] [Ferret] Contains imidacloprid and an insect growth regulator
2. Advantage II [Small Cat] [Ferret] Contains imidacloprid and pyriproxyfen
3. Advantage II [Small Cat] [Ferret] Contains imidacloprid and the insect growth regulator [IGR] pyriproxyfen
4. A single topical application remains effective for up to 4 weeks [a month]
5. Convenient, easy-to-apply [and fragrance free] [monthly] topical solution
6. One treatment prevents further flea infestations for up to 4 weeks [a month]
7. Stops existing flea infestations by killing adult fleas
8. Prevents reinfestations by killing adult fleas, so they can't lay eggs
9. Prevents flea eggs from hatching [into biting adults]
10. Effectively breaks the flea life cycle
11. Kills [Controls] flea life stages
12. Flea prevention and treatment – kills adult fleas, flea larvae, and flea eggs
13. 3-way protection against fleas kills [controls] adults, larvae, and eggs
14. Prevents flea eggs from developing into [biting] [adult] fleas
15. Flea adulticide, larvicide, and ovicide
16. Kills fleas [flea eggs] [and flea larvae]
17. Kills fleas and their eggs
18. Advantage II [Small Cat] [Ferret] May be used year-round for flea prevention
19. Contains an insect growth regulator (IGR) to kill flea eggs and prevent reinfestation
20. Remains effective after bathing
21. Remains effective following shampooing
22. Waterproof, after 24 hours
23. Remains effective after exposure to rain or sunlight
24. Fragrance-free
25. In child-resistant packaging
26. Starts working through contact. [Fleas do not have to bite to die.]
27. Kills fleas which may cause Flea Allergy Dermatitis and flea bite anemia
28. 1[2][3][4][5][6][8][10][12] monthly doses

{Optional marketing claims for long neck applicator:}

29. Long-neck applicator
30. Long-neck makes it easy to get through fur to skin
31. Applicator makes it easy to get through long or thick fur and direct to skin

{Optional marketing claims appearing on any location for Sublabel A Cat & Ferret:}

1. For use on cats and kittens 8 weeks of age and older and weighing 5 – 9 lbs. [For use on ferrets (and kits) 10 weeks of age and older and weighing 1 lb. or greater]
2. Once a month topical flea prevention and treatment for cats and kittens 8 weeks of age or older and weighing 5 – 9 lbs. [and ferrets 10 weeks of age and older and weighing 1 lb. or greater]
3. Advantage II Small Cat is for the prevention and treatment of fleas on cats and kittens 8 weeks of age and older and weighing 5 – 9 lbs. [and ferrets 10 weeks of age and older and weighing 1 lb. or greater]
4. For the prevention and treatment of flea infestations on cats [ferrets]
5. Kills fleas within 12 hours on cats [, 24 hours on ferrets,] and continues to prevent infestations for up to four weeks [a month]
6. Kills fleas on cats within 12 hours [and on ferrets within 24 hours] of application
7. Controls flea problems on cats [ferrets]
8. Controls existing flea infestations on your cat [or ferret] and prevents further infestations

9. Prevents [flea] reinfestations on cats [ferrets]
10. Protects cats [ferrets] against fleas [flea infestations]
11. Kills fleas that may cause transmission of tapeworms [a tapeworm infestation] in cats [ferrets]
12. Prevents fleas on treated cats [or ferrets] from reinfesting
13. Prevents [establishment of] new flea infestations on cats [ferrets] after application
14. Protection from fleas for your cat [ferret] when applied monthly
15. Kills fleas even if your cat gets wet
16. Treatment of cats with Advantage II Small Cat kills fleas which may cause flea allergic dermatitis [FAD] or flea bite hypersensitivity
17. Monthly use of Advantage II Small Cat on cats kills fleas which may cause flea allergy dermatitis [flea bite hypersensitivity]
18. Reinfesting fleas on cats are killed within 2 hours providing protection against further flea infestation
19. Cats or Kittens should be dry prior to application and remain dry for 24 hours after application [for product to work effectively]

{Optional marketing claims appearing on any location for Sublabel B Cat only:}

1. For use on cats and kittens 8 weeks of age and older and weighing 5 – 9 lbs.
2. Once a month topical flea prevention and treatment for cats and kittens 8 weeks of age or older and weighing 5 – 9 lbs.
3. Advantage II Small Cat is for the prevention and treatment of fleas on cats and kittens 8 weeks of age and older and weighing 5 – 9 lbs.
4. For the prevention and treatment of flea infestations on cats
5. Kills fleas within 12 hours on cats and continues to prevent infestations for up to four weeks [a month]
6. Kills fleas on cats within 12 hours of application
7. Controls flea problems on cats
8. Controls existing flea infestations on your cat and prevents further infestations
9. Prevents [flea] reinfestations on cats
10. Protects cats against fleas [flea infestations]
11. Kills fleas that may cause transmission of tapeworms [a tapeworm infestation] in cats
12. Prevents fleas on treated cats from reinfesting
13. Prevents [establishment of] new flea infestations on cats after application
14. Protection from fleas for your cat when applied monthly
15. Kills fleas even if your cat gets wet
16. Treatment of cats with Advantage II Small Cat kills fleas which may cause flea allergic dermatitis [FAD] or flea bite hypersensitivity
17. Monthly use of Advantage II Small Cat on cats kills fleas which may cause flea allergy dermatitis [flea bite hypersensitivity]
18. Reinfesting fleas on cats are killed within 2 hours providing protection against further flea infestation
19. Cats or Kittens should be dry prior to application and remain dry for 24 hours after application [for product to work effectively]

{Optional marketing claims appearing on any location for Sublabel C Ferret only:}

1. For use on ferrets (and kits) 10 weeks of age and older and weighing 1 lb. or greater
2. Once a month topical flea prevention and treatment for ferrets 10 weeks of age and older and weighing 1 lb. or greater
3. Advantage II Ferret is for the prevention and treatment of fleas on ferrets 10 weeks of age and older and weighing 1 lb. or greater
4. For the prevention and treatment of flea infestations on ferrets
5. Kills fleas within 24 hours on ferrets, and continues to prevent infestations for up to four weeks [a month]
6. Kills fleas on ferrets within 24 hours of application
7. Controls flea problems on ferrets

8. Controls existing flea infestations on your ferret and prevents further infestations
9. Prevents [flea] reinfestations on ferrets
10. Protects ferrets against fleas [flea infestations]
11. Kills fleas that may cause transmission of tapeworms [a tapeworm infestation] in ferrets
12. Prevents fleas on treated ferrets from reinfesting
13. Prevents [establishment of] new flea infestations on ferrets after application
14. Protection from fleas for your ferret when applied monthly

{Optional pictograms which may appear on the label}

{image of short neck tube will only appear with the instructions for the short neck tube}



Fleas



Flea Larvae



Flea Eggs



Waterproof



{Alternate Brand Names}


- | | |
|---|--|
| 1. Advantage™ II Plus Small Cat* | 3. Advantage™ II Advanced Small Cat |
| *Contains imidacloprid and pyriproxyfen | |
| 2. Advantage™ II Max Small Cat** | 4. Advantage™ II Advanced Care Small Cat |
| ** Contains imidacloprid and pyriproxyfen | |
| as compared to Advantage for Small Cats | 5. Advantage™ II Protect Small Cat |
| containing only imidacloprid. | |

{Alternate Brand Names Sub-Label C only}

- | | |
|---|---------------------------------------|
| 1. Advantage™ II Ferret | 3. Advantage™ II Advanced Ferret |
| 2. Advantage™ II Plus Ferret* | 4. Advantage™ II Advanced Care Ferret |
| *Contains imidacloprid and pyriproxyfen | 5. Advantage™ II Protect Ferret |

{Representative cat; each outer carton will have a representative picture of a cat in the appropriate age and weight range for the product.}

{Representative Cat picture optional on insert/leaflet, blister pack, pouch, tube.}



{Representative ferret; each outer carton will have a representative picture of a ferret in the appropriate age and weight range for the product.}

{Representative ferret picture optional on insert/leaflet, blister pack, pouch, tube.}

