
Controlling cross-contamination is of major housekeeping concern not only in hospitals, but in schools, institutions, and industry. STERISOL DISINFECTANT CLEANER is formulated for this problem area. It both cleans and disinfects effectively when used as directed. Its hard surface disinfecting action will reduce the hazard of cross-contamination.

WARNING

Keep out of reach of children. Causes skin irritation. May cause eye damage. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, give milk, raw egg white, or gelatin solution. Call a physician immediately.



CLEANER-DISIN FOR HOSP

AOAC PHENOL COEFFICIE Staph. aureau (ATCC No. 65; Salmonella typhosa (ATCC N


ACTIVE INGREDIENTS

Didecyl dimethyl ammonium
Tetrasodium ethylenediamine
Sodium carbonate
Sodium metasilicate anhydrou
Inert Ingredients

EPA REGISTRATION NO. 1

Keep out of reach of chil ary statements.

CON Manu



STERISOL

CLEANER-DISINFECTANT-DEODORIZER-FUNGICIDE FOR HOSPITAL AND INSTITUTIONAL USE

HENOL COEFFICIENTS	
ireau (ATCC No. 6538)	100
la typhosa (ATCC No. 6539)	43.8
INGREDIENTS	
dimethyl ammonium chloride	4.25%
ium ethylenediamine tetraacetate	1.60%
carbonate	2.00%
netasilicate anhydrous	0.50%
redients	91.65%
	100.00%

GISTRATION NO. 11559-1

WARNING

ut of reach of children. See left panel for additional precautionements.

CONTENTS ONE GALLON Manufactured By SV Chemicals

Ta ora, Aashington

USE DIRECTIONS

Add two ounces per gallon water.

Apply STERISOL DISINFECTANT CLEANER to walls, floors, and other hard surfaces or mop. For heavily soiled areas, a preliminary cleaning may be required.

At two ounces per gallon use-level, STERISOL DISINFECTANT CLEANER is effect pseudomonas aeruginosa, staphylococcus aureaus, salmonella choleraesuis, and trichople digitale. Germicidal performance against the first three organisms has been confirmed by Use-Dilution Test. Fungicidal performance against T. interdigitale was determined by Fungicidal Test.

Rinse empty container with water before discarding it.