


11561-13

2-2-1988

1/5

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460	EPA REGISTRATION NO. 11561-13 TERM OF ISSUANCE	DATE OF ISSUANCE 2 FEB 1988
NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i>	NAME OF PESTICIDE PRODUCT Warner Enterprises General Purpose Insecticide	
NAME AND ADDRESS OF REGISTRANT (Include ZIP code) <div style="display: flex; justify-content: space-between;"> <div> <input type="checkbox"/> Warner Enterprises P.O. Box 621 Ranson, WV 25433 </div> <div style="text-align: center;"> <div style="border: 1px solid black; padding: 10px; transform: rotate(45deg); display: inline-block;"> 161 181 </div> </div> </div>		
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.		
On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.		
A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.		
Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.		
This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:		
1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.		
2. Add the phrase "EPA Registration No. 11561-13" to your label before you release the product for shipment.		
3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.		
If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.		
A stamped copy of the label is enclosed for your records.		
<div style="text-align: right;"> Arturo N. Castillo Product Manager (17) Insecticide-Rodenticide Branch Registration Division (TS-767C) </div>		
Enclosures		
<input type="checkbox"/> ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL 		DATE 2/2/88



WARNER ENTERPRISES

KEEP FROM FREEZING.

2/5

GENERAL PURPOSE INSECTICIDE

A Concentrated PYRENONE® Micro Emulsion - Clear-Stable - Leaves No Oily or Objectionable Residues

Effective, Fast Acting Insecticide

Kills and Repels Biting Flies

For Use in Homes, Institutions, Restaurants

Food Processing Plants, Dairies, Livestock and Outdoors for Mosquito Control

For Control of Fleas and Brown Dog Ticks on premises and on pets

Apply with Conventional Mechanical or Compressed Air Equipment

ACTIVE INGREDIENTS:

	BY WEIGHT
Pyrethrins.....	0.1%
*Piperonyl Butoxide, Technical.....	1.0%
INERT INGREDIENTS:.....	98.9%
TOTAL.....	100.0%

*Equivalent to min. 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

Contains petroleum distillates.

EPA Reg. No. 11561-RG

EPA Est. No. 72-PA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

CAUCIÓN

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.
- IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.
- IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Manufactured For
WARNER ENTERPRISES
Ranson, West Virginia 25438

NET CONTENTS: 8 OUNCES through 1 GALLON LIQUID

ACCEPTED
with COMMENTS
in EPA 10000 Label
2 FEB 1988

11561-13

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS: CAUTION--Harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove pets, birds and cover fish aquariums before space spraying or surface application.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, ponds, streams, tidal marshes or estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a cool dry place inaccessible to children and pets. Protect from freezing.

DISPOSAL: Wrap container in several layers of newspaper and discard in trash.

DIRECTIONS

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned with potable water before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended.

SURFACE SPRAYING: To control cockroaches, palmetto bugs, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites and cadellles: Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers, behind and beneath sinks, stoves, refrigerators and cabinets, along baseboards, around door and window frames, into dark corners and in and around garbage compactors, garbage chutes and cans. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet or surface.

In case of infested foods in cabinets, dispose of the infested material. Remove all contents and clean thoroughly. Spray interior surfaces of empty cabinets thoroughly. After spray has dried, cover shelves with paper or otherwise protect foods from direct contact with sprayed surfaces.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moulding and floors. Repeat treatment as needed.

PET AREAS: To control fleas and brown dog ticks on premises: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moulding, window and door frames and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

SPACE SPRAYING: Close doors, windows and transoms and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of house flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths and tobacco moths: Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1,000 cubic feet of space. Do not remain in treated areas. If insects have not dropped to floor in 3 minutes, repeat spraying as quantity sprayed was insufficient. Ventilate the area after treatment is completed. To aid in the prevention of breeding, spray on standing water and in garbage pails.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

ON LIVESTOCK: To protect cattle, horses and hogs from horn flies, house flies, mosquitoes and gnats apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies and deer flies, apply at a rate of 2 ounces per adult animal sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

To control poultry lice, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For control of bedbugs and mites in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

FLEAS AND TICKS ON DOGS AND CATS: Start spraying at the animal's head - avoiding the eyes, nose and mouth as you spray. Make sure that the animal's entire body is covered, including the legs, under body and tail. While spraying, fluff the hair so that spray will penetrate to the skin. Make sure spray wets thoroughly. Avoid contact with genitalia. Repeat as necessary.

OUTDOOR APPLICATIONS

ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest (low, marshy places, stagnant pools or ponds and shore lines). Apply at a rate of 2.5 pints per acre (0.1 ounce per 1,000 square feet).

Using truck-mounted conventional misting equipment such as Rotomist® models 51C, 91 or 100 HT traveling at 5 m.p.h. to treat a 300 foot swath calibrate equipment to deliver 0.9 gallon per minute.

OUTDOORS: Camping, fishing, hunting. Spray outer clothing and surroundings. Repels biting flies, mosquitoes, midges, gnats -- if sprayed every hour or so will aid in keeping surroundings free from mosquitoes. If in rare cases skin irritation results, avoid continued contact. Do not pour or squirt - use a good sprayer which will throw a heavy fine mist.

PYRENONE - Registered Trademark of Fairfield American Corporation

LIMITED WARRANTY: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.