


11561-15

3-7-1988

1/4

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460		EPA REGISTRATION 11561-15	DATE OF ISSUANCE 7 MAR 1988
NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)		NAME OF PESTICIDE PRODUCT Warner Enterprises Tomato & Vegetable Spray	
NAME AND ADDRESS OF REGISTRANT (Include ZIP code) Warner Enterprises P.O. Box 621 Ranson, WV 25438 <div style="position: absolute; top: 150px; right: 100px; font-size: 2em; transform: rotate(-15deg);">161/2161 434 18/1</div>			
<p>NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p> <p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p> <p>Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none">1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.2. Add the phrase "EPA Registration No. 11561-15" to your label before you release the product for shipment.3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-70 Enclosure for a further description of final printed labeling. <p>If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 3(c). Your release for shipment of the product constitutes acceptance of these conditions.</p> <p>A stamped copy of the label is enclosed for your records.</p> <div style="text-align: right; margin-top: 20px;">Arturo E. Castillo Product Manager (17) Insecticide-Rodenticide Branch Registration Division (TS-767C)</div> <p>Enclosures</p> <p><input type="checkbox"/> ATTACHMENT IS APPLICABLE</p> <p>SIGNATURE OF APPROVING OFFICIAL </p> <p>DATE 3/7/88</p>			



WARNER ENTERPRISES

KEEP FROM FREEZING. 2/4

TOMATO & VEGETABLE SPRAY

KILLS GYPSY MOTH CATERPILLARS AND JAPANESE BEETLES.

AN EFFECTIVE INSECTICIDE FOR USE ON VEGETABLE AND ORNAMENTAL PESTS.

A CONCENTRATED PYRENONE® MICRO-EMULSION, CLEAR, STABLE.

APPLY WITH CONVENTIONAL MECHANICAL COMPRESSED AIR EQUIPMENT OR NON-AEROSOL PUMP.

ACTIVE INGREDIENTS:	BY WEIGHT
Pyrethrins.....	0.02%
*Piperonyl Butoxide, Technical.....	0.20%
INERT INGREDIENTS:.....	99.78%
TOTAL.....	100.00%

*Equivalent to min. 0.16% (butylcarbityl) (6-propylpiperonyl) ether and 0.04% related compounds.

PYRENONE - Registered trademark of Fairfield American Corporation.

EPA Reg. No. 11561-

EPA Est. No. 72-PA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

CAUCIÓN

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.
- IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.
- IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Manufactured For
WARNER ENTERPRISES
Ranson, West Virginia 25438

NET CONTENTS: 8 OUNCES through 1 GALLON LIQUID

ACCEPTED
7 MAR 1988

11561-15

3/4

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS: CAUTION--Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning or equipment or disposal of wastes. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in tightly closed, original container in a locked area away from children and domestic animals. Do not reuse empty container. If freezing occurs, thaw and agitate vigorously. FOR DISPOSAL, securely wrap original container in several layers of newspaper and discard in trash.

DIRECTIONS

TO OPERATE: SHAKE BEFORE USING. Hold sprayer 18 inches from foliage and spray with circular motion. Direct spray so that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when air is still.

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects names, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest.

Use on the following crops: Asparagus, Beans, Broccoli, Cauliflower, Cabbage, Brussels Sprouts, Celery, Cranberries, Eggplant, Lettuce, Mustard Greens, Kale, Collards, Turnips, Peppers, Potatoes, Radishes, Spinach, and Tomatoes. To Control: asparagus beetles, aphids, leafhoppers, whiteflies, Mexican bean beetles, 12-Spotted cucumber beetles, cabbage loopers, cross-striped cabbageworms, diamondback moth larvae, flea beetles, harlequin bugs, imported cabbageworms, stink bugs, green peach aphids, leaf tiers, fireworms, blister beetles, Colorado potato beetles, and webworms.

Use on the following ornamentals: African Violets, Azaleas, Begonias, Camellias, Dahlias, Dogwood, Roses, Wandering Jew, Asters, Carnations, Chrysanthemums, Geraniums, Gladiolus and Marigolds. To Control: aphids, flea beetles, leafhoppers and whiteflies.

To kill gypsy moth caterpillars and adults, adult Japanese beetles and Eastern and forest tent caterpillars on contact:

47/4

On Trees and Shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

INDOORS AS A SPACE SPRAY

To control house flies, fruit flies, mosquitoes, gnats and small flying moths: apply using a good space sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure filling the room with mist. Use 4 to 5 ounces per 1,000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

OUTDOORS, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at a rate of 5 to 6 quarts per acre (4 ounces per 1,000 square feet).

To control mosquito larvae, apply at the rate of 7.75 gallons per acre or .18 gallons per 1,000 square feet. Apply as a coarse wetting spray to the border of, or on shallow waters of wetlands, swamps and marshes where mosquito larvae are known to breed.

LIMITED WARRANTY: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.