



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

February 4, 2022

Lisa Setliff
Regulatory Agent for OAT Agrio Co., Ltd.
OAT AGRIO CO., LTD.
c/o Landis International, Inc.
PO Box 5126 Valdosta, GA 31603-5126

Subject: PRIA Label Amendment – R350 Amendment to Add Aerial Application
Product Name: GATTEN
EPA Registration Number: 11581-6
Application Date: 3/10/2021
Decision Number: 572373

Dear Lisa Setliff:

The application referred to above, submitted under the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable under FIFRA sec 3 (c)(5). You must submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 11581-6
Decision No. 572373

with FIFRA section 6. If you have any questions, please contact Carmen Swinger by phone at 202-566-2923, or via email at swinger.carmen@epa.gov.




Nathan Mellor, Product Manager 21
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

GATTEN®

[Alternate Brand Name: Gatten® Fungicide, Gatten® CA Fungicide]



A fungicide for control of powdery mildew.

Active Ingredient: Flutianil†‡.....4.7%
 Other Ingredients:.....95.3%
 Total 100.0%
 0.423 lbs flutianil/gal

®Trademark of OAT Agrio Co., Ltd.

†(2Z)-2-[[2-fluoro-5-(trifluoromethyl)phenyl]thio]-2-[3-(2-methoxyphenyl)-2-thiazolidinylidene]acetonitrile
 ‡ [CAS No. 958647-10-4]

**KEEP OUT OF REACH OF CHILDREN
 WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

First Aid	
If in eyes:	– Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. – Call a poison control center or doctor for treatment advice.
If swallowed:	– Call a poison control center or doctor immediately for treatment advice. – Have person sip a glass of water if able to swallow. – DO NOT induce vomiting unless told to do so by a poison control center or doctor. – DO NOT give anything to an unconscious person.
If on skin or clothing:	– Take off contaminated clothing. – Rinse skin immediately with plenty of water for 15-20 minutes. – Call a poison control center or doctor for treatment advice.
If inhaled:	– Move person to fresh air. – If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. – Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For non-emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 (NPIC Web site: www.npic.orst.edu)	
See Additional Precautionary Statements [on/in side panel, back panel, booklet]	
Manufactured [by/for]: OAT Agrio Co., Ltd.	
1-3-1 Kanda Ogawa-machi, Chiyoda-ku Tokyo 101-0052, Japan	c/o Landis International, Inc. PO Box 5126 Valdosta, GA 31603-5126 Phone 1-800-526-3471

EPA Registration Number: 11581-6 EPA Establishment Number: 11581-JPN-2
 NET CONTENTS: _____gallons/fluid ounces (_____Liters/Milliliters)
 (Note to Reviewer: Optional text in [brackets])

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING/AVISO: Causes substantial but temporary eye injury. **DO NOT** get in eyes or on clothing. Wear protective eyewear (goggles, face shield, safety glasses). Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Mixers, loaders, and other handlers must wear the following:

- Long-sleeved shirt and long pants
- Socks
- Shoes
- Chemical-resistant gloves made of barrier laminate or butyl rubber ≥ 14 mils.
- Protective Eyewear (goggles, face shield, or safety glasses)

User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users must remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users must remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For terrestrial uses: **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, or pets either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR par 170. This Standard contains requirements for the protection of agricultural workers on farms, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow workers to enter treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated including plants, soil, or water), is:

- Coveralls worn over short-sleeved shirt and short pants
- Socks
- Shoes
- Chemical-resistant gloves made of barrier laminate or butyl rubber ≥ 14 mils.
- Protective Eyewear

Product Information

GATTEN® is a fungicide for control of powdery mildew on crops listed in the **Directions for Application** section of this label. GATTEN® may be used in programs that are compatible with the principles of Integrated Pest Management (IPM), which include the use of disease resistant crop varieties, cultural practices, pest scouting and disease forecasting systems which reduce unnecessary applications of pesticides.

Flutianil, the active ingredient in GATTEN®, is a new chemical class of fungicide of the thiazolidine group and belongs to FRAC mode of action group U13. GATTEN® is effective for strategic use in programs that attempt to minimize disease resistance to fungicides.

This product is intended for use by professional applicators.

Directions for Ground or Aerial Spray Application

Users must read, understand, and follow the label use rates and restrictions. Minimum label rates may be used under low disease pressure conditions while maximum label rates and shortened specified spray intervals are advised under high disease pressure. For application, determine the number of acres to be treated, the specified label use rate, and the spray volume per acre. Prepare only the amount of spray solution that is necessary to spray the measured acres. Calibrate spray equipment prior to use.

For optimal disease control, use thorough spray coverage. Thorough spray coverage is a function of spray pressure, spray volume per acre, nozzle type and spacing and application equipment speed. Calibrate spray equipment prior to use.

MANDATORY SPRAY DRIFT REQUIREMENTS

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use $\frac{1}{2}$ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Ground Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.

- **DO NOT** apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.”

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control.

While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size- Ground Boom

- Volume- Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure- Use the lowest spray pressure specified for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle- Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size- Aircraft

- Adjust Nozzles- Follow nozzle manufacturer’s directions for setting up nozzles. To reduce fine droplets, nozzles must be oriented parallel with the airflow in flight.

BOOM HEIGHT - Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom must remain level with the crop and have minimal bounce.

RELEASE HEIGHT- Aircraft

Higher release heights increase the potential for spray drift. When applying aerially to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind

conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-less Ground Applications:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

Take precautions to minimize spray drift.

Restrictions

NEW YORK: Not for Sale, Sale Into, Distribution and/or Use in Nassau and Suffolk Counties of New York State.

- **NEW YORK: DO NOT** apply more than 16 fl oz product/acre per year. **DO NOT** exceed 0.0528 lb ai/acre per year.
- **[[CALIFORNIA:] DO NOT** exceed maximum application rate of 8 fl oz product/acre (0.0264 lb ai/acre) per application.]
- **[[CALIFORNIA:] DO NOT** apply more than 1 application per year.]
- **[[CALIFORNIA:] DO NOT** apply in California Ground Water Protection Areas (https://www.cdpr.ca.gov/docs/emon/grndwtr/gwpa_locations.htm)]
- **[[CALIFORNIA:] DO NOT** apply directly to soil.]
- **DO NOT** apply via chemigation.
- **[DO NOT** apply via aerial application.]
- **NOT** for use in greenhouses.

[[CALIFORNIA:] Prevent movement into soil: Minimize pesticide contact with the soil surface by directing spray application at foliage only. **DO NOT** apply product unless plant surfaces have time to fully dry prior to expected rainfall or scheduled overhead irrigation.]

[[CALIFORNIA:] Irrigation requirements: Apply on fields irrigated with drip or sprinkler irrigation only and ensure that irrigation does not exceed 125% of evapotranspiration. **DO NOT** apply to fields irrigated by flood, furrow or other surface flow delivery methods]

Mixing directions

DO NOT combine GATTEN® in the spray tank with other pesticides, surfactants, or fertilizers unless prior use has shown the combination physically compatible, effective, and non-injurious under conditions of use.

GATTEN® is physically and biologically compatible with many registered pesticides, fertilizers or micronutrients. Contact your supplier or a local Extension for advice when considering mixing GATTEN® with other products. If you have no experience with the combination you are considering, you must conduct a test to determine physical compatibility. To do so, add the specified proportions of both chemicals with the same proportion of water, as will be present in the chemical supply tank, into a suitable container; mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be readily remixed, the mixture must be physically compatible.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank

mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Rotational Crops Directions

Any crop may be planted immediately after the last application.

Resistance-Management

Fields must be scouted prior to application to identify the disease that is present and to determine if the intended application may be effective. Fields must be scouted at the appropriate period after application to verify that the treatment was effective.

For further information or to report suspected resistance contact OAT Agrio Co., Ltd. at https://www.oat-agrio.co.jp/en/contact_en/. You can also contact your pesticide distributor or university extension specialist to report resistance. If resistance is suspected, treat with an effective pesticide, if available, having a different mode of action and/or use non-chemical means to best manage the pest.

For resistance management, GATTEN®, contains a Group U13 fungicide. Any fungal population may contain individuals naturally resistant to GATTEN® and other Group U13 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay fungicide resistance, take one or more of the following steps:

- **DO NOT** apply more than maximum number of applications for each labeled crop in the directions for application table below and avoid consecutive sprays of GATTEN® or other fungicides in the same group in a season.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.

Directions for Application

Crop	PHI	Target Diseases	Application Rate fl oz product /A (lb a.i./A)	Directions	Restrictions
Apples	14	<i>P. leucotricha</i> (Powdery Mildew)	5.5 (0.017) – 12.8 (0.04)	<p>Apply by air using a minimum of 10 gallons of water per acre.</p> <p>Apply by ground using water volumes of 20 to 100 gallons per acre for concentrate sprays and 101 to 200 gallons per acre for dilute sprays.</p> <p>Repeat applications at 7 to 14-day intervals, depending on disease pressure.</p> <p>Apply to foliage during the fruiting stage [following rapid canopy development] using airblast ground equipment or by air.</p>	<p>Maximum Application rate: 12.8 fl oz product /A (0.04 lb a.i./A)</p> <p>DO NOT apply more than 4 applications per year.</p> <p>The minimum re-treatment interval (RTI) is 7 days.</p> <p>DO NOT exceed a maximum of 0.16 lb ai per acre per year.</p> <p>DO NOT apply within 14 days of harvest.</p> <p>NEW YORK: DO NOT apply more than 16 fl oz product/acre per year. DO NOT exceed 0.0528 lb ai/acre per year.</p> <p>CALIFORNIA: DO NOT exceed maximum application rate of 8 fl oz product/acre (0.0264 lb ai/acre). DO NOT apply more than 1 application per year.</p>

Crop	PHI	Target Diseases	Application Rate fl oz product /A (lb a.i./A)	Directions	Restrictions
<p>Cherry subgroup 12-12A including:</p> <p>Capulin Cherry, black Cherry, Nanking Cherry, sweet Cherry, tart Cultivars, varieties, and/or hybrids of these</p>	3	<i>P. clandestina</i> (Powdery Mildew)	5.5 (0.017) – 12.8 (0.04)	<p>Apply by air using a minimum of 10 gallons of water per acre.</p> <p>Apply by ground using water volumes of 30 to 100 gallons per acre for concentrate sprays and 101 to 300 gallons per acre for dilute sprays.</p> <p>Repeat applications at 7 to 14-day intervals, depending on disease pressure.</p> <p>Apply to foliage during the fruiting stage [following rapid canopy development] using airblast ground equipment or by air.</p>	<p>Maximum Application rate: 12.8 fl oz product /A (0.04 lb a.i./A)</p> <p>DO NOT apply more than 4 applications per year.</p> <p>The minimum re-treatment interval (RTI) is 7 days.</p> <p>DO NOT exceed a maximum of 0.16 lb ai per acre per year.</p> <p>DO NOT apply within 3 days of harvest.</p> <p>NEW YORK: DO NOT apply more than 16 fl oz product/acre per year. DO NOT exceed 0.0528 lb ai/acre per year.</p> <p>CALIFORNIA: DO NOT exceed maximum application rate of 8 fl oz product/acre (0.0264 lb ai/acre). DO NOT apply more than 1 application per year.</p>

Crop	PHI	Target Diseases	Application Rate fl oz product /A (lb a.i./A)	Directions	Restrictions
<p>Melon subgroup 9A and Squash/cucumber subgroup 9B including:</p> <p>All types and hybrids of:</p> <p>Chayote Chinese waxgourd Citron melon Cucumber[*] Gherkin Gourd, edible; Chinese okra Cucuzza Hyotan Hechima <i>Momordica</i> spp. Balsam apple Balsam pear Bitter melon Chinese cucumber Muskmelon Cantaloupe Casaba Crenshaw melon Golden pershaw melon Honeydew melon Honey balls Mango melon Persian melon Pineapple melon Santa Claus melon Snake melon Pumpkin Squash, summer Crookneck squash Scallop squash Straightneck squash Vegetable marrow Zucchini Squash, winter Acorn squash Butternut squash Calabaza Hubbard squash Spaghetti squash Watermelon</p>	0	<i>G. cichoracearum</i> (Powdery Mildew)	5.5 (0.017) – 12.8 (0.04)	<p>Apply by air using a minimum of 5 gallons of water per acre.</p> <p>Apply by ground using water volumes of 30 to 100 gallons per acre for sprays.</p> <p>Repeat applications at 7 to 14-day intervals, depending on disease pressure.</p> <p>Apply to foliage during seedling to the fruiting stage using broadcast ground equipment or by air.</p>	<p>Maximum Application rate: 12.8 fl oz product /A (0.04 lb a.i./A)</p> <p>DO NOT apply more than 5 applications during each growing season. Maximum of two growing seasons per year.</p> <p>The minimum re-treatment interval (RTI) is 7 days.</p> <p>DO NOT exceed a maximum of 0.20 lb ai per acre per year. Crop can be harvested after the product has dried.</p> <p>NEW YORK: DO NOT apply more than 16 fl oz product/acre per year. DO NOT exceed 0.0528 lb ai/acre per year.</p> <p>CALIFORNIA: DO NOT exceed maximum application rate of 8 fl oz product/acre (0.0264 lb ai/acre). DO NOT apply more than 1 application per year.</p>

Crop	PHI	Target Diseases	Application Rate fl oz product /A (lb a.i./A)	Directions	Restrictions
<p>Small fruit, vine climbing, except fuzzy kiwifruit, subgroup 13-07F including:</p> <p>Amur river grape Gooseberry Grape Kiwifruit, hardy Maypop Schisandra berry Cultivars, varieties, and/or hybrids of these</p>	14	<i>E. necator</i> (Powdery Mildew)	5.5 (0.017) – 12.8 (0.04)	<p>Apply by air using a minimum of 10 gallons of water per acre</p> <p>Apply by ground using water volumes of 20 to 100 gallons per acre for concentrate sprays and 101 to 400 gallons per acre for dilute sprays.</p> <p>Repeat applications at 7 to 14-day intervals, depending on disease pressure.</p> <p>Apply to foliage during the fruiting stage [from bloom when canopy has completed rapid shoot growth period] using airblast ground equipment or by air.</p>	<p>Maximum Application rate: 12.8 fl oz product /A (0.04 lb a.i./A)</p> <p>DO NOT apply more than 4 times applications per year.</p> <p>The minimum re-treatment interval (RTI) is 7 days.</p> <p>DO NOT exceed a maximum of 0.16 lb ai per acre per year.</p> <p>DO NOT apply within 14 days of harvest.</p> <p>NEW YORK: DO NOT apply more than 16 fl oz product/acre per year. DO NOT exceed 0.0528 lb ai/acre per year.</p> <p>CALIFORNIA: DO NOT exceed maximum application rate of 8 fl oz product/acre (0.0264 lb ai/acre). DO NOT apply more than 1 application per year.</p>

Crop	PHI	Target Diseases	Application Rate fl oz product /A (lb a.i./A)	Directions	Restrictions
<p>Berry, low growing, subgroup 13-07G including:</p> <p>Bearberry Bilberry Blueberry, lowbush Cloudberry Cranberry Lingonberry Muntries Partridgeberry Strawberry Cultivars, varieties, and/or hybrids of these</p>	0	<i>P. aphanis</i> (Powdery Mildew)	5.5 (0.017) – 12.8 (0.04)	<p>Apply by air using a minimum of 10 gallons of water per acre.</p> <p>Apply by ground using water volumes of 50 to 200 gallons per acre for sprays.</p> <p>Repeat applications at 7 to 14-day intervals, depending on disease pressure.</p> <p>Apply to foliage during the fruiting stage using broadcast ground equipment or by air.</p>	<p>Maximum Application rate: 12.8 fl oz product /A (0.04 lb a.i./A)</p> <p>DO NOT apply more than 5 applications per year.</p> <p>The minimum re-treatment interval (RTI) is 7 days.</p> <p>DO NOT exceed a maximum of 0.20 lb ai per acre per year. Crop can be harvested after the product has dried.</p> <p>NEW YORK: DO NOT apply more than 16 fl oz product/acre per year. DO NOT exceed 0.0528 lb ai/acre per year.</p> <p>CALIFORNIA: DO NOT exceed maximum application rate of 8 fl oz product/acre (0.0264 lb ai/acre). DO NOT apply more than 1 application per year.</p>

Crop	PHI	Target Diseases	Application Rate fl oz product /A (lb a.i./A)	Directions	Restrictions
Hops[*]	7	<i>P. macularis</i> (Powdery mildew)	5.5 (0.017) – 12.8 (0.04)	<p>Apply by air using a minimum of 10 gallons of water per acre.</p> <p>Apply by ground using water volumes of 50 to 200 gallons per acre for sprays.</p> <p>Repeat applications at 7 to 14-day intervals, depending on disease pressure.</p> <p>Apply to foliage during the fruiting stage using broadcast ground equipment or by air.</p>	<p>Maximum Application rate: 12.8 fl oz product /A (0.04 lb a.i./A)</p> <p>DO NOT apply more than 4 applications per year.</p> <p>The minimum re-treatment interval (RTI) is 7 days.</p> <p>DO NOT exceed a maximum of 0.16 lb ai per acre per year.</p> <p>NEW YORK: DO NOT apply more than 16 fl oz product/acre per year. DO NOT exceed 0.0528 lb ai/acre per year.</p> <p>CALIFORNIA: DO NOT exceed maximum application rate of 8 fl oz product/acre (0.0264 lb ai/acre). DO NOT apply more than 1 application per year.</p>

[* indicates Not for Use in California]

Storage and Disposal

DO NOT contaminate water, food, or feed by storage or disposal.

Storage:

Store product in a secure dry, cool and well ventilated place. Store product in original container.

DO NOT store for prolonged periods in direct sunlight.

Pesticide Disposal:

Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling:

Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

(5 gallons or less) Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(Larger than 5 gallons) Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Then offer for [recycling if available or] reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

WARRANTY AND CONDITIONS OF SALE

SELLERS OFFER AND THE BUYER ACCEPTS THIS PRODUCT UNDER THE FOLLOWING AGREED CONDITIONS OF SALE AND WARRANTY

The directions for the use of this product are believed to be adequate and must be followed carefully. However, it is impossible to take into account all the variables and to eliminate risks associated with its use. Injury or damage may result because of conditions that are beyond the control of the SELLER.

To the extent consistent with applicable law, SELLER warrants that this product conforms to the chemical description on this label and is believed to be reasonably fit for the purposes stated on this label when used in accordance with the directions on this label under conditions of normal use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SELLER MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

To the extent consistent with applicable law, the SELLER shall not be liable for consequential, special or indirect damages resulting from the use or handling of the product. Any variation or exemption from this warranty must be in writing and signed by an authorized representative of the SELLER.

20220204LAS