



WEED-FREE® S

(DSMA & SURFACTANT)

Patent Pending

FOR SELECTIVE POST-EMERGENCE WEED CONTROL

READ THIS LABEL BEFORE USING PRODUCT

INGREDIENT STATEMENT

ACTIVE INGREDIENT:
Disodium Methanearsonate*..... 29.11%
INERT INGREDIENTS..... 70.89%
TOTAL..... 100.00%

Total Arsenic, all in water soluble form, expressed as elemental..... 12.28%

*Contains 3.2 lbs. DSMA per gallon (equivalent to 5 lbs. disodium methanearsonate hexahydrate per gallon.)

WARNING

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contact with skin. Avoid breathing dust or spray mist. Wash hands after using. Avoid spray near feed and food products. Keep away from children and domestic animals. Keep children and animals off treated area until washed into soil.

ANTIDOTE: If taken internally, induce vomiting and CALL PHYSICIAN AT ONCE.

Do not contaminate water used for domestic and irrigation purposes.


DO NOT FEED ANY PART OF TREATED COTTON PLANTS FORAGE OR TREATED VEGETATION TO LIVESTOCK. DO NOT GRAZE TREATED AREAS.

Clean application equipment thoroughly after every use. Do not use this product for any purpose or in any manner not shown on this label. To do so may result in unlawful residues on crop plants, injury to nearby susceptible desirable vegetation or may not give desired results.

Applications which result in some of the spray solution being deposited onto the cotton may result in injury to or kill of the cotton and a subsequent loss of yield.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

20



DIRECTIONS FOR USE

COTTON

Use WEED-FREE SUPER for post emergence control of certain annual and perennial grasses, weeds and vines such as watergrass, goosegrass, nutgrass, cocklebur, annual morning glory and Johnsongrass. WEED-FREE SUPER should be used from the time the cotton plants are 3 inches high until first bloom, and not before this period. Apply WEED-FREE SUPER when grasses and weeds are three inches high. Best results are obtained when the air temperature is above 70° F.

Add 2 quarts of WEED-FREE SUPER to approximately forty gallons of water and mix thoroughly. This solution should be applied at the rate of 40 gallons per acre of soil surface actually sprayed. When making directed band applications on the drill, use an amount of spray solution in proportion to the amount of land surface actually being sprayed. Thus, if a total band width of 18 inches with rows spaced 40 inches apart, use approximately 18 gallons of spray solution per acre of crop treated. When weed and grass control is desired on the middles, an extra nozzle may be mounted to the side to apply a separate directed spray on these areas as well. Spray nozzles should be positioned to give complete coverage of the weeds and grasses to be controlled without contact with the cotton foliage. The rate of application for cotton must not exceed three pounds DSMA (anhydrous) per acre.

Do not make more than 2 applications of any combination of products containing methanearsonates per crop season nor apply them after the first bloom. To do so may result in unlawful residues of arsenic in cottonseed.

EPA Reg. No. 1022403

MANUFACTURED BY **CHAPMAN CHEMICAL COMPANY**

WEED-FREE[®] SUPER

(DSMA & SURFACTANT)

Patent Pending

FOR SELECTIVE POST-EMERGENCE WEED CONTROL

NET CONTENTS 5 GALLONS

READ THIS LABEL BEFORE USING PRODUCT

INGREDIENT STATEMENT

ACTIVE INGREDIENT:

sodium Methanearsonate*	29.11%
INERT INGREDIENTS	70.89%
TOTAL	100.00%

arsenic, all in water soluble form, expressed as elemental..... 12.28%

contains 3.2 lbs. DSMA per gallon (equivalent to 3.2 lbs. disodium methanearsonate hexahydrate per gallon.)

WARNING

KEEP OUT OF REACH OF CHILDREN

Do not swallow. Avoid contact with skin. Avoid breathing or spray mist. Wash hands after using. Avoid storage and food products. Keep away from children and domestic animals. Keep children and animals off treated areas until into soil.

If taken internally, induce vomiting and CALL A DOCTOR AT ONCE.

Contaminate water used for domestic and irrigation purposes.

DO NOT GRAZE ANY PART OF TREATED COTTON PLANTS FORAGE OR OTHER VEGETATION TO LIVESTOCK. DO NOT GRAZE TREATED AREAS.

Wash application equipment thoroughly after every use. Do not use this product for any purpose or in any manner not given on label. To do so may result in unlawful residues in soil, injury to nearby susceptible desirable vegetation, and will not give desired results.

Do not use on crops which result in some of the spray solution getting on cotton. Cotton may result in injury to or kill of the cotton and consequent loss of yield.

Contaminate water by cleaning of equipment, or disposal

20

GENERAL WEED AND GRASS CONTROL

WEED-FREE SUPER is useful in controlling many noxious weeds and grasses including cocklebur, goosegrass, watergrass, nutgrass, pigweed, ragweed and Johnsongrass. Add two quarts of WEED-FREE SUPER to 30 gallons of water. The unwanted vegetation should be thoroughly covered with spray to the point of run-off. Spot retreatments may be necessary and should be made where regrowth occurs. Any spray equipment which will give good spray coverage may be used, with power sprayers being particularly effective for such areas as rights-of-way, drainage ditch banks, fence rows, implement storage yards and similar non-crop areas where control of the grasses and weeds mentioned above is desired.

Do not reuse empty container. Destroy by burying in a safe place.

NON-WARRANTY

Neither Chapman Chemical Company nor the seller makes any representation or warranty, expressed or implied, concerning this product, except that it is of the Chapman Chemical Company standard quality. Neither shall be held responsible in any manner for any personal injury, property damage or other type of loss resulting from the improper handling storage and use of this product. The buyer accepts and uses this product on these conditions.

21

CHAPMAN CHEMICAL COMPANY MEMPHIS, TENNESSEE