

11600-4

2-3-2003

1/10

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

FEB 3 2003

Ms. Christina M. Swick
Agent for Conklin Company, Inc.
Lewis & Harrison
122 C Street, NW
Suite 740
Washington, DC. 20001

Subject: Sanox II
EPA Registration No. 11600-4
Application Date: 11/05/02
Receipt Date: 11/06/02

Dear Ms. Swick:

The following amendment, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable with the conditions listed below:

- Update First Aid Statements
Add pest claims against a number of organisms

CONDITIONS

Revise the label as follows:

- 1. On page 2 of the label, delete the surface claims "cutting boards, chopping blocks, coolers, ice chests, refrigerator meat, fruit, vegetables and egg bins. These claims were not cited on the last accepted stamp label nor the cited product 6836-75.
2. Page 3, add the word "glazed" before "enameled surfaces" and before "porcelain".
3. Page 5, revise the Veterinary Practice/Animal Care/Animal Laboratory/Zoos/Pet Shop/Kennels' Disinfection Directions by inserting the following statement "Wipe or allow to air dry" between "minutes" and "ventilate" followed by "Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and manure". Also, you will need to decide whether your product will be used to disinfect feeding and watering equipment because the statement to rinse the equipment with potable water is contradictory to the phrase that this product should not be used on such

CONCURRENCES

Table with columns for SYMBOL, SURNAME, and DATE, containing handwritten entries like 75100, Jones, and 11-06-03.

2/10

equipment. If you decide to disinfect feeding and watering equipment, you must add the statement, "Empty all feeding and watering appliances" to the directions before the clean with detergent statement.


4. Page 6, under Heavy Duty (1:16 dilution), revise the sentence to read, "to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer. Brush or swab thoroughly, allow to stand for 10 minutes and flush.

GENERAL COMMENTS

A stamped copy of the accepted labeling is enclosed. Submit three (3) copies of your final printed labeling before distributing or selling the product bearing the revised labeling.

Should you have any questions or comments concerning this letter, please contact Zenobia Jones at (703) 308-6198.

Sincerely,



Velma Noble
Product Manager 31
Regulatory Management Branch 1
Antimicrobial Division (7510C)

SANOX[®] II

Highlighted Label 3/10

Disinfectant - Cleaner - Sanitizer - Fungicide
Mildewstat - Virucide* - Deodorizer

For Hospital, Institutional, Industrial and Home Use

Virucide*, kills HIV-1 on pre-cleaned, hard, nonporous surfaces
Effective in hard water

Formulated for Effective Poultry Premises Sanitation ✓
Formulated for Effective Swine Premises Sanitation ✓
Formulated for Effective Farm Premises Sanitation ✓
Formulated for Veterinary Practice / Animal Care / Animal Laboratory Disinfection ✓

Active Ingredients:

Octyl decyl dimethyl ammonium chloride.....	1.650%
Dioctyl dimethyl ammonium chloride.....	0.660%
Didecyl dimethyl ammonium chloride.....	0.990%
Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%).....	2.200%
Dimethyl benzyl ammonium chloride	
Inert Ingredients	94.500% ✓
Total	100.000%

KEEP OUT OF REACH OF CHILDREN

DANGER

SEE LEFT / SIDE / RIGHT / BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No.: 11600-4
EPA Establishment No.: 11600-MN-1

Net Contents:
(insert container size here)

Manufacturer by:
Conklin Company, Inc.
Consumer Products Division
551 Valley Park Drive
P. O. Box 155
Shakopee, MN 55379-0155
(952) 445-6010
www.conklin.com

ACCEPTED
with COMMENTS
in EPA Letter Dated:

FEB 3 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.

11600-4

4/10

✓ Sanox® II is a proven "one-step" disinfectant-cleaner-sanitizer-fungicide-mildewstat-virucide* which is effective in water up to 400 ppm hardness in the presence of 5% serum contamination.

✓ Cross-contamination is a major housekeeping concern not only in hospitals, but in schools, homes, institutions, and industry. Sanox II has been formulated to aid in the reduction of cross-contamination in these areas.

Sanox II is a one-step disinfectant-cleaner that is effective against a broad spectrum of bacteria, is virucidal*, fungicidal and will prevent the growth of mold and mildew and their odors when used as directed. Sanox II is an effective one-step sanitizer-cleaner for use on non-food contact surfaces.

✓ Sanox II is an effective, yet economical heavy-duty disinfectant-cleaner that disinfects, cleans and deodorizes in one labor-saving step. It can be diluted for use with a mop and bucket, trigger sprayer, sponge or by soaking.

✓ Sanox II is formulated for use in daily maintenance programs with a balance of detergents, biocides and malodor counteractants that deliver effective cleaning, disinfecting and malodor control.

✓ Sanox II is a complete, chemically balanced disinfectant and sanitizer that provides clear use solutions even in the presence of hard water.

✓ Sanox II deodorizes by killing microorganisms that cause offensive odors.

✓ Sanox II is a concentrated one-step (Hospital Use) disinfect-cleaner that is effective against a broad spectrum of bacteria, is virucide* (including HIV-1, the AIDS virus), fungicidal, and inhibits the growth of mold and mildew and their odors when used as directed.

✓ Sanox II meets AOAC Use-Dilution Test Standards for hospital disinfectants.

RECOMMENDED USE AREAS

Sanox II will clean and deodorize surfaces in toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas, and other places where bacterial growth can cause malodors.

✓ Use Sanox II for non-scratch cleaning of showers and tubs, shower doors and curtains, fixtures and toilet bowls.

✓ Use Sanox II to clean and disinfect finished floors, sinks and tubs.

✓ Use Sanox II in kitchens, bathrooms and other household areas.

Sanox II will not leave grit or soap scum.

✓ Use Sanox II for sanitizing and disinfecting of ultrasound transducers, probes, mammography compressor plates and other hard nonporous surfaces.

✓ Use Sanox II on coils and drain pans of air conditioning and refrigeration equipment and heat pumps. Follow the directions for sanitation of non-food contact surfaces.

✓ Sanox II is a versatile cleaner and broad-spectrum disinfectant formulated for use only on non-critical instruments in Ultrasonic Baths (Ultrasonic cleaning units).

✓ Use Sanox II to clean and disinfect nonporous salon / barber tools and instruments, combs, brushes, scissors, blades and manicure instruments.

Use Sanox II to clean and disinfect finished floors. Can be applied with damp mop or autoscrubber. Cleans and disinfects without dulling gloss. To limit gloss reduction, use at 2 oz. per gallon of water (1:64 dilution).

Apply Sanox II to walls, floors, and other hard (inanimate) nonporous surfaces such as tables, chairs, countertops, stovetops, appliances, refrigerators, microwave ovens, cutting boards and chopping blocks, sinks, tile, coolers and ice chests, outdoor furniture, shower stalls and tubs, kennel runs, cages, garbage cans.

ACCEPTED
with COMMENTS
in EPA Letter Dated:
FEB 3 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
registered under F.F.A. Reg. No
11600-4

5/10

refrigerator, meat, fruit, vegetable and egg bins, cabinets, desks, telephones, doorknobs, whirlpool bathtubs, coil and drain pans of air conditioning and refrigeration equipment, heat pumps, conductive flooring, fiberglass, enameled surfaces, finished woodwork, vinyl, plastic upholstery, porcelain, toilet bowls and urinals surfaces with a cloth, mop or mechanical spray so as to thoroughly wet surfaces. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution daily or when use-solution becomes visibly dirty.

Use Sanox II in homes, hospitals, medical and dental offices, nursing homes, day care centers, nurseries, restaurants, bars, cafeterias, institutional kitchens, food preparation and storage areas, convenience stores, public rest rooms, institutions, schools, colleges, athletic facilities, locker rooms, dressing rooms, shower and bath areas, exercise facilities, camp grounds, play grounds, public facilities, food processing plants, transportation terminals, hotels and motels, factories, business and office buildings, barber shops, salons, boats, ships, campers, trailers, mobile homes, cars, buses, trains, airplanes, emergency vehicles, prisons, correctional facilities and ultrasonic baths.

Use Sanox II in USDA inspected food processing facilities, dairy farms, hog farms, poultry and turkey farms, kennels, pet animal quarters, zoos, pet shops and animal laboratories.

Use Sanox II for Poultry Premise Sanitation (Hatcheries) on hard nonporous surfaces in: Egg Receiving Area, Egg Holding Area, Tray Dumping Area, Trays, Racks, Egg Flats, Chick Holding Area, Hatchery Room, Chick Processing Area, Chick Loading Area, Poultry Buildings, Ceilings, Sidewalls and Floors, Drinkers and Other Poultry House Related Equipment.

Use Sanox II for Swine Premises Sanitation of hard, nonporous surfaces: Waterers and Feeders, Hauling Equipment, Dressing Plants, Loading Equipment, Farrowing Barns and Areas, Nursery Blocks, Creep Area, Chutes.

Use Sanox II for Farm Premise Sanitation of hard, nonporous surfaces: Floors, Walls, Feed Racks, Managers, Troughs, Automatic Feeders, Fountains and Waterers, Forks, Shovels, Scrapers and other nonporous surfaces in Barns, Pens, Stalls, Chutes and other facilities and fixtures occupied or crossed by animals.

Sanox II is a versatile disinfectant for Veterinarian, Animal Care and Farm Premise applications.

ACCEPTED with COMMENTS in EPA Letter Dated:

FEB 9 2003

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 11600-4

RECOMMENDED DISINFECTION DIRECTIONS

DISINFECTION – To disinfect hard, nonporous surfaces in hospitals, add 2 oz. of Sanox II per gallon (1:64 dilution) of water. Treated surfaces must remain wet for 10 minutes. At 1 1/4 oz. of Sanox II per gallon (1:16 dilution) of water, Sanox II will disinfect hard, nonporous surfaces in schools, industry and non-medical institutions.

Preparation of Disinfectant Use-Solution:

1. Add (mix) 2 oz. of Sanox II per gallon of water (1:64 dilution)
2. For heavy duty use, add (mix) 8 oz. of Sanox II per gallon of water (1:16 dilution)

Other dilution ratios -

1. Add (mix) 2 oz. of Sanox II per 4 gallons of water (1:256 dilution) to achieve 200 ppm active
2. Add (mix) 2 oz. of Sanox II per 5 1/2 gallons of water (1:352 dilution) to achieve 150 ppm active

Two (2) Ounces Per Gallon of Water Use Level (1:64 dilution):

The activity of Sanox II has been evaluated in the presence of 5% serum and 400 ppm hard water by the AOAC Use-Dilution Test Standards and found to be effective against a broad spectrum of gram negative and gram positive organisms. Sanox II is also Bactericidal according to the current AOAC Use-Dilution Test Method, Fungicidal according to the AOAC Fungicidal Test and Virucidal according to the virucidal qualification modified in the presence of 400 ppm hard water plus 5% serum. Below is a list of organisms that Sanox II is effective against.

6/10

Organisms	Organisms
Bacteria:	Streptococcus pyogenes (Strep) ✓
Pseudomonas aeruginosa (Pseudomonas) ✓	Vibrio cholerae ✓
Staphylococcus aureus (Staph) ✓	Viruses:
Salmonella choleraesuis (Salmonella) ✓	*Herpes Simplex Virus Types 1 & 2 (Herpes) ✓
Acinetobacter baumannii (Acinetobacter) ✓	*HIV-1 (the AIDS Virus) ✓
Brevibacterium ammoniagenes (Brevibacterium) ✓	*Influenza Type A / Brazil (Influenza) ✓
Campylobacter jejuni (Campylobacter) ✓	*Respiratory Syncytial Virus (RSV) ✓
Enterobacter aerogenes ✓	*Vaccinia (Pox Virus) ✓
Enterococcus faecalis (Enterococcus) ✓	*Adenovirus Type 5 (Adenovirus) at 1:16 dilution - 8 oz./gallon ✓
Enterococcus faecalis - Vancomycin resistant (VRE) ✓	Animal Viruses:
Escherichia coli (E. coli) ✓	Avian Influenza Virus ✓
Klebsiella pneumoniae (Klebsiella) ✓	Avian Infectious Bronchitis Virus ✓
Pseudomonas cepacia (Pseudomonas) ✓	Newcastle's Disease Virus ✓
Salmonella schottmueleri (Salmonella) ✓	Canine Distemper Virus ✓
Salmonella typhi (Salmonella) ✓	Pseudorabies Virus ✓
Serratia marcescens ✓	Fungi:
Shigella dysenteriae (Shigella) ✓	Trichophyton mentagrophytes (the Athlete's Foot Fungus) ✓
Staphylococcus aureus Methicillin-Resistant (MRSA) ✓	Aspergillus niger ✓
Staphylococcus aureus Multi-Drug Resistant ✓	
Staphylococcus aureus - Vancomycin Intermediate Resistant (VISA) ✓	

Disinfection / Virucidal Directions:

1. Apply use-solution to hard, nonporous surfaces with a cloth, mop, sponge, coarse sprayer, or, by immersing, thoroughly wetting surfaces.
2. For spray applications, use a coarse spray device. Spray 6 to 8 inches from surface and, rub with brush, sponge or cloth. Avoid breathing spray.
3. Treated surfaces must remain wet for 10 minutes. Wipe dry with a cloth, sponge or mop, or, allow to air dry.
4. Prepare a fresh solution daily, or more often if the use-solution becomes visibly diluted or soiled.
5. After disinfecting food contact surfaces, rinse with potable water. DO NOT use on utensils, glasses or dishes.
6. If heavily soiled, a preliminary cleaning is recommended.

Virucidal* Performance (1:64 dilution):

At 2 oz. of Sanox II per gallon use level, Sanox II was evaluated in the presence of 5% serum and 400 ppm hard water for a contact time of 10 minutes at room temperature (20° to 25° C) against the following viruses found on inanimate, hard, nonporous environmental surfaces.

Organisms	Organisms
Viruses:	Animal Viruses:
*Herpes Simplex Virus Types 1 & 2 (Herpes) ✓	Avian Influenza Virus ✓
*HIV-1 (the AIDS Virus) ✓	Avian Infectious Bronchitis Virus ✓
*Influenza Type A / Brazil (Influenza) ✓	Newcastle's Disease Virus ✓
*Respiratory Syncytial Virus (RSV) ✓	Canine Distemper Virus ✓
*Vaccinia (Pox Virus) ✓	Pseudorabies Virus ✓
*Adenovirus Type 5 (Adenovirus) at 1:16 dilution - 8 oz./gallon ✓	

ACCEPTED with COMMENTS in EPA Letter Dated: FEB 9 2003

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 11600-4

Fungicidal Control (1:64 dilution):

Sanox II is an effective fungicide against Trichophyton mentagrophytes (the Athlete's Foot Fungus) when used on surfaces such as locker rooms, dressing rooms, shower and bath areas, exercise facilities, etc., at 2 oz of Sanox II per gallon of water in 250 ppm hard water and 5% serum.

Mold and Mildew Control (1:64 dilution):

At 2 oz. of Sanox II per gallon of water, Sanox II will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces (as indicated in general instructions above). Allow to air dry on surface and repeat when mildew returns.

Salon / Barber Instruments and Tools Disinfection (Bactericide / Virucide* / Fungicide Directions (1:32 dilution):

Mix 4 oz. of Sanox II per gallon of water (1 oz. of Sanox II per quart) to disinfect hard, nonporous instruments and tools. Completely immerse combs, brushes, rollers, manicure and other salon instruments and tools for 10

7/10

minutes. For heavily soiled instruments and tools, a preliminary cleaning is required. Rinse thoroughly and dry before use. Prepare a fresh use-solution daily, or more often if solution becomes diluted or soiled.

Note: Plastics may remain immersed until ready to use. Stainless steel instruments must be removed after 10 minutes, rinsed, dried and kept in a clean non-contaminated receptacle.

Ultrasonic Bath Disinfectant Directions (1:64 dilution):

Use Sanox II to disinfect hard, nonporous, non-critical objects compatible with Ultrasonic cleaning units. Pour a fresh use-solution of 2 oz. of Sanox II per gallon of water directly into bath chamber. Pre-clean heavily soiled objects. Place objects into bath chamber and operate for a minimum of 10 minutes, according to the manufacturer's directions. Remove objects and wash with sterile water, or allow to air dry. Prepare a fresh use-solution for each disinfection use.

Disinfection of Whirlpool Units (1:64 dilution):

After using the unit, drain and refill with fresh water to just cover the intake valve. Add 2 oz of Sanox II for each gallon of fresh water added. Start the pump to circulate the solution. Wash down the sides, seat and any related equipment with a clean swab, brush or sponge. Treated surfaces must remain wet for 10 minutes. After the unit has been thoroughly cleaned, drain the solution from the unit and rinse cleaned surfaces with fresh water.

Poultry and Swine Premise Disinfection Directions (1:64 dilution):

Remove all animals and feeds from premises. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 2 oz of Sanox II per gallon of water use-solution, and saturate surfaces with the recommended disinfecting solution for 10 minutes. Immerse all types of equipment used in handling and restraining animals, as well as forks, shovels and scrapers. DO NOT house animals or use equipment until treatment has been absorbed, set or dried. Thoroughly scrub feeding and watering equipment with soap or detergent, and rinse with potable water before reuse.

Hatcheries (1:64 dilution):

Use 2 oz of Sanox II per gallon of water to treat the following hard, nonporous surfaces: hatchers, setters, trays, racks, carts, sexing tables, delivery vehicles and other hard surfaces. Leave all treated surfaces wet for 10 minutes or more. Then allow to air dry.

Vehicles (1:64 dilution):

Clean all vehicles including mats, crates, cabs and wheels with high pressure water and Sanox II. Use a 2 oz. of Sanox II per gallon of water mix. Leave all treated surfaces wet for 10 minutes or more.

Dressing Plant Disinfection Directions (1:64 dilution):

Cover or remove all food and packaging materials, and remove all heavy soils prior to application. Saturate all surfaces (equipment, utensils, walls, floors and loading platforms) with a 2 oz of Sanox II per gallon of water use-solution. Scrub all surfaces with disinfectant, leave wet for 10 minutes, then thoroughly rinse with potable water before operations are resumed.

Veterinary Practice / Animal Care / Animal Laboratory / Zoos / Pet Shop Kennels Disinfection Directions (1:64 dilution):

Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls, equipment and utensils used or occupied by animals. Clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 2 oz of Sanox II per gallon of water (or equivalent dilution) for a period of 10 minutes. Ventilate buildings and other closed spaces. DO NOT house animals or use equipment until treatment has been absorbed, set or dried.

Farm Premises Disinfection Directions (1:64 dilution):

Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls, facilities and fixture surfaces utilized by animals. Thoroughly clean all surfaces with soap or detergent, and rinse with water. Saturate all surfaces with a use-solution of 2 oz of Sanox II per gallon of water (or, equivalent dilution) for a period of 10 minutes. Ventilate buildings, cars, trucks, boats and other closed spaces. DO NOT house livestock or use equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feeding equipment with soap or detergent, and rinse with potable water before reuse.

ACCEPTED with COMMENTS in EPA Letter dated 11/6/00-4 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended for the purpose of...

8/10

*KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1, associated with AIDS).

"SPECIAL INSTRUCTION FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES / OBJECTS SOILED WITH BLOOD / BODY FLUIDS" (1:64 dilution)

Personal Protection: Clean up should always be done wearing protective latex gloves, gowns, masks and eye protection.

Cleaning Procedure: Blood and other body fluids containing HIV-1 must be thoroughly cleaned from surfaces and objects before the application of Sanox II.

Contact Time: Leave surface wet for 1 minute (60 seconds) for HIV-1 with a 2 oz. per gallon use-solution. Use a 10 minute contact time for disinfection against all other viruses, bacteria and fungi claimed.

Disposal of Infectious Material: Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to local regulations for infectious waste disposal.

For Schools, Homes, Industry and Non-Medical Institutional Use (≅ 1:100 dilution):

At 1 ¼ oz. Sanox II per gallon of water, Sanox II delivers excellent cleaning and germicidal effectiveness. It is effective against Staphylococcus aureus (Staph), Salmonella choleraesuis (Salmonella), Escherichia coli (E. coli) and Serratia marcescens. The same AOAC tests used to confirm performance for hospitals were used.

Non-Acid Toilet Bowl Disinfectant / Cleaner Application -

Concentrate:

- 1. Remove gross filth prior to disinfection.
- 2. Add 2 oz of Sanox II to the toilet bowl (0.75 gallon) and mix. Brush over exposed surfaces, and allow to stand for 10 minutes and flush.

Use-solution (1:64 dilution):

- 1. Remove gross filth prior to disinfection.
- 2. Empty toilet or urinal and apply 2 oz. of Sanox II per gallon of water use-solution to exposed surfaces with a cloth, mop, sponge or mechanical spray, and allow to stand for 10 minutes and flush.

Heavy-Duty (1:16 dilution):

- 1. Remove gross filth prior to disinfection.
- 2. Empty toilet or urinal and apply 8 oz. of Sanox II per gallon of water use-solution to exposed surfaces, and allow to stand for 10 minutes and flush.

ACCEPTED with COMMENTS in EPA Letter Dated:

FEB 3 2003 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the products registered under EPA Reg. No 11600-4

RECOMMENDED SANITIZATION DIRECTIONS

SANITIZATION – To sanitize hard, porous and nonporous environmental non-food contact surfaces, add 2 oz. of Sanox II per 4 gallon of water (1:256 dilution). Treated surfaces must remain wet for 1 minute (60 seconds).

Sanox II meets EPA's Sanitizer Test Standards for non-food contact sanitizers at 2 oz per 4 gallons of water (1:256 dilution, providing 200 ppm active) modified in the presence of 500 ppm hard water plus 5% organic serum.

Sanitizing Non-Food Contact Surfaces (such as floors, walls, tables, etc.) (1:256 dilution):

At 2 oz. of Sanox II per 4 gallon of water (½ oz. Sanox II per gallon of water) use level, Sanox II is an effective sanitizer against Staphylococcus aureus and Klebsiella pneumoniae on hard, porous and nonporous, environmental surfaces. Treated surfaces must remain wet for 60 seconds. Wipe dry with a sponge, mop or cloth; or, allow to air dry. Sanox II eliminated 99.9% of these organisms in 1 minute (60 seconds).

Sanitizing Barber / Salon Instruments and Tools (1:256 or 1:352 dilution):

Immerse precleaned barber / salon instruments and tools (such as combs, brushes, razors, scissors, blades and manicure instruments) in a use-solution of 2 oz. of Sanox II per 4 to 5 ½ gallons of water (or equivalent dilution) for a least 1 minute (60 seconds). Rinse thoroughly and dry before use. Prepare a fresh solution daily, or more often if solution becomes visibly diluted, cloudy or soiled.

Note: Plastic instruments and tools may remain immersed until ready to use. Stainless steel instruments must

9/10

be removed after 10 minutes, rinsed, dried, and kept in a clean non-contaminated receptacle.

Sanitizing Shoe Baths (1:256 dilution):

To prevent tracking harmful organisms into animal areas, shoe baths containing one inch of freshly made use-solution should be utilized. Scrape footwear and place in a 2 oz of Sanox II per 4 gallon of water solution for a minimum of 1 minute (60 seconds). Change the use-solution in the bath daily or more often if it appears diluted or soiled.

Refill Applicator Bottle For Disinfection and Sanitization:

Sanox II can be used with Conklin's Sanox II Refill Applicator Bottle. When using the Sanox II Refill Applicator Bottle, follow the use instructions on the Refill Applicator Bottle.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER. Keep Out of Reach of Children. Corrosive. Causes eye and skin damage. DO NOT get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
If in eyes	<ul style="list-style-type: none"> •Hold eye open and rinse slowly and gently with water for 15-20 minutes. •Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. •Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> •Take off contaminated clothing. •Rinse skin immediately with plenty of water for 15-20 minutes. •Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> •Call poison control center or doctor immediately for treatment advice. •Have person sip a glass of water if able to swallow. •Do not induce vomiting unless told to do so by the poison control center or doctor. •Do not give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN	
Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.	

ACCEPTED
with COMMENTS
in EPA Letter Dated:
FEB 3 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.
11660-4

(If container size is 5 gallon or greater, use the following ENVIRONMENTAL HAZARDS statements)

ENVIRONMENTAL HAZARDS

DO NOT discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. DO NOT discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

(If container size is greater than one gallon, use the following STORAGE AND DISPOSAL statements)

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE -

DO NOT store on side. DO NOT reuse empty container. Always replace cap securely. DO NOT substitute cap with pump or other dispensers. AVOID creasing or impacting of side walls.

PESTICIDE DISPOSAL -

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

10/10

CONTAINER DISPOSAL -

Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(if container size is one gallon or less, use the following STORAGE AND DISPOSAL statements)

STORAGE AND DISPOSAL

Store in areas inaccessible to children. Always replace cap securely. DO NOT substitute cap with pump or other dispensers. Securely wrap original container in several layers of newspaper and dispose of in trash. DO NOT reuse container.

For U.S. Health Hazard information, call 1-800-228-5635

HEALTH	3 (Serious)
FLAMMABILITY	0 (Minimal)
REACTIVITY	0 (Minimal)
PERSONAL PROTECTION	B

ACCEPTED
with COMMENTS
in EPA Letter Dated:

FEB 3 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 11600-4