# SAR!NE B1 - 30 **BROAD SPECTRUM FUNGICIDE BACTERICIDE - ALGAECIDE**

EPA Reg. #11620-3

CAUTION

**KEEP OUT OF THE** 

**REACH OF CHILDREN** 

Harmful if swallowed. Do not get

in eyes. Avoid skin contact. In case of contact flush with plenty of water for at least 15 minutes. If

eye irritution persists get medical

not discharge effluent to lakes,

streams or ponds. Do not contam-

ingte water by cleaning of equip-


ment or disposal of waste. Use

Do not reuse empty drum, Re-

only as directed on label.

and burying in a safe place.

food.



Active ingredient ..... Poly[oxyethylene(dimethyl ethylene dichloride] Inert ingredient ..... Weight per gallon ... Weight of active ingr

Sarine B1 30 is used to control algae, bacteria, and fungi in recirculating attention. Avoid contamination of commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime and other deposits. This product is taxic to fish. Do An initial slug addition of 2.00 to 5.00 fluid ounces of Sarine B1 30 per 1,000 gallons of water. to provide a concentration of 16 to 40 parts per million of Sarine B1 30 based on the total weight of water in the system is recommended. Subsequent slug additions of 0.50 to 5.00 fluid ounces of Sarine B1 30 per 1,000 gallons of water (4 to 40 parts per million of Sarine B1 30) should be employed every 2 to 5 days or as needed. The frequency turn to drum recorditioner or de- of addition depends upon the relative amount of bleedoff and the severity stroy by perforating or crushing of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

### MANUFACTURED BY

OLSEN CHEMICAL CO, INC

-62-64 E -26th Street + Paterson, N J 07514 + (201) 525-2600

# SARINE B1 30

Percent by weight
liminio)ethylene(dimethyliminio)-
30 percent
redient per gallon 2.67 pound

## APPLICATION