



11623-49

2-28-2000

1/2

Office of Pesticide Programs  
Registration Division (H7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

Number:  
11623-49      FEB 28 2000

NOTICE OF PESTICIDE:  
  x   Registration  
       Reregistration

Term of Issuance:  
Conditional

Name of Pesticide Product:  
Appollo Contact II

under FIFRA, as amended

Name and Address of Registrant (include ZIP Code):  
Apollo Industries, Inc.  
c/o Adams Technology Systems  
5145 Forest Run Trace - Suite B  
Alpharetta, GA 30022-4504

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Change the label by revising the EPA Registration Number to read, "EPA Reg. No. 10623-49".
3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Signature of Approving Official:  
*Marion J. Johnson, Jr.*  
Marion J. Johnson, Jr.

Date:  
February 28, 2000

ACCEPTED  
WITH COMMENTS  
In EPA Label dated:  
FEB 28 2000

Under Federal Insecticide,  
Fungicide, and Rodenticide Act,  
an amendment for the pesticide  
registered under EPA Reg. No.  
11623-49



# CONTACT II

**SBP-1382® INSECTICIDE  
AQUEOUS PRESSURIZED  
SPRAY 0.25%**

**AUTHORIZED BY USDA FOR USE IN EDIBLE PRODUCT  
AREAS OF OFFICIAL ESTABLISHMENTS OPERATING UNDER  
THE MEAT, POULTRY, SHELL EGG GRADING, AND EGG  
PRODUCTS INSPECTION PROGRAM.**

**ACTIVE INGREDIENT:**

\* Resmethrin ..... 0.25%

**OTHER INGREDIENTS:** ..... 99.75%  
..... 100.00%

\* [5-(phenylmethyl)-3-furanyl] methyl 2,2-dimethyl-3-(2-methyl-1-propenyl) cyclopropanecarboxylate - and - cis/trans isomers ratio max. 30% (±) cis and min. 70% (±) trans

AgrEvo Environmental Health Inc.'s SBP-1382® brand of resmethrin insecticide.

**PRECAUCION AL CONSUMIDOR:** Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente. (To the user: If you cannot read English, do not use this product until the label has been fully explained to you.)

EPA REG. NO. 11623-

EPA EST. NO. 11623-GA-1

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Side Panels For Additional  
Precautionary Statements

NET WEIGHT: 13 OZS. (369 g)

MANUFACTURED BY  
**APOLLO INDUSTRIES, INC.**

1850 S. COBB INDUSTRIAL BLVD.  
SMYRNA, GEORGIA 30082  
(770) 433-0210  
www.apolloind.com

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SHAKE WELL BEFORE USING**

Not for use on plants being grown for sale or other commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks or lawns and grounds.

For use in and around hotels, motels, homes, food handling establishments, packaging and processing plants, seed houses, commercial, industrial, office, institutional and storage buildings, milk handling areas of milk, cheese and ice cream plants, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, boats, ship cabins and hospitals. In hospitals, patients should be removed prior to treatment. Also for use in buses, automobiles, trucks, boxcars and ship cargo holds. Also for use in dog kennels and horse stables. In horse stables, not for application if horses will be used for food.

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray.

**INDOORS**

**FOR HOUSE FLIES, MOSQUITOES, GNATS, FLYING MOTHS (Millers), FRUIT FLIES AND VINEGAR FLIES:** Close all doors and windows. Point nozzle upwards; direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

**FOR CRAWLING INSECTS: Cockroaches (Waterbugs), Earwigs, Grain Mites, Sowbugs (Pillbugs), Silverfish (Firebrats), Spiders, Centipedes, Millipedes, Fleas, Ants, Booklice and Crickets:** Spray thoroughly and frequently into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, door frames, bookcases and storage areas. Spray directly on insects where possible. Repeat as necessary.

**TO KILL ANTS - INCLUDING FIRE ANTS:** Spray doors, around window frames, and other points of entry. Spray ant trails, nests, and hills contacting as many ants directly as possible.

**TO KILL BEDBUGS AND LICE:** Apply as a spot treatment to cracks and crevices around baseboards, floorboards, bedboards and walls. First, thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor pests, concentrating on tufts, folds, and seams. Repeat as necessary. For persistent infestations, consult professional applicator.

**TO PROTECT HOUSE PLANTS:** Spray infested African violets, philodendron, begonias, geraniums and coleus for control of *Aphids*, *Exposed Thrips*, *Adult Whiteflies*, *Spider Mites*, and *Fungus Gnats*. Use with caution. To ascertain possible phytotoxicity, spray only a few plants, wait 24 hours, and check for possible injury.

**FOR PROTECTION OF WOOLENS:** Apply only to garments, blankets, carpets, and other woollen fabrics before placing in storage for protection from larvae of webbing clothing moths and black carpet beetles.

Dry clean or wash garments and blankets before treating. Clean carpets, draperies and other fabrics by thoroughly brushing before treating. Hold the container 14 to 18 inches (30 to 46 cm) from the fabric and apply the spray while moving evenly in a forward and back motion across the fabric while covering one foot (30 cm) per second. Spray the entire fabric surface until it is slightly moist or damp to the touch. When possible treat both sides of the fabric. Do not soak or wet fabric. For maximum protection, place treated fabrics after drying in plastic or paper bags. Also treat cracks and crevices of closets, chests and trunks where these fabrics are being stored. Treat the cracks and crevices at the rate of 2 seconds per linear foot (30 cm). When applied as directed on fabric and in storage areas, this treatment will offer protection for 6 months. To insure protection, inspect the fabric in the storage area once every three months. Repeat application only when necessary. Avoid heavy applications. When possible make applications outdoors. If application is made indoors do not remain in treated areas. Ventilate the area after treatment is complete.

**OUTDOORS**

**TO KILL FLYING INSECTS:** Spray areas infested with *Mosquitoes*, *Gnats (Midges)*, *House Flies*, *Stable Flies*, *Deer Flies* and *Flying Moths (Millers)*. Follow precautions listed above. Hold container 18 to 24 inches (46 to 61 cm) from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard (0.76 cu. m) of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

**TO PROTECT ORNAMENTAL PLANTS:** To control *Aphids*, *Exposed Thrips*, *Adult Whiteflies*, *Leafhoppers*, *Gypsy Moths*, *Plant Bugs*, *Spittlebugs*

and *Japanese beetles* on smaller rhododendrons, pines, spruce, hemlock, spruce, dogwood, palm, mimosa, toyon, viburnum, euonymus, oaks, maples, locust, Forsythia and althea, and mimosa webworm larvae on mimosa, spray directly on insects. Hold container 18 inches (46 cm) from the infested plants. Spray in short bursts of one to three seconds or spray at the rate of one linear foot (30 cm) of growth per second. Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control. Spray peonies, iris, zinnia, snapdragons, ageratum, chrysanthemums, nasturtiums, daisy, geranium, marigolds, daphne, lathedera, iris, tobia, queen sago and broad leaf acacia for control of *aphids*, *exposed thrips*, *adult whiteflies*, *leafhoppers*, *plant bugs*, *spittlebugs*, and *Japanese beetles*. Spray as directed above.

**Ornamental Quince, Juniper, Cypress, Ornamental Mulberry, Ash, Cotoneaster, Rose, Camellia, Azalea, Pyracantha and Caenothus** may also be treated to control the above pests, but use with extreme caution. Certain varieties have shown some leaf burn under certain conditions of use such as heavy dosage. Repeat as necessary to obtain control.

**TO KILL TENT CATERPILLARS:** Application should be made when caterpillars are young and tents are first noticed (usually late spring). For best results, apply in the late afternoon or evening when caterpillars have returned to their tents. Aim spray directly at tents, and totally saturate on all sides. Repeat as necessary.

**TO KILL GYPSY MOTH CATERPILLARS:** Application should be made when caterpillars are young and first noticed (usually late spring). Spray plants evenly to the point of run-off, applying to both sides of leaves and other areas of plants where insects are suspected. For dense evergreens and shrubs, be sure spray reaches interior leaves and branches.

**TO CONTROL NUISANCE PESTS: CRICKETS, SILVERFISH, EARWIGS, SOWBUGS, CENTIPEDES AND MILLIPEDES:** Same as directed above for indoor use.

**ANTS:** Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

**FIRE ANTS:** Kills and temporarily reduces annoyance of these insects. To reduce annoyance from these ants and keep them from entering homes, spray patios, barbecue areas, driveways, door sills, window sills, other entrance areas, foundations of buildings, ant trails, around pools and areas where the ants are noticed. Spray until the area is wet contacting as many insects as possible. This treatment will not kill the colonies, but will reduce the immediate population for up to four hours. Repeat as necessary. Avoid contacting plants, shrubbery and large areas of lawns.

**SPIDERS:** (Including Black Widow and Brown Recluse): Direct short bursts of spray on spiders and repeat as necessary.

**WASPS, HORNETS, YELLOW JACKETS AND BEES:** Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary.

**PRECAUTIONARY STATEMENTS  
Hazards to Humans & Domestic Animals**

**CAUTION**

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packing plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, thoroughly with an effective cleaning compound and rinse with potable water. In the home, all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use. Remove pets, birds and cover fish aquariums before spraying.

**FIRST AID**

**IF IN EYES OR ON SKIN:** Flush with plenty of water. Get medical attention if irritation persists.

**ENVIRONMENTAL HAZARDS**

This pesticide is highly toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated sites may be hazardous to fish in adjacent waters. Do not contaminate water when disposing of equipment washwaters.

**PHYSICAL AND CHEMICAL HAZARDS**

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F (54.5°C) may cause bursting. Do not apply this water-based product in conduits, motor windings, junction and switch boxes or other electrical equipment because of possible shock hazards. Do not use on surfaces that can be harmed or stained by water.

**STORAGE & DISPOSAL**

Storage: Keep in locked storage area.  
Container Disposal: Replace cap and discard container in trash.

V110999

2/2