

SODIUM HYPOCHLORITE

CAUTION: KEEP OUT OF REACH OF CHILDREN

NEVER MIX WITH ACID AS
THIS WOULD RELEASE CHLORINE GAS
AVOID CONTACT WITH SKIN AND EYES

DO NOT DRINK
HARMFUL IF SWALLOWED

SEE ANTIDOTE STATEMENT ON SIDE PANEL

CONTENT

Active Ingredients: Sodium Hypochlorite 10.5% by weight.
Inert Ingredients: 89.5% by weight.

Directions for Use

Regrades with appropriate test kit and increase dosage as necessary to obtain required level of available chlorine.

ANTIDOTE

EXTERNAL — Wash with Water

INTERNAL — Drink mucilage, raw egg white, milk or rice gruel. Follow with emetic (tablespoon of mustard in glass of water) to induce vomiting.

CALL A PHYSICIAN

EPA Reg. No. 11632-1

11632-1 ALBAN DANIEL CHEMICAL CORPORATION
5012 - 46th Avenue, Hyattsville, Maryland 20781
927-1008

ACCEPTED
MAY 22 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. ~~11632-1~~

The
dir
cat
has
For
ide
for
obt
rep
ppr
que
the
pac
Kee
Pod
10
30
50
100

ORITE
OF CHILDREN
AS
RINE GAS
AND EYES

MED
SIDE PANEL

10.5% by weight.

SODIUM HYPOCHLORITE

Degrades with age. Use a test kit and increase dosage as necessary to obtain required level of available chlorine.

ANTIDOTE

EXTERNAL — Wash with Water
INTERNAL — Drink mucilage, raw egg white, milk or rice gruel. Follow with emetic (tablespoon of mustard in glass of water) to induce vomiting.

CALL A PHYSICIAN

EPA Reg. No. 11632-1

ACCEPTED
MAY 22 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO. ~~74123~~

11632-1
ALBAN DANIEL CHEMICAL CORPORATION
5012 - 46th Avenue, Hyattsville, Maryland 20781
927-9008

Directions For Use in Swimming Pools

The chlorine in swimming pools tends to dissipate in direct sunlight. Therefore it is best to make an application in the evening or preferably after use of the pool has been discontinued for the day.

For newly filled pools—to establish a chlorine residual and quantity indicated by size of pool below. Repeat dosage until a chlorine residual of 0.6 to 1.0 ppm is obtained as determined by use of a test kit. Thereafter, repeat dosage daily or as needed to maintain 0.6 to 1.0 ppm chlorine residual at all times. Test pool water frequently with a test kit. Before entering the pool, check the chlorine residual and if above 2.0 ppm, allow the pool to stand until the residual drops below 2.0 ppm.

Keep the pool PH in the 7.2 to 7.6 range.

Pool Capacity	Daily Dosage
10,000 gallons	2 quarts
30,000 gallons	5 quarts
50,000 gallons	2 gallons
100,000 gallons	4 gallons