

DIRECTIONS FOR USE

Avitrol is a pesticide for the control of certain species of pest birds. It is composed of a choice bird food impregnated with an active bird management chemical. Birds ingesting Avitrol react with distress symptoms and calls. By limiting the amount of bait available to relatively few birds, the remainder of the flock can be frightened away from most roosts and feeding sites with a minimum of mortality. For best results the applications should be made at the proper time and place under the direction of trained personnel.

Preceding with untreated feeds of the same composition as the Avitrol carrier is usually essential to the effective persistence of birds. Careful observation of bird habits should be made to establish proper feeding locations and to determine that no desirable or protected bird species are present that may feed on Avitrol. After pest bird feeding is established, immediately replace untreated grain with Avitrol. Repeat Avitrol treatment until bird numbers have been reduced to acceptable levels on the premises.

For the control of starlings in feedlots, corn bait is rendered much more attractive by mixing 4 parts with 1 part of peanut butter. This mixture must be used in elevated feeders out of reach of pets and farm animals.

Avitrol grain is lethal to most birds that ingest enough to cause flock-frightening symptoms. To obtain the desired minimal mortality, the distribution of Avitrol grain should be limited to scattered spot placements that will provide feeding opportunity for the necessary number of demonstrating birds. The remainder of the feeding area may be covered with untreated grain to provide desirable clutter.

AVITROL MUST NOT BE EXPOSED IN ANY MANNER THAT MAY ENDANGER DEERABLE AND PROTECTED BIRD FEEDS. IF THERE IS A QUESTION OF SUCH HAZARD, CONTACT LOCAL, STATE AND FEDERAL GAME, WILDLIFE, RECREATION AND FEDERAL BIRD MANAGEMENT AGENCIES BEFORE UNDERTAKING BIRD MANAGEMENT WITH AVITROL. IN VIOLATION OF CALIFORNIA LAWS THAT MAY PROHIBIT THE USE OF ANY TOXIC CHEMICAL IN BIRD CONTROL.

Avitrol

DOUBLE STRENGTH CORN CHOPS

For the Control of
CERTAIN BLACKBIRDS, COWBIRDS,
AND STARLINGS

Consult your local or nearest Federal, State or local government agencies or pest control operators. Not for use on the public.

Active Ingredients	
4-Aminopyridine	1.0%
Inert Ingredients	99.0%
Corn	98.6%
Hydrogen chloride	0.39%

Net Contents


AVITROL CORPORATION
TULSA, OKLAHOMA

CA No. 1649
Printed 3-15-64

CAUTION

KEEP OUT OF REACH OF CHILDREN

May produce toxic symptoms if swallowed or by prolonged and repeated contact with the skin. Handle with protective gloves. Wash thoroughly with soap and water after handling before eating or smoking. Change to clean clothing after handling. If swallowed, induce vomiting and call a physician. Do not feed to livestock or chickens. Do not mix with grain for livestock feed or for human consumption. Do not scatter on streams, lakes or ponds with the material. Do not allow this material to remain on property for more than 24 hours after application unless it is completely dried. Do not use where food is eaten or prepared.



NOTICE: Buyer and user must read and follow all directions and instructions on the label and on the container of this material.