# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. May cause irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

## **ENVIRONMENTAL** HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

This pesticide is toxic to birds and fish. Wildlife feeding on treated bait may be killed. Do not contaminate water by cleaning equipment or disposal of waste. Do not feed to livestock or poultry. Do not mix with grain for livestock or poultry feed

Ŋ

# DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### USE RESTRICTIONS

Do not apply to food or feed crops. Do not use where stored food or feed (such as grain or meat) might become contaminated.

Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. If there is a question of such a hazard, consult local, state and federal game authorities before undertaking bird management with Avitrol

Investigate local laws that may prohibit the use of any toxic chemical in bird control

### **STORAGE & DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

Storage: Store only in original container in a cool, dry location inaccessible to children and pets. Store apart from food or animal feed and protect against rodent penetration of the carton. Store apart from other pesticides, fertilizers, food or feed that may cause cross-contamination of odor or insect infestation.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If thes wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Fiber Drums with Liners: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

(a) A set of the se

# **RESTRICTED USE** PESTICIDE DUE TO ACUTE AVIAN TOXICITY

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification



# **DOUBLE STRENGTH CORN CHOPS**

Active ingredient: 4-Aminopyridine Inert Ingredients:

Total Avitrol is a poison with flock alarming properties used for the control of Red-Winged, Yellow-Headed, Rusty and Brevers Blackbirds, Grackles, Cowbirds and Starlings in, on or in the area of leedlots, structures, nesting, roosting and feeding sites, in such a way that a part of a flock may react and frighten the rest away. Birds that react and alarm a flock usually die.

## **KEEP OUT OF REACH OF CHILDREN** CAUTION FIRST AID

Have label with you when obtaining treatment advice.

	r		sary nu
If swallowed	<ul> <li>Call a poison control center, doctor, or 1-800-424-9300 immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> </ul>		sites wh not appl uneaten mals, it
lf in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.     Remove contact lenses, if present, after the first 5 minutes, then		product
	continue rinsing eye. • Call a poison control center, doctor or treatment advice.	\$80 <b>0</b> 424.9300 immediately (4	ground.
	Manufactured By AVITROL CORPOF 7644 East 46th Strr Tulsa, Oklahoma 7414 (918) 622-7763 • (800) 6 EPA Reg. No. 1164	ATION he Federal gebungielde, and Ro	dentidule ACC, he new tide
Avitrol Lo	EPA Est. 11649-OK t No	<	Notice: use and label ins

### PREBAITING DIRECTIONS

This product must not be applied where nontarget birds are feeding. Careful observations of the birds' feeding habits must therefore be made to establish proper feeding locations and to determine that no nontarget birds are feeding on the prebait. Prebaiting with untreated grains of the same composition as the Avitrol carrier is usually essential to the effective deterrence of the birds with Avitrol. The target species are primarily ground feeders and bait should be placed at ground level if possible.

For the control of Starlings in feedlots, corn bait may be rendered more attractive by mixing four parts with one part of peanut butter. This mixture must be kept off the ground, out of reach of cattle.

### **DILUTION DIRECTIONS**

Avitrol treated grain should be thoroughly mixed with untreated grain of the same composition as the Avitrol carrier at the dilution ratios indicated below. For the control of Blackbirds, Cowbirds, Grackles and Starlings with Avitrol, no dilution ratio less than one part treated to nine parts untreated is recommended

#### **BAITING DIRECTIONS**

To obtain minimal mortality, the distribution of Avitrol should be limited to scattered spot placements that will provide feeding opportunities only for the necesmber of target birds. After the birds' feeding has been established through prebaiting, untreated bait with diluted treated bait only at ere the target birds are actively feeding. Do y treated bait to inactive feeding sites. Where bait may be a hazard to other birds or anishould be picked up at the end of each day. is product is used in or around occupied buildk up and dispose of dead birds by burial. This must not applied where nontarget bird species

1.0%

99.0%

100.0%

d with peanut butter must be kept off the out of reach of cattle.

he first few days, baiting with Avitrol may successive applications until control is Note that retreatment may also be necesrainfall.

Buyer and/or user assumes all risks of /or handling of these materials contrary to structions

5/03