

DIRECTIONS FOR USE

GENERAL

Avitrol Concentrate pesticide may be used when directed below for the control of herring gulls at sanitary landfills and in, on, or near structures where they feed, nest or roost.

Avitrol Concentrate is lethal to most birds that ingest enough to cause flock frightening distress symptoms and die. By Avitrol, limiting the amount of bait available to relatively few birds, the remainder of the flock can be frightened away from the feed and feeding sites with a minimum of mortality. For best results, the applications must be made at the proper time and place under the direction of trained personnel.

PRE-BAITING

Pre-feeding with bread is essential to the effective baiting of birds with Avitrol. Careful observation is required to determine to establish proper feeding locations and to determine that non-target bird species are present in the area.

Tossing bread squares into the air sometimes attracts the birds. Bread tossed from a very slowly moving vehicle is often readily accepted. Bait can also be effectively broadcast by a person on foot. In this case, it is desirable to wear the same hat and clothing each time. As soon as the bait is applied, move away from the area. Early morning placement, before natural feeding occurs, is best. Continue pre-baiting until the pest birds have established their feeding pattern.

BAIT PREPARATION

To prepare the bait, cut each 264 square standard size bread into 12 equal pieces. Open and pour into the plastic bag, close by twisting the top or trapping as much air as possible. Tumble and shake the contents until all the powder is evenly distributed over the pieces of bread with none remaining on the walls of the bag. Handle treated bait with a gloved hand and remove with soap and water any of the powder that may have adhered to the skin.

DILUTION

Prior to baiting mix one part (24 squares) bread treated with Avitrol Concentrate with 10 additional parts (240 squares) of untreated bread of similar size. Use rubber glove if mixing is done by hand.

BAITING

After pest bird feeding is established immediately replace untreated bread with diluted Avitrol treated bread at those sites used for pre-baiting where target species are present. For each application of bait in flocks of over 5,000 broadcast the 264 bread squares (24 treated and 240 untreated) prepared from one bag of powder among eight widely separated sites. For flocks fewer than 5,000 use half of the above quantity. Repeat treatment until bird numbers have been reduced to acceptable levels on the premises. Note that retreatment may also be necessary after rain.

Additional Avitrol baits are available for the control of these and other pest birds. For list of baits and more details of material consult Avitrol brochures or manufacturer.

AVITROL

CONCENTRATE

For The Control Of

HERRING GULLS AT, IN, OR NEAR SANITARY LANDFILLS AND STRUCTURES WHERE GULLS FEED, NEST OR ROOST


Registered under the Pesticide Control Act, 1957, and the Pesticides Act, 1972, of the Government of the State of Oklahoma. Registered under the Pesticide Control Act, 1957, and the Pesticides Act, 1972, of the State of Oklahoma. Both Federal and State laws prohibit the sale of this product to anyone under the age of 18.

Avitrol Concentrate	25
Net Contents	280 g (10.0 oz)

AVITROL CORPORATION
TULSA, OKLAHOMA

EPA Reg No. 11649-10
U.S. Patent 3,150,041

EPA Est. 11649-OK-1



DANGER POISON

KEEP OUT OF REACH OF CHILDREN

Poisonous if swallowed or may produce toxic symptoms upon prolonged contact with the skin. Do not allow the dust to escape from the plastic bag during the preparation of the bait. Avoid inhalation or contact with eyes. Use protective gloves when handling the treated bait. Wash thoroughly with soap and water before eating or smoking. Do not contaminate lakes, streams, feed or foodstuffs with the material. Keep away from livestock, poultry or pets. Do not allow Avitrol treated bread to remain in uncontrolled places where it may be mistaken for food by people or animals if where any kind of contamination will result.

ANTIDOTE

CALL A PHYSICIAN AT ONCE

Induce vomiting by giving plenty of warm salt water (1 tablespoon of salt per glass), repeat until vomit fluid is clear. Have patient lie down and be quiet.

AVITROL

NOTICE: Buyer and/or user assumes all risks of use and/or handling of these materials contrary to label instructions.

ENVIRONMENTAL HAZARD CAUTIONS:

This product is toxic to birds and other wildlife. Birds feeding on treated area may be killed. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. Consult local state and federal game authorities before undertaking bird management with Avitrol. Investigate local laws that may prohibit the use of any toxic chemical in bird control.

STORAGE AND DISPOSAL

- 1. **PROHIBITIONS**
Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.
- 2. **PESTICIDE DISPOSAL**
Pesticide that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
- 3. **CONTAINER DISPOSAL**
Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place.
- 4. **GENERAL**
Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burying.