

Reg # 11649-10

PM-11

ACCEPTED

191

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Poisonous if swallowed. Harmful if absorbed through skin. May cause eye irritation. Do not breathe dust. Do not get in eyes, on skin, or on clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds and fish. Wildlife feeding on treated bait may be killed. Do not contaminate water by cleaning of equipment or disposal of waste. Do not feed to livestock or poultry. Do not mix with grain for livestock or poultry feed.

Pick up and dispose of dead birds by burial.

STORAGE AND DISPOSAL

1. PESTICIDE DISPOSAL

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

2. CONTAINER DISPOSAL

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

Do not use where food (grain or meat) might become contaminated.

Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. If there is a question of such hazard, consult local, state and Federal game authorities before undertaking bird management with Avitrol.

Investigate local laws that may prohibit the use of any toxic chemical in bird control.

Applicators must wear protective clothing, including long sleeves, gloves and respirators when using this product.

RESTRICTED USE PESTICIDE

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

AVITROL

Concentrate

Active Ingredient:	4-Aminopyridine	25.0%
Inert Ingredients:		75.0%
	Total	100.0%

Avitrol is a poison with flock alarming properties used for the control of Gulls at, in, near or in the vicinity of Sanitary Landfills, Airports and structures where Gulls feed, nest, loaf or roost, in such a way that a part of a flock may react and frighten the rest away. Birds that react and alarm a flock usually die.

Herring Gulls (*Larus argentatus*), as are all Gulls, are protected by law and treaty. Both Federal and State permits are required to use the Avitrol method of Gull control.

Avitrol Lot No. _____

KEEP OUT OF REACH OF CHILDREN

DANGER POISON

STATEMENT OF PRACTICAL TREATMENT

If swallowed: If the patient is unconscious, maintain breathing and heart beat (CPR: Cardiopulmonary resuscitation). Contact your local Poison Control Center, hospital or physician immediately.

If patient is conscious: Induce vomiting with syrup of Ipecac (if not available, stimulate the back of the throat with finger). Never give anything by mouth to an unconscious person. Contact your Poison Control Center, hospital or physician immediately.

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

If on skin: Wash thoroughly with soap and water.

See side panel for additional precautionary statements.

Manufactured By

AVITROL

Corporation
320 S. Boston
Tulsa, Oklahoma 74103
EPA Reg. No. 11649-10
EPA Est. 11649-0K-1

Net Contents: 0.281 oz. (8 grams)

1984
EPA Federal Insecticide,
Fungicide and Rodenticide Act
Registration for the pesticide
11649-10

Prebaiting Directions

This product must not be applied where nontarget birds are feeding. Careful observations of the birds' feeding habits must therefore be made to establish proper feeding locations and to determine that no nontarget birds are feeding on the prebait.

Prebaiting with bread is essential to the effective deterrence of birds with Avitrol. Tossing bread squares into the air sometimes attracts the birds. Bread tossed from a very slowly moving vehicle is often readily accepted. Bait can also be effectively broadcast by a person on foot. In this case it is desirable to wear the same hat and clothing each time. Immediately after the bait is applied, move away from the area. Early morning placement, before natural feeding occurs, is best. Continue prebaiting until the pest birds have established their feeding pattern.

Bait Preparation

To prepare the bait, cut each of two (2) standard slices of fresh bread into 12 equal pieces. Open and place into the plastic bag in which a portion of the product was shipped; close by twisting the top entrapping as much air as possible. Tumble and shake the contents until all the powder is evenly distributed over the pieces of bread with none remaining on the walls of the bag. Handle treated bait with a gloved hand and remove with soap and water any of the powder that may have adhered to the skin.

Dilution Directions

Prior to baiting, mix one part (24 squares) bread treated with Avitrol Concentrate with 10 additional parts (240 squares) of untreated bread of similar size. Use rubber glove if mixing is done by hand.

Baiting Directions

To obtain minimal mortality, the distribution of Avitrol should be limited to scattered spot placements that will provide feeding opportunities only for the necessary number of target birds. After the birds' feeding pattern has been established through prebaiting, replace untreated bait with diluted treated bait only at sites where the target birds are actively feeding. Do not apply treated bait to inactive feeding sites. Where uneaten bait may be a hazard to other birds or animals, it should be picked up at the end of each day. Do not allow Avitrol treated bread to remain in unprotected places where it may be mistaken for food by people or animals. This product must not be applied where nontarget bird species are feeding. For each application of bait to flocks of over 5,000, broadcast the 264 bread squares (24 treated and 240 untreated) prepared from one bag of powder among eight widely separated sites. For flocks fewer than 5,000, use half of the above quantity. Repeat treatment until bird numbers have been reduced to acceptable levels on the premises. Note that retreatment may also be necessary after rainfall.

Notice: Buyer and/or user assumes all risks of use and/or handling of these materials, contrary to label instructions.