

ACCEPTED

JUN 11 1984

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 11649-11

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS & DOMESTIC ANIMALS**

DANGER: Poisonous if swallowed. May be fatal if inhaled. Do not breathe dust. Wear a mask or pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and the National Institute for Occupational Safety and Health. Harmful if absorbed through skin. May cause eye irritation. Do not get in eyes, on skin, or on clothing. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds and fish. Wildlife feeding on treated bait may be killed. Do not contaminate water by cleaning of equipment or disposal of waste. Do not feed to livestock or poultry. Do not mix with grain for livestock or poultry feed.

Pick up and dispose of dead birds by burial.

STORAGE AND DISPOSAL

1. PESTICIDE DISPOSAL
Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

2. CONTAINER DISPOSAL
Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Use Restrictions

Do not use where food (grain or meat) might become contaminated. Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. If there is a question of such a hazard, consult local, state and Federal game authorities before undertaking bird management with Avitrol.

Investigate local laws that may prohibit the use of any toxic chemical in bird control.

Applicators must wear protective clothing, including hat, sleeves, gloves and respirators when using this product.

**RESTRICTED USE
PESTICIDE**

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

AVITROL**Powder Mix**

Active Ingredient:	4-Aminopyridine	50.0%
Inert Ingredients:		50.0%
	Total	100.0%

Avitrol is a poison with flock alarming properties used for the control of Starlings in, or in the immediate vicinity of feed lots in such a way that a part of a flock may react and frighten the rest away. Birds that react and alarm a flock usually die.

Avitrol Lot No. _____

KEEP OUT OF REACH OF CHILDREN

DANGER POISON

STATEMENT OF PRACTICAL TREATMENT

If swallowed: If the patient is unconscious, maintain breathing and heart beat (CPR: Cardiopulmonary resuscitation). Contact your local Poison Control Center, hospital or physician immediately.

If patient is conscious: Induce vomiting with syrup of Ipecac (if not available, stimulate the back of the throat with finger). Never give anything by mouth to an unconscious person. Contact your Poison Control Center, hospital or physician immediately.

If on skin: Wash thoroughly with soap and water.

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration. Get medical attention.

See side panel for additional precautionary statements.

Manufactured by

AVITROL

Corporation
320 - Boston
Tulsa, OK 74103
EPA Reg. No. 11649-11
EPA Est. 11549-OK-1
Net Contents 0.707 oz. (20 grams)

Prebaiting Directions

Prebaiting with the bait material to be used is usually essential to the effective deterrence of birds with Avitrol. This product must not be used where nontarget birds are feeding. Careful observations of the birds' feeding habits must therefore be made to establish proper feeding locations and to determine that no nontarget birds are feeding on the prebait. The Avitrol concentrate in this bag is for deposition on a suitable bait for the control of Starlings in and around feedlots. Livestock feed varies in different sections of the country and this provides a method of preparing alternative baits from attractive food materials.

Bait Preparation

To prepare bread as bait, cut each of ten (10) standard slices of fresh bread into 16 equal pieces. Open and place into the plastic bag; close by twisting the top entrapping as much air as possible. Tumble and shake the contents until all the powder is even distributed over the pieces of bread with none remaining on the walls of the bag.

To prepare an alternative bait, such as french fries, steam flaked corn, etc. as baits; introduce approximately two pounds of the selected alternative bait material into the plastic bag and proceed as directed above for bread. Handle treated bait with a gloved hand and remove with soap and water any of the powder that may have adhered to the skin.

Application Directions

Using a gloved hand, scatter the contents of each bag of prepared bait over piles of untreated bait of the same composition at a ratio of approximately one part treated to nine parts untreated in areas where prebait was well accepted.

Baiting Directions

To obtain minimal mortality, the distribution of Avitrol should be limited to scattered spot placements that will provide feeding opportunities only for the necessary number of target birds. After the birds' feeding pattern has been established through prebaiting, replace untreated bait with diluted treated bait only at sites where the target birds are actively feeding. Do not apply treated bait to inactive feeding sites. Early morning applications are recommended. Repeat treatment on succeeding days until control is obtained. Where untreated bait may be a hazard to other birds or animals, it should be picked up at the end of each day. Do not allow Avitrol treated bait to remain in unprotected places where it may be mistaken for food by people or animals. This product must not be applied where nontarget bird species are feeding. Note that retreatment may also be necessary after rainfall.

Notice: Buyer and/or user assumes all risks of use and/or handling of these materials contrary to label instructions.

PM 14

11649-11

page 1 of 1

ACCEPTED

JUN 11 1984

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 11649-11

RESTRICTED USE
PESTICIDE

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

AVITROL

Powder Mix

Active Ingredient: 4-Aminopyridine	50.0Z
Inert Ingredients:	50.0Z
Total	100.0Z

Avitrol is a poison with flock alarming properties used for the control of Starlings in, or in the immediate vicinity of feed lots in such a way that a part of a flock may react and frighten the rest away. Birds that react and alarm a flock usually die.

Avitrol Lot No. _____

KEEP OUT OF REACH OF CHILDREN

DANGER POISON

STATEMENT OF PRACTICAL TREATMENT

If swallowed: If the patient is unconscious, maintain breathing and heart beat (CPR: Cardiopulmonary resuscitation). Contact your local Poison Control Center, hospital or physician immediately.

If patient is conscious: Induce vomiting with syrup of Ipecac (if not available, stimulate the back of the throat with finger). Never give anything by mouth to an unconscious person. Contact your Poison Control Center, hospital or physician immediately.

If on skin: Wash thoroughly with soap and water.

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration. Get medical attention.

See side panel for additional precautionary statements.

Manufactured by

AVITROL
Corporation
320 S. Boston
Tulsa, OK 74103
EPA Reg. No. 11649-11
EPA Est. 11649-OK-1
Net Contents 0.707 oz. (20 grmm)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Poisonous if swallowed. May be fatal if inhaled. Do not breathe dust. Wear a mask or pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and the National Institute for Occupational Safety and Health. Harmful if absorbed through skin. May cause eye irritation. Do not get in eyes, on skin, or on clothing. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds and fish. Wild life feeding on treated bait may be killed. Do not contaminate water by cleaning of equipment or disposal of waste. Do not feed to livestock or poultry. Do not mix with grain for livestock or poultry feed.

Pick up and dispose of dead birds by burial.

STORAGE AND DISPOSAL

1. PESTICIDE DISPOSAL

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

2. CONTAINER DISPOSAL

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Use Restrictions

Do not use where food (grain or meat) might become contaminated. Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. If there is a question of such a hazard, consult local, state and Federal game authorities before undertaking bird management with Avitrol.

Investigate local laws that may prohibit the use of any toxic chemical in bird control.

Applicators must wear protective clothing, including long sleeves, gloves and respirators when using this product.

Prebaiting Directions

Prebaiting with the bait material to be used is usually essential to the effective deterrence of birds with Avitrol. This product must not be used where nontarget birds are feeding. Careful observations of the birds' feeding habits must therefore be made to establish proper feeding locations and to determine that no nontarget birds are feeding on the prebait. The Avitrol concentrate in this bag is for deposition on a suitable bait for the control of Starlings in and around feedlots. Livestock feed varies in different sections of the country and this provides a method of preparing alternative baits from attractive food materials.

Bait Preparation

To prepare bread as bait, cut each of ten (10) standard slices of fresh bread into 16 equal pieces. Open and place into the plastic bag; close by twisting the top entrapping as much air as possible. Tumble and shake the contents until all the powder is even distributed over the pieces of bread with none remaining on the walls of the bag.

To prepare an alternative bait, such as french fries, steam flaked corn, etc. as baits; introduce approximately two pounds of the selected alternative bait material into the plastic bag and proceed as directed above for bread. Handle treated bait with a gloved hand and remove with soap and water any of the powder that may have adhered to the skin.

Dilution Directions

Using a gloved hand, scatter the contents of each bag of prepared bait over piles of untreated bait of the same composition at a ratio of approximately one part treated to nine parts untreated in areas where prebait was well accepted.

Baiting Directions

To obtain minimal mortality, the distribution of Avitrol should be limited to scattered spot placements that will provide feeding opportunities only for the necessary number of target birds. After the birds' feeding pattern has been established through prebaiting, replace untreated bait with diluted treated bait only at sites where the target birds are actively feeding. Do not apply treated bait to inactive feeding sites. Early morning applications are recommended. Repeat treatment on succeeding days until control is obtained. Where uneaten bait may be a hazard to other birds or animals, it should be picked up at the end of each day. Do not allow Avitrol treated bait to remain in unprotected places where it may be mistaken for food by people or animals. This product must not be applied where nontarget bird species are feeding. Note that retreatment may also be necessary after rainfall.

Notice: Buyer and/or user assumes all risks of use and/or handling of these materials contrary to label instructions.

BEST AVAILABLE COPY