

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed. May cause irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to birds and fish. Wildlife feeding on treated bait may be killed. Do not contaminate water by cleaning of equipment or disposal of waste. Do not feed to livestock or poultry. Do not mix with grain for livestock or poultry feed.

**STORAGE AND DISPOSAL****1. PESTICIDE DISPOSAL**

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

**2. CONTAINER DISPOSAL**

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Use Restrictions**

Do not use where food (grain or meat) might become contaminated.

Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. If there is a question of such hazard, consult local, state and Federal game authorities before undertaking bird management with Avitrol.

Investigate local laws that may prohibit the use of any toxic chemical in bird control.

**RESTRICTED USE  
PESTICIDE**

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

**AVITROL**

Corn Chops - 99

Active Ingredient: 4-Aminopyridine	0.03%
Inert Ingredients:	99.97%
Total	100.0%

Avitrol is a poison with flock alarming properties used for the control of Common Grackles, Red-Winged, Yellow-Headed Blackbirds, Cowbirds and Starlings that depredate ripening ears of field and sweet corn; in such a way that a part of a flock may react and frighten the rest away. Birds that react and alarm a flock usually die.

**KEEP OUT OF REACH OF CHILDREN****CAUTION****STATEMENT OF PRACTICAL TREATMENT**

If swallowed: If the patient is unconscious, maintain breathing and heart beat (CPR: Cardiopulmonary resuscitation). Contact your local Poison Control Center, hospital or physician immediately.

If patient is conscious: Induce vomiting with syrup of Ipecac (if not available, stimulate the back of the throat with finger). Never give anything by mouth to an unconscious person. Contact your Poison Control Center, hospital or physician immediately.

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

See side panel for additional precautionary statements.

Manufactured by

**AVITROL**  
Corporation  
320 S. Boston  
Tulsa, OK 74103

EPA Reg. No. 11649-12

EPA Est. \_\_\_\_\_  
Net Contents 50 lbs.

**Pre-treatment Considerations**

Avitrol Corn Chops-99 is composed of medium chopped corn coated with an active bird management chemical. Birds ingesting Avitrol react with distress symptoms and calls. By limiting the amount of bait available to relatively few birds, the remainder of the flock can be frightened away from feeding sites with a minimum of mortality.

When bird damage become apparent, careful observations of the flight patterns should be made to establish feeding locations. See that no desirable or protected birds are present that may be affected by Avitrol.

**Application Directions**

This product must be applied in accordance with 40 CFR Part 170. Confine treatment to areas 50 feet in from field edge. Do not apply more than four applications per crop season. No application rate in excess of .0144 ounces of 4-aminopyridine per acre on the portion being treated may be used.

Using calibrated equipment, apply the bait evenly by air or by ground in swaths 10 to 15 rows wide and 30 to 35 rows between swaths. The amount for one treatment is one pound per acre based on the total area (applied at 3 pounds per acre on treated portion). Repeat after a heavy rain (.5 inches) or when the material becomes depleted. For field corn, three or four treatments may be necessary when bird damage is severe. For sweet corn, because of the shorter damage season (7 to 10 days), two treatments are normally sufficient. Treatment areas or swaths should be alternated so that the total material applied in a season is evenly distributed over the field.

It is important that Avitrol Corn Chops-99 be widely scattered and the recommended amount per acre be strictly observed. The single treated kernels, 1 in 100, are toxic to blackbirds but are so thinly distributed by bulking with clean grain that very few birds are exposed. Applied in this way it is not likely that enough treated grain can be found to constitute a hazard to large game birds such as pheasants.

Notice: Buyer and/or user assumes all risks of use and/or handling of these materials contrary to label instructions.

