

Reg # 11649-12 PM-14 142

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. May cause irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds and fish. Wildlife feeding on treated bait may be killed. Do not contaminate water by cleaning equipment or disposal of waste. Do not feed to livestock or poultry. Do not mix with grain for livestock or poultry feed.

RESTRICTED USE PESTICIDE

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until dusts have settled. Protective clothing means at least, Respirator, Rubber Gloves, Lightweight Protective Coveralls and Lightweight Boots.

Because certain states may require more restrictive re-entry intervals for various crops treated by this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and/or oral warnings should include: CAUTION: This area was treated with the pesticide AVITROL CORN CHOPS-99 on (date of application). Do not enter without appropriate protective clothing until dusts have settled. In case of accidental exposure, refer to "Statement of Practical Treatment."

USE RESTRICTIONS

Do not use where food (grain or meat) might become contaminated.

Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. If there is a question of such hazard, consult local, state and federal game authorities before undertaking bird management with Avitrol.

Investigate local laws that may prohibit the use of any toxic chemical in bird control.

CORN CHOPS-99

Active Ingredient: 4-Aminopyridine 0.03%
Inert ingredients: 99.97%
100.0%

Avitrol is a poison with flock alarming properties used for the control of Common Grackles, Red-Winged, Yellow-Headed Blackbirds, Cowbirds and Starlings that depredate ripening ears of field and sweet corn; in such a way that a part of a flock may react and frighten the rest away. Birds that react and alarm a flock usually die.

with comments in EPA Letter Dated

DEC 27 1988

Under the Federal Insecticide, Fungicide, and Herbicide Act, amended, for the pesticide registered under EPA Reg. No.

11649-12

KEEP OUT OF REACH OF CHILDREN CAUTION STATEMENT OF PRACTICAL TREATMENT

If swallowed: If the patient is unconscious, maintain breathing and heart beat (CPR: Cardiopulmonary resuscitation). Contact your local Poison Control Center, hospital or physician immediately.

If patient is conscious: Induce vomiting with syrup of Ipecac (if not available, stimulate the back of the throat with finger). Never give anything by mouth to an unconscious person. Contact your Poison Control Center, hospital or physician immediately.

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

See side panel for additional precautionary statements.

BEST AVAILABLE COPY

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

1. STORAGE INSTRUCTIONS:

Store in original container only. Store apart from food or animal feed and protect against rodent penetration of the carton. Store apart from other pesticides, fertilizers, food or feed that may cause cross-contamination of odor

ENDANGERED SPECIES RESTRICTIONS

The following restrictions apply to use of this product after February 1, 1988.

Before using this pesticide on corn, in the counties listed below, you must obtain the PESTICIDE USE BULLETIN FOR PROTECTION OF ENDANGERED SPECIES for the county in which the product is to be used. The bulletin is available from your County Extension Agent, State Fish and Game Office, or your pesticide dealer. Use of this product in a manner inconsistent with the PESTICIDE USE BULLETIN FOR PROTECTION OF ENDANGERED SPECIES is a violation of Federal laws.

CALIFORNIA
COLUSA, INYO, LOS ANGELES, MERCED, ORANGE, RIVERSIDE, SAN BERNARDINO, SAN DIEGO, SANTA BARBARA, STANISLAUS, SUTTER, AND VENTURA

FLORIDA
ALACHUA, BAKER, BRADFORD, BREVARD, BROWARD, CHARLOTTE, CITRUS, CLAY, COLLIER, COLUMBIA, DADE, DE SOTO, DIXIE, DUVAL, FLAGLER, GADSDEN, GILCHRIST, GLADES, HARDEE, HENDRY, HERNANDO, HIGHLANDS, HILLSBOROUGH, INDIAN RIVER, JEFFERSON, LAFAYETTE, LAKE, LEE, LEON, LEVY, MADISON, MANATEE, MARION, MARTIN, MONROE, NASSAU, ORANGE, OKEECHOBEE, OSCEOLA, PALM BEACH, PASCO, PINELLAS, POLK, PUTNAM, ST. JOHNS, ST. LUCIE, SARASOTA, SEMINOLE, SUMTER, SUWANNEE, TAYLOR, UNION, VOLUSIA AND WAKULLA

GEORGIA
BRANTLEY, BRYAN, BULLOCH, BURKE, CAMDEN, CANDLER, CHARLTON, CHATHAM, EFFINGHAM, EMANUEL, EVANS, GLASCOCK, GLYNN, JEFFERSON, JENKINS, JOHNSON, LIBERTY, LONG, MCINTOSH, PIERCE, RICHMOND, SCREVEN, WARE, WASHINGTON AND WAYNE

KANSAS
CLARK, COMANCHE, MEADE AND STAFFORD

MONTANA
GARFIELD, MCCONE, SHERIDAN AND VALLEY

NEBRASKA
BOYD, BROWN, BUFFALO, BUTLER, CASS, CEDAR, COLFAX, DAWSON, DODGE, DOUGLAS, HALL, HAMILTON, HOLT, HOWARD, KEARNEY, KEWA PAHA, KNOX, MERRICK, NANCE, PHELPS, PLATTE, POLK, ROCK, SAPPY AND SAUNDERS

NORTH DAKOTA
EANSON, BOTTINEAU, BURKE, BURLEIGH, DIVIDE, DUNN, LIDDY, EMMONS, FOSTER, KIDDER, LOGAN, MCHENRY, MCINTOSH, MCKENZIE, MCLEAN, MERCER, MORTON, MOUNTAIN, NELSON, OLIVER, PIERCE, RAMSEY, RANVILLE, ROLETTE, SHERIDAN, SIOUX, STUTSMAN, TOWNER, WARD, WELLS AND WYOMING

SOUTH CAROLINA
AIKEN, BARNWELL, BEAUFORT, BERKELY, CHARLESTON, COLLETON, DORCHESTER, GEORGETOWN, HAMPTON, Horry, JASPER AND MARION

SOUTH DAKOTA
CLAY, HAAKON, HUGHS, POTTER, STANLEY, SULLY, UNION, VALMORRIS, YANKTON AND ZIEBACH

TEXAS
ARANSAS, AUSTIN, CAMERON, COLORADO, FORT BEND, GOLIAD, REFUGIO AND VICTORIA

PRETREATMENT CONSIDERATIONS

Avitrol Corn Chops-99 is composed of medium chopped corn coated with an active bird management chemical. Birds ingesting Avitrol react with distress symptoms and calls. By limiting the amount of bait available to relatively few birds, the remainder of the flock can be frightened away from feeding sites with a minimum of mortality.

When bird damage becomes apparent, careful observations of the flight patterns should be made to establish feeding locations. See that no desirable or protected birds are present that may be affected by Avitrol.

APPLICATION DIRECTIONS

This product must be applied in accordance with 40 CFR Part 170. Confine treatment to areas 50 feet in from field edge. Do not apply more than four applications per crop season. No application rate in excess of .0144 ounces of 4-aminopyridine per acre on the portion being treated may be used.

Using calibrated equipment, apply the bait evenly by air or by ground in swaths 10 to 15 rows wide and 30 to 35 rows between swaths. The amount for one treatment is one pound per acre based on the total area (applied at 3 pounds per acre on treated portion). Repeat after a heavy rain (5 inches) or when the material becomes depleted. For field corn, three or

contaminated water by leaching equipment or disposal of waste. Do not feed to livestock or poultry. Do not mix with grain for livestock or poultry feed.

AVITROL CORPORATION

292

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until dusts have settled. Protective clothing means at least, Respirator, Rubber Gloves, Lightweight Protective Coveralls and Lightweight Boots.

Because certain states may require more restrictive re-entry intervals for various crops treated by this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and/or oral warnings should include: CAUTION: This area was treated with the pesticide AVITROL CORN CHOPS-99 on (date of application). Do not enter without appropriate protective clothing until dusts have settled. In case of accidental exposure, refer to "Statement of Practical Treatment."

USE RESTRICTIONS

Do not use where food (grain or meat) might become contaminated.

Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. If there is a question of such hazard, consult local, state and federal game authorities before undertaking bird management with Avitrol.

Investigate local laws that may prohibit the use of any toxic chemical in bird control.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

1. STORAGE INSTRUCTIONS:

Store in original container only. Store apart from food or animal feed and protect against rodent penetration of the carton. Store apart from other pesticides, fertilizers, food or feed that may cause cross-contamination of odor or insect infestation.

2. DISPOSAL INSTRUCTIONS:

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinseate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and Local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

CORN CHOPS—99

Active ingredient: 4-Aminopyridine 0.03%
Inert ingredients: 99.97%
100.0%

Avitrol is a poison with flock alarming properties used for the control of Common Grackles, Red-Winged, Yellow-Headed Blackbirds, Cowbirds and Starlings that deplete ripening ears of field and sweet corn; in such a way that a part of a flock may react and frighten the rest away. Birds that react and alarm a flock usually die.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

DEC 27 1988

Under the Federal Insecticide, Fungicide, and Rodenticide Act, amended, for the pesticide registered under EPA Reg. No.

11649-12

KEEP OUT OF REACH OF CHILDREN CAUTION STATEMENT OF PRACTICAL TREATMENT

If swallowed: If the patient is unconscious, maintain breathing and heart beat (CPR: Cardiopulmonary resuscitation). Contact your local Poison Control Center, hospital or physician immediately.

If patient is conscious: Induce vomiting with syrup of Ipecac (if not available, stimulate the back of the throat with finger). Never give anything by mouth to an unconscious person. Contact your Poison Control Center, hospital or physician immediately.

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

See side panel for additional precautionary statements.

BEST AVAILABLE COPY

Manufactured By
AVITROL CORPORATION
Tulsa, Oklahoma 74145
EPA Reg. No. 11649-12
EPA Est. 11649-OK-1

Lot No. _____

Net Contents _____ Pounds

ALACHUA, BAKER, BRADFORD, BREVARD, BROWARD, CHARLOTTE, CITRUS, CLAY, COLLIER, COLUMBIA, DADE, DE SOTO, DIXIE, DUVAL, FLAGLER, GADSDEN, GILCHRIST, GLADES, HARDEE, HENDRY, HERNANDO, HIGHLANDS, HILLSBOROUGH, INDIAN RIVER, JEFFERSON, LAFAYETTE, LAKE, LEE, LEON, LEVY, MADISON, MANATEE, MARION, MARTIN, MONROE, NASSAU, ORANGE, OKEECHOBEE, OSCEOLA, PALM BEACH, PASCO, PINELLAS, POLK, PUTNAM, ST. JOHNS, ST. LUCIE, SARASOTA, SEMINOLE, SUMTER, SUWANNEE, TAYLOR, UNION, VOLUSIA AND WAKULLA

GEORGIA

BRANTLEY, BRYAN, BULLOCH, BURKE, CAMDEN, CANDLER, CHARLTON, CHATHAM, EFFINGHAM, EMANUEL, EVANS, GLASCOCK, GLYNN, JEFFERSON, JENKINS, JOHNSON, LIBERTY, LONG, MCINTOSH, PIERCE, RICHMOND, SCREVEN, WARE, WASHINGTON AND WAYNE

KANSAS

CLARK, COMANCHE, MEADE AND STAFFORD

MONTANA

GARFIELD, MCCONE, SHERIDAN AND VALLEY

NEBRASKA

BOYD, BROWN, BUFFALO, BUTLER, CASS, CEDAR, COLFAX, DAWSON, DODGE, DOUGLAS, HALL, HAMILTON, HOLT, HOWARD, KEARNEY, KEYSER, KNOX, MERRICK, NANCE, PHELPS, PLATTE, POLK, ROCK, SARPY AND SAUNDERS

NORTH DAKOTA

BANSON, BOTTINEAU, BURKE, BURLEIGH, DIVIDE, DUNN, EDDY, EMMONS, FOSTER, KIDDER, LOGAN, MCHENRY, MCINTOSH, MCKENZIE, MCLEAN, MERCER, MORTON, MOUNTAIN, NELSON, OLIVER, PIERCE, RAMSEY, RAMVILLE, POLETTE, SHERIDAN, SIOUX, STUTSMAN, TOWNER, WARD, WELLS AND WILLIAMS

SOUTH CAROLINA

AIKEN, BARNWELL, BEAUFORT, BERKELY, CHARLESTON, COLLETON, DORCHESTER, GEORGETOWN, HAMPTON, Horry, JASPER AND MARION

SOUTH DAKOTA

CLAY, HAAKON, HIGHS, POTTER, STANLEY, SULLY, UNION, WALWORTH, YANKTON AND ZIEBACH

TEXAS

ARANSAS, AUSTIN, CAMERON, COLORADO, FORT BEND, GOLIAD, REFUGIO AND VICTORIA

PRETREATMENT CONSIDERATIONS

Avitrol Corn Chops-99 is composed of medium chopped corn coated with an active bird management chemical. Birds ingesting Avitrol react with distress symptoms and calls. By limiting the amount of bait available to relatively few birds, the remainder of the flock can be frightened away from feeding sites with a minimum of mortality.

When bird damage becomes apparent, careful observations of the flight patterns should be made to establish feeding locations. See that no desirable or protected birds are present that may be affected by Avitrol.

APPLICATION DIRECTIONS

This product must be applied in accordance with 40 CFR Part 170. Confine treatment to areas 50 feet in from field edge. Do not apply more than four applications per crop season. No application rate in excess of .0144 ounces of 4-aminopyridine per acre on the portion being treated may be used.

Using calibrated equipment, apply the bait evenly by air or by ground in swaths 10 to 15 rows wide and 30 to 35 rows between swaths. The amount for one treatment is one pound per acre based on the total area (applied at 3 pounds per acre on treated portion). Repeat after a heavy rain (.5 inches) or when the material becomes depleted. For field corn, three or four treatments may be necessary when bird damage is severe. For sweet corn, because of the shorter damage season (7 to 10 days), two treatments are normally sufficient. Treatment areas or swaths should be alternated so that the total material applied in a season is evenly distributed over the field.

It is important that Avitrol Corn Chops-99 be widely scattered, and the recommended amount per acre be strictly observed. The single treated kernels, 1 in 100, are toxic to blackbirds but are so thinly distributed by bulk with clean grain that very few birds are exposed. Applied in this way it is not likely that enough treated grain can be found to constitute a hazard to large game birds such as pheasants.

Notice: Buyer and/or user assumes all risks of use and/or handling of these materials contrary to label instructions.