

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. May cause eye irritation. Wash thoroughly with soap and water after handling.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to birds and fish. Wildlife feeding on treated bait may be killed. Do not contaminate water by cleaning equipment or disposal of waste. Do not feed to livestock or poultry. Do not mix with grain for livestock or poultry feed.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**RE-ENTRY STATEMENT**

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until dusts have settled. Protective clothing means at least Respirator, Rubber Goggles, Lightweight Protective Coveralls and Lightweight Boots.

Because certain states may require more restrictive re-entry intervals for various crops treated by this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and/or oral warnings should include **CAUTION**. This area was treated with the pesticide **AVITROL CORN CHOPS-99** on (date of application). Do not enter without appropriate protective clothing until dusts have settled. In case of accidental exposure, refer to "Statement of Practical Treatment."

**USE RESTRICTIONS**

Do not use where food (grain or meal) might become contaminated.

Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. If there is a question of such hazard, consult local, state and federal game authorities before undertaking bird management with Avitrol.

Investigate local laws that may prohibit the use of any toxic chemical in bird control.

**STORAGE & DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**1. STORAGE INSTRUCTIONS:**  
Store in original container only. Store apart from food or animal feed and protect against rodent penetration of the carton. Store apart from other pesticides, fertilizers, food or feed that may cause cross-contamination of odor or insect infestation.

**2. DISPOSAL INSTRUCTIONS:**  
Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and Local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

**RESTRICTED USE PESTICIDE**

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.



**CORN CHOPS—99**

Active Ingredient 4-Aminopyridine	0.03%
Inert ingredients	99.97%
Total	100.0%

Avitrol is a poison with flock alarming properties used for the control of Common Grackles, Red-Winged, Yellow-Headed Blackbirds, Cowbirds and Starlings that depredate ripening ears of field and sweet corn, in such a way that a part of a flock may react and frighten the rest away. Birds that react and alarm a flock usually die.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

If swallowed: If the patient is unconscious, maintain breathing and heart beat (CPR Cardiorespiratory resuscitation). Contact your local Poison Control Center, hospital or physician immediately.

If patient is conscious: Induce vomiting with syrup of Ipecac (if not available, stimulate the back of the throat with finger). Never give anything by mouth to an unconscious person. Contact your Poison Control Center, hospital or physician immediately.

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

See side panel for additional precautionary statements.

**BEST AVAILABLE COPY**

Manufactured By  
**AVITROL CORPORATION**

Tulsa, Oklahoma 74145  
EPA Reg No 11649-12  
EPA Est 11649-OK-1

Lot No


**ENDANGERED SPECIES RESTRICTIONS**

Do not use where food (grain or meal) might become contaminated. Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. If there is a question of such hazard, consult local, State and Federal game authorities before undertaking bird management with Avitrol. Investigate local laws that may prohibit the use of any toxic chemical in bird control.

**PRETREATMENT CONSIDERATIONS**

Avitrol Corn Chops-99 is composed of medium chopped corn coated with an active bird management chemical. Birds ingesting Avitrol react with distress symptoms and calls. By limiting the amount of bait available to relatively few birds, the remainder of the flock can be frightened away from feeding sites with a minimum of mortality.

When bird damage becomes apparent, careful observations of the flight patterns should be made to establish feeding locations. See that no desirable or protected birds are present that may be affected by Avitrol.



**APPLICATION DIRECTIONS**

This product must be applied in accordance with 40 CFR Part 170. Confine treatment to areas 50 feet in from field edge. Do not apply more than four applications per crop season. No application rate in excess of 0.144 ounces of 4-aminopyridine per acre on the portion being treated may be used.

Using calibrated equipment, apply the bait evenly by air or by ground in swaths 10 to 15 rows wide and 30 to 35 rows between swaths. The amount for one treatment is one pound per acre based on the total area (applied at 3 pounds per acre on treated portion). Repeat after a heavy rain (5 inches) or when the material becomes depleted. For field corn, three or four treatments may be necessary when bird damage is severe. For sweet corn, because of the shorter damage season (7 to 10 days), two treatments are normally sufficient. Treatment areas or swaths should be alternated so that the total material applied in a season is evenly distributed over the field.

It is important that Avitrol Corn Chops 99 be widely scattered and the recommended amount per acre be strictly observed. The single treated kernels, 1 in 100, are toxic to blackbirds but are so thinly distributed by bulking with clean grain that very few birds are exposed. Applied in this way it is not likely that enough treated grain can be found to constitute a hazard to large game birds such as pheasants.