

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. May cause eye irritation. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

This pesticide is toxic to birds and fish. Wildlife feeding on treated bait may be killed. Do not contaminate water by cleaning equipment or disposal of waste. Do not feed to livestock or poultry. Do not mix with grain for livestock or poultry feed.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until dusts have settled. Protective clothing means at least: Respirator, Rubber Gloves, Lightweight Protective Coveralls and Lightweight Boots.

Because certain states may require more restrictive re-entry intervals for various crops treated by this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written and/or oral warnings should include: CAUTION: This area was treated with the pesticide AVITROL FC CORN CHOPS-99 on (date of application). Do not enter without appropriate protective clothing until dusts have settled. In case of accidental exposure, refer to "First Aid."

USE RESTRICTIONS

This product may be used only in fields of ripening sweet corn, field corn, or sunflowers as a toxicant/frightening agent to mitigate damage to these crops caused by flocks of common grackles (*Quiscalus quiscula*), red-winged blackbirds (*Agelaius phoeniceus*), yellow-headed blackbirds (*Xanthocephalus*), brown-headed cowbirds (*Molothrus ater*), and starlings (*Sturnus vulgaris*).

Do not use where stored food or feed (such as grain or meat) might become contaminated.

Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. If there is a question of such hazard, consult local, state and federal game authorities before undertaking bird management with Avitrol.

Investigate local laws that may prohibit the use of any toxic chemical in bird control.

Notice: Buyer and/or user assumes all risks of use and/or handling of these materials contrary to label instructions.

RESTRICTED USE PESTICIDE

DUE TO ACUTE AVIAN TOXICITY

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.



FC CORN CHOPS—99

| | |
|------------------------------------|--------|
| Active ingredient: 4-Aminopyridine | 0.03% |
| Inert Ingredients: | 99.97% |
| Total | 100.0% |

Avitrol is a poison with flock alarming properties used for the control of Common Grackles, Red-Winged, Yellow-Headed Blackbirds, Cowbirds and Starlings that deplete ripening ears of field and sweet corn and ripening sunflower fields in such a way that a part of a flock may react and frighten the rest away. Birds that react and alarm a flock usually die.

KEEP OUT OF REACH OF CHILDREN CAUTION FIRST AID

Have label with you when obtaining treatment advice.

| | |
|---------------------|---|
| If swallowed | <ul style="list-style-type: none"> Call a poison control center, doctor, or 1-800-424-9300 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. |
| If in eyes | <ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center, doctor or 1-800-424-9300 immediately for treatment advice. |

ACCEPTED
FEB 5 2004
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticide AVITROL FC CORN CHOPS-99
EPA Reg. No. 11649-12

Manufactured By
AVITROL CORPORATION
7644 East 46th Street
Tulsa, Oklahoma 74145-6370
(918) 622-7763 • (800) 633-5069
EPA Reg. No. 11649-12

EPA Est. 11649-OK-1

Lot No.

Net Contents _____ Pounds

DIRECTIONS FOR USE (continued)

PRETREATMENT CONSIDERATION

Avitrol FC Corn Chops-99 is composed of medium chopped corn coated with an active bird management chemical. Birds ingesting Avitrol react with distress symptoms and calls. By limiting the amount of bait available to relatively few birds, the remainder of the flock can be frightened away from feeding sites with a minimum of mortality.

When bird damage becomes apparent, careful observations of the flight patterns should be made to establish feeding locations. See that no desirable or protected birds are present that may be affected by Avitrol.

APPLICATION DIRECTIONS

This product must be applied in accordance with 40 CFR Part 170. Confine treatment to areas 50 feet in from field edge. For corn do not apply more than four applications per crop season. For sunflowers do not apply more than five applications per season. No application rate in excess of .0144 ounces of 4-aminopyridine per acre on the portion being treated may be used.

Using calibrated equipment, apply the bait evenly by air or by ground in swaths 10 to 15 rows wide and 30 to 35 rows between swaths. The amount for one treatment is one pound per acre based on the total area (applied at 3 pounds per acre on treated portion). Repeat after a heavy rain (.5 inches) or when the material becomes depleted. For field corn, three or four treatments may be necessary when bird damage is severe. For sweet corn, because of the shorter damage season (7 to 10 days), two treatments are normally sufficient. Treatment areas or swaths should be alternated so that the total material applied in a season is evenly distributed over the field.

It is important that Avitrol FC Corn Chops-99 be widely scattered and the recommended amount per acre be strictly observed. The single treated kernels, 1 in 100, are toxic to blackbirds but are so thinly distributed by bulking with clean grain that very few birds are exposed. Applied in this way it is not likely that enough treated grain can be found to constitute a hazard to large game birds such as pheasants. After October 1, additional care should be taken in baiting with Avitrol, since migratory grain eating birds may be in the area in increased numbers.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store only in original container in a cool, dry location inaccessible to children and pets. Store apart from food or animal feed and protect against rodent penetration of the carton. Store apart from other pesticides, fertilizers, food or feed that may cause cross-contamination of odor or insect infestation.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Fiber Drums with Liners: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.