

PRECAUTIONARY STATEMENTS

HAZARD TO MAN & DOMESTIC ANIMALS

CAUTION: Irritant if swallowed. May cause irritation. Wash contact with eyes or skin immediately with soap and water after handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds and fish. Wildlife feeding on treated bait may be killed. Do not contaminate water by cleaning equipment or disposal of wastes. Do not feed to livestock or poultry. Do not mix with grain for livestock or poultry feed.

USE AND DISPOSAL

1. PESTICIDE DISPOSAL
Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to federal or approved state procedures. Contact your local or state Department of Agriculture for disposal instructions.

2. NEVER REUSE
Complete container by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into applicator. Wash container thoroughly of bags by a suitable method or by incineration if allowed by state and local authorities.

3. STORAGE

Do not store in violation of local, state or federal law to use. Do not store in a manner inconsistent with its labeling.

4. FEEDING

Do not use where it might be fed to man or animal. Do not use in any manner that might be injurious to man and protected animals. Do not use in proximity of water, ponds, streams, or other bodies of water. Do not use in areas where it might be fed to man or animal.

Do not use in areas where it is prohibited by local, state or federal law.

RESTRICTED USE PESTICIDE

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

AVITROL

EC Corn Chops 99%

Active Ingredient: 4-Aminopyridine 0.012
Inert Ingredients: 99.988
Total 100.000

Avitrol is a poison with flock alarming properties used for the control of Red-Winged, and Yellow-Headed Blackbirds, Cowbirds, Common Grackles, and Starlings in Ripening Sunflower Fields in such a way that a part of a flock may react and frighten the rest away. Birds that react and alarm a flock usually die.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed: If the patient is unconscious, maintain breathing and heart beat (CPR, Cardiopulmonary resuscitation). Contact your local Poison Control Center, hospital or physician immediately.

If patient is conscious: Induce vomiting with Syrup of Ipecac (if not available), stimulate the back of the throat with fingers. Never give anything by mouth to an unconscious person. Contact your Poison Control Center, hospital or physician immediately.

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

See side panel for additional precautionary statements.

Manufactured by

AVITROL
Corporation
1205 S. Boston
Tulsa, Oklahoma 74104

EPA Reg. No. 11649-15

EPA USE
See instructions on label


Pretreatment Considerations

Avitrol EC Corn Chops-99% is composed of medium chopped corn coated with an active bird management chemical. Birds ingesting Avitrol react with distress symptoms and calls. By limiting the amount of bait available to relatively few birds, the remainder of the flock can be frightened away from feeding sites with a minimum of mortality. When bird damage reaches 200 treated damaged heads, or when at least 200 feeding target birds are observed in a field, baiting procedures may be discontinued.

Application Directions

This product must be applied in accordance with 40 CFR Part 170. Confine treatment to areas 50 feet in from field edge. Do not apply more than five applications per crop season. No application rate in excess of 0.014 ounces of 4-aminopyridine per acre on the portion being treated may be used. If properly calibrated equipment, apply the bait evenly by air or by ground in swaths 10 to 15 rows wide and 30 to 35 rows between swaths. The amount for one treatment is one pound per acre based on the total area (applied at 1 pound per acre on treated portion). Repeat at 10 to 15 day intervals (5 inches) or when the material becomes depleted. Treatment areas or swaths should be alternated so that the total material applied in a season is evenly distributed over the field. It is important that Avitrol EC Corn Chops 99% be widely scattered and the recommended amount per acre be strictly observed. The single treated acre (1 in 100) are most toxic to blackbirds, but are thinly distributed by the birds with clean grain that very few birds are exposed. Applied in this way it is not likely that enough treated grain can be found to constitute a hazard to large waterfowl, such as geese. After set back, additional applications should be taken in baiting with Avitrol, since migratory grain-eating birds may be in the area in increased numbers.

Notice: Buyer and/or user assumes all risks of use and handling of these materials in accordance with the restrictions.



BEST AVAILABLE COPY

11649-15

Reg no: 11649-15

PM-16

181

NOT REVIEWED

In Accordance with FR Notice 82-77
Based on Draft Labeling Dated 6/11/82

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. May cause irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds and fish. Wildlife feeding on treated bait may be killed. Do not contaminate water by cleaning equipment or disposal of waste. Do not feed to livestock or poultry. Do not mix with grain for livestock or poultry feed.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

Do not use where food (grain or meat) might become contaminated.

Avitrol must not be exposed in any manner that may endanger desirable and protected bird species. If there is a question of such hazard, consult local state and federal game authorities before undertaking bird management with Avitrol.

Investigate local laws that may prohibit the use of any toxic chemical in bird control.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

1. STORAGE INSTRUCTIONS

Store in original container only. Store apart from food or animal feed and protect against rodent penetration of the carton. Store apart from other pesticides, fertilizers, food or feed that may cause cross-contamination of odor or taste.

2. DISPOSAL INSTRUCTIONS

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mix, or empty container is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill, if incineration is allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

RESTRICTED USE PESTICIDE

For sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.



FC CORN CHOPS—99S

Active Ingredient: 4-Aminopyridine	0.03%
Inert Ingredients	99.97%
Total	100.0%

Avitrol is a poison with flock alarming properties used for the control of Red Winged and Yellow Headed Blackbirds, Cowbirds, Common Grackles and Starlings in Ripening Sunflower Fields in such a way that a part of a flock may react and frighten the rest away. Birds that react and alarm a flock usually die.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed, the patient is unconscious, maintain breathing and heart beat. Call Poison Control Center, hospital, or physician immediately.

If patient is conscious, induce vomiting with syrup of Ipecac. If not available, stimulate the back of the throat with finger. Never give anything by mouth to an unconscious person. Contact your Poison Control Center, hospital, or physician immediately.

If in eyes, flush with plenty of water. Get medical attention if irritation persists.

See side panel for additional precautionary statements.

Manufactured By
AVITROL CORPORATION

Tulsa, Oklahoma 74145
EPA Reg. No. 11649-15
EPA Est. No. 74-1001

EPA No.

Avitrol (4-Aminopyridine)

PRETREATMENT CONSIDERATIONS

Avitrol FC Corn Chops-99S is composed of medium chopped corn coated with an active bird management chemical. Birds ingesting Avitrol react with distress symptoms and calls. By limiting the amount of bait available to relatively few birds, the remainder of the flock can be frightened away from feeding sites with a minimum of mortality. When bird damage reaches 200 freshly damaged heads, or when at least 200 feeding target birds are observed in a field, baiting procedures may be initiated.

APPLICATION DIRECTIONS

This product must be applied in accordance with 40 CFR Part 170. Confine treatment to areas 50 feet in from field edge. Do not apply more than five applications per crop season. No application rate in excess of 0.144 ounces of 4-aminopyridine per acre on the portion being treated may be used. Using calibrated equipment, apply the bait evenly by air or by ground in swaths 10 to 15 rows wide and 30 to 35 rows between swaths. The amount for one treatment is one pound per acre based on the total area (applied at 3 pounds per acre on treated portion). Repeat after a heavy rain (5 inches or more) when the material becomes depleted. Treatment areas or swaths should be alternated so that the total material applied in a season is evenly distributed over the field. It is important that Avitrol FC Corn Chops 99S be widely scattered and the recommended amount per acre be strictly observed. The single treated kernels, 1 in 100 are toxic to blackbirds, but are so thinly distributed by mixing with clean grain that very few birds are exposed. Applied in this way it is noted that enough treated grain can be found to constitute a hazard to large game birds such as pheasants. After October 1, additional care should be taken in baiting with Avitrol since migratory grain-eating birds may be in the area in increased numbers.

Notice: Buyer and/or user assumes all risks of use and of injury from these materials contrary to label instructions.