


For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



## .2% GRAIN BAIT FOR POCKET G

CONTAINS \*GOPHACIDE FOR HAND OR BURROW MACHINE BAITING

Active Ingredient: O,O-bis(p-chlorophenyl) acetimidoylphosphoroamidothicate 0.2% Inert Ingredients ..... 99.8% 100.0%

### WARNING KEEP OUT OF REACH OF CHILDREN

Additional CAUTION, WARNING, RESTRICTION and ANTIDOTE STATEMENTS on right panel.

> NET WEIGHT POUNDS

Available in 1#, 3#, 5#, 10#, 25# and 504 bags.

GOPHACIDE is Reg. TM in U.S. Pat. Off. by Farbenfabriken Bayer A.G., Chemagro Corporation licensee.

EPA Reg. No. 11656-67

### STOP — READ THIS LABEL BEFORE USE WARNING

May be harmful or fatal if swallowed or absorbed through the skin. Do not get in eyes or on skin. This material Repeated contact with the skin may increase hazard of handling. Wear cotton or natural rubber gloves whi cotton gloves are used, they must be laundered or discarded after each day's use. Always wash thoroughly warm water after handling. Wash contaminated clothing with soap and water before reuse.

Do not store or use in an area where there is possibility of children or animals gaining access to the mat carcasses of animals poisoned by the bait.

If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression induce vomiting by giving a tablespoonful of salt in a glass of warm water. Administer milk or water freely and vomiting until ve mit fluid is clear. If eyes are contaminated, wash with flowing water for at least 15 minutes.

FISH AND WILDLIFE: This product is toxic to fish and wildlife. Keep out of any body of water. Treated bait  $\epsilon$ surface will be hazardous to birds and other wildlife. Apply this product only as specified on this label. ANTIDOTE: Although atropine may be less effective as an antidote with this than with some other organic ph

pounds, it should be used in large therapeutic doses in cases of acute poisoning. Repeal as necessary to the pois RESTRICTIONS: For use only by personnel experienced in pocket gopher control. Consult your local, co rodent control experts for advice on how to best use this product safely and efficiently. Apply only in burro above ground surface. Control of other rodent species is not recommended, since evaluations have not yet be

### DIRECTIONS FOR USE

.2% Grain Bait for Pocket Gophers is very effective for the control of the valley (Thomomys bottae) pocket gopher. Bait preparations ar and death usually occurs within 24 to 48 hours after the bast is eaten. The mountain (Thomomys talpoides) and plains (Geomys bursariu controlled by this concentration. HOW to APPLY: Place in artificial burrows made with suitable burrow building equipment. Use 112 Bast for Pocket Gophers per acre in burrows spaced at 20 foot intervals or 1 lb. per acre in burrows spaced at 30 foot intervals. Treatment to natural burrows using 1 teaspoon of .2% Grain Bast for Pocket Gophers per burrow, but not to exceed 1½ per acre of area to be treat

DO NOT REUSE THIS CONTAINER -- Completely empty the contents and bury the unused chemical at least 18 inches deep in an isol from water supplies.

Burn the empty container completely. Exercise extreme caution and stay well away from the smoke

Do not use the container in any connection with feed, food, or drinking water

– PLANT – Second and M Streets Telephone 353-2660 IMPERIAL, CALIFORNIA

WF5 67 ( - 1 60

**WESTERN FARM SERVICE, INC.** 


**EPA** Est. No. 11(

3075 CITRUS CIRCLE **WALNUT CREEK, CA 94598** 



### RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



## .2% GRAIN BAIT FOR POCKET GOPHERS

CONTAINS ® GOPHACIDE FOR HAND OR BURROW MACHINE BAITING

OF REACH OF

TION, WARNING, RES-

NTIDOTE STATEMENTS

IILDREN

# STOP — READ THIS LABEL BEFORE USE WARNING

May be harmful or fatal if swallowed or absorbed through the skin. Do not get in eyes or on skin. This material is cumulative. Repeated contact with the skin may increase hazard of handling. Wear cotton or natural rubber gloves while handling. If cotton gloves are used, they must be laundered or discarded after each day's use. Always wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and water before reuse.

Do not store or use in an area where there is possibility of children or animals gaining access to the material or to the carcasses of animals poisoned by the bait.

If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression. If swallowed, induce vomiting by giving a tablespoonful of salt in a glass of warm water. Administer milk or water freely and again induce vomiting until vomit fluid is clear. If eyes are contaminated, wash with flowing water for at least 15 minutes.

FISH AND WILDLIFE: This product is toxic to fish and wildlife. Keep out of any body of water. Treated bait exposed on soil surface will be hazardous to birds and other wildlife. Apply this product only as specified on this label.

ANTIDOTE: Although atropine may be less effective as an antidote with this than with some other organic phosphorus compounds, it should be used in large therapeutic doses in cases of acute poisoning. Repeat as necessary to the point of tolerance.

RESTRICTIONS: For use only by personnel experienced in pocket gopher control. Consult your local, county, or state rodent control experts for advice on how to best use this product safely and efficiently. Apply only in burrows. Do not use above ground surface. Control of other rodent species is not recommended, since evaluations have not yet been completed.

\_\_ POUNDS

14, 34, 54, 104, 254 and ' )4 bags.

is Reg. TM in U.S. Pat. Off. in Bayer A.G., Chemagro isee.

1656-6 7

#### DIRECTIONS FOR USE

2% Grain Bait for Pocket Gophers is very effective for the control of the valley (Thomomys bottae) pocket gopher. Bait preparations are readily accepted and death usually occurs within 24 to 48 hours after the bait is eaten. The mountain (Thomomys talpoides) and plains (Geomys bursarius) gophers are also controlled by this concentration. HOW to APPLY: Place in artificial burrows made with suitable burrow building equipment. Use 1½ lbs. of 2½ Grain Bait for Pocket Gophers per acre in burrows spaced at 20 foot intervals or 1 lb. per acre in burrows spaced at 30 foot intervals. Treatment may also be made to natural burrows using 1 teaspoon of .2% Grain Bait for Pocket Gophers per burrow, but not to exceed 1½ per acre of area to be treated.

DO NOT REUSE THIS CONTAINER — Completely empty the contents and bury the unused chemical at least 18 inches deep in an isolated location away from water supplies.

Burn the empty container completely. Exercise extreme caution and stay well away from the smoke

Do not use the container in any connection with feed, food, or drinking water

## WESTERN FARM SERVICE, INC.

EPA Est. No. 11656-CA-27

3075 CITRUS CIRCLE WALNUT CREEK, CA 94598

-7 GL LBO

LANT -

end M Streets one 353-2660 CALIFORNIA