

# FIRST CHOICE

## DUSTING SULFUR

### An Agricultural Insecticide and Fungicide

ACTIVE INGREDIENT	BY WEIGHT
Sulfur .....	98%
INERT INGREDIENTS .....	2%
EPA Reg. No. 11656-3-ZA	EPA Est. No. 2935-CA-1
	TOTAL 100%

ACCEPTED

9-30-97

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 11656-3

**READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT**

### KEEP OUT OF REACH OF CHILDREN CAUTION

FOR AGRICULTURAL USE ONLY OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN. SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

#### STATEMENT OF PRACTICAL TREATMENT

If an adverse reaction occurs after exposure: Promptly start the recommended procedures below. Seek medical attention if further treatment is required or call (800) 424-9300.

If swallowed: Give several glasses of water, induce vomiting by gagging the victim with a finger placed on the back of the victim's tongue. Give fluids until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If inhaled: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

In case of contact: Flush all affected areas with plenty of water for several minutes.

For eye contact: Hold eyelids apart and flush with plenty of water for 15 minutes. Get medical attention if irritation occurs.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if inhaled or absorbed through skin. Avoid breathing dust. Do not take internally. Avoid contact with skin, eyes and clothing.

#### PERSONAL PROTECTIVE EQUIPMENT:

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Goggles or face shield

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATION

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if restrictive materials are worn. Then wash thoroughly and clean clothing.

#### DOSEAGE RECOMMENDATION

##### FRUIT CROPS

CITRUS: Six-spotted mite, citrus rust mite, citrus flat mite, Yuma spider mite, citrus thrips- Apply 80-100 lbs. per acre. Follow recommendations of State Agricultural Experiment Stations.

CAUTION: Avoid applications during or preceding high temperatures. Do not apply within 2 months of an oil spray.

GRAPES: Powdery mildew-Apply 8-20 lbs. per acre. Make first application when shoots are 6 in. to 8 in. long. Make a second application when shoots are 12 in. to 15 in. long and the third about 14 days later. Repeat applications at 10-14 days intervals to protect new growth.

CAUTION: Avoid over application in hot water. Farm workers are not permitted to enter the treated area to engage in activity requiring substantial contact with treated foliage within one (1) day after application.

APPLES: Scab, powdery mildew- Apply 30-35 lbs. per acre in the green tip to pink bud stage.

CAUTION: Use on sulfur tolerant varieties only.

CANEBERRIES (BOYSENBERRIES): Powdery mildew- Apply 15-20 lbs. per acre.

Dust when disease appears and re-treat as necessary. Check for plant safety before treating commercially.

STRAWBERRIES: Powdery mildew- Apply 10-20 lbs. per acre. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commercially.

FIGS: Fig rust mite- Apply 40 lbs. per acre when mites first appear. Since the effectiveness of sulfur varies in different localities, State Agricultural Experiment Stations should be consulted as to the effectiveness before application.

CHERRIES: Brown rot blossom and twig blight and brown rot of fruit, powdery mildew- Apply 40-45 lbs. per acre. Make first application at pink bud stage, repeat at petal fall and shuck fall stages. Repeat as necessary through growing season.

NECTARINES AND PEACHES: Powdery mildew, brown rot blossom and twig blight and brown rot of fruit, rust and peach silver mite- Apply 35-60 lbs. per acre. Dust is necessary during bloom and when rain threatens during growing season up to harvest for brown rot control. Apply when mildew and mites first appear. Repeat as necessary.

CAUTION: Farm workers are not permitted to enter the treated area to engage in activity requiring substantial contact with treated foliage within one (1) day after application.

PLUM AND PRUNES: Brown rot of fruit and prune rust- Apply 35-50 lbs. per acre starting 5 weeks before harvest and thereafter at weekly intervals if wet weather occurs.

##### VEGETABLE AND FIELD CROPS

ALFALFA (Seed): Spider mites- Apply 15-50 lbs. per acre. Apply when spider mites first appear & repeat at 14 day intervals or as necessary.

CAUTION: Sulfur may cause petal burn to open flowers.

BEANS: Spider mites, powdery mildew- Apply 15-30 lbs. per acre for early season mite control when populations first appear.

CAUTION: May cause injury to some varieties of beans. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commercially.

CARROTS, ONIONS, GARLIC: Potato mite- Apply 20-35 lbs. per acre during warm weather.

COTTON: Atlantic mite, two-spotted mite- Apply 25-40 lbs. per acre when mites first appear & repeat as necessary.

CLOVER: Powdery mildew, spider mite- Apply 15-25 lbs. per acre. Dust when disease appears and re-treat

2 of 3

PM21

11656-3

Exact Copy Available

9-30-97

1043

3063

**APPLICATIONS AND OTHER HANDLES MUST WEAR:**

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Goggles or face shield

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATION**

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS:**

Do not apply directly to water, to areas where surface water is present or to riparian areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on label. Do not apply when weather conditions favor drift from treated areas.

**PHYSICAL OR CHEMICAL HAZARDS:**

Barring sulfur emits highly toxic fumes. Sulfur dust suspended in air ignites easily. Keep away from heat, sparks and flame. Avoid creating dust in handling. Keep container closed. Use with adequate ventilation. Keep from contact with oxidizing materials.

In Case of Fire, Use water spray, fog, CO<sub>2</sub>, or sand. Wear NIOSH approved self-contained breathing apparatus.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations. Use the higher dosage during periods of frequent and heavy rainfall. Do not apply when temperatures exceed or are likely to exceed 90°F. Do not use on Eastern American type grapes, apricots, apple and conifer pears, spinach, fibroids, walnuts or other sulfur sensitive plants. Do not use on sulfur sensitive cucurbits. Do not use sulfur within 30 days of an oil spray (60 days on olives). If the crop to be treated will be used for canning or freezing purposes, consult food processors before applying sulfur.

**GENERAL USE DIRECTIONS**

Apply this product uniformly to the foliage when air is calm, using ground or aircraft. Timing and dosage may vary as a result of local weather conditions. Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information concerning timing, number and rate of applications needed, and safe re-entry interval.

**CAUTIONS IN USING**

Some fruit and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Sulfur may burn foliage and fruit during periods of high temperature and under certain climatic conditions. Except on cotton, do not apply when temperatures exceed or are likely to exceed 90°F. Do not use on Eastern American type grapes, apricots, apple and conifer pears, spinach, fibroids, walnuts or other sulfur sensitive plants. Do not use on sulfur sensitive cucurbits. Do not use sulfur within 30 days of an oil spray (60 days on olives). If the crop to be treated will be used for canning or freezing purposes, consult food processors before applying sulfur.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: Coveralls, Waterproof gloves, shoes plus socks, goggles or face shield.

harvest and thereafter at weekly intervals if wet weather occurs.

**VEGETABLE AND FIELD CROPS**

**ALFALFA (Seed):** Spider mite- Apply 18-20 lbs per acre. Apply when spider mites first appear & repeat at 14 day intervals as necessary.

**CAUTION:** Sulfur may cause petal burn to open flowers.

**BEANS:** Spider mite, powdery mildew- Apply 15-30 lbs per acre for early season mite control when populations first appear.

**CAUTION:** May cause injury to some varieties of beans. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commercially.

**CARROTS, ONIONS, GARLIC:** Petal mite- Apply 20-35 lbs per acre during warm weather.

**COTTON:** Atlantic mite, two-spotted mite- Apply 25-40 lbs per acre when mites first appear & repeat as necessary.

**CLOVER:** Powdery mildew, spider mite- Apply 15-25 lbs per acre. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commercially.

**CUCUMBERS, SQUASH, MELONS (Sulfur resistant varieties only):** Powdery mildew- Apply 15-25 lbs per acre. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commercially.

**PEAS:** Powdery mildew- Apply 15-20 lbs per acre. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commercially.

**SUGAR BEETS:** Spider mite- Apply 30-35 lbs per acre. Powdery Mildew: For use in California only. Apply 10-20 lbs per acre when first spots of powdery mildew appear on leaves. Apply second application 3 to 4 weeks after the initial application only if the disease is not adequately controlled by the initial application. Use the higher rates when moderate to severe disease pressure exists. Application made later than 5 weeks before harvest may not increase yield of sugar.

**TOMATOES:** Resat mite, two-spotted mite- Apply 15-50 lbs per acre when mites first appear & repeat as necessary.

**TURNIPS AND RUTABAGAS:** Powdery mildew- Apply 15-20 lbs per acre. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commercially.

**ORNAMENTALS**

Make full coverage application on both sides of leaves. Keep new growth protected. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commercially.

**CHRYSANTHEMUM:** Powdery mildew, rust- Do not apply when flowering. Apply 20-40 lbs per acre.

**HOLLYHOCK:** Rust- Apply 20-40 lbs per acre.

**ROSE:** Black spot, powdery mildew, rust- Apply 10-25 lbs per acre.

**CAUTION:** DO NOT use over 15 lbs per acre when temperatures approach 90°F. Soil must be moist.

**STORAGE AND DISPOSAL**

1. Prohibitions: Do not contaminate water, food or feed by storage or disposal.
2. Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at any approved waste disposal facility.
3. Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.
4. General: Store in original container only and keep in closed storage area.

**CONDITIONS OF SALE AND WARRANTY**

WESTERN FARM SERVICE MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE EXPRESSED OR IMPLIED concerning this product or its uses which exceed beyond the use of the product under normal conditions in accord with the statements made on this label.

**NET WEIGHT: 50 LBS. (22.68 kgs.)**

Manufactured for:  
**WESTERN FARM SERVICE, INC.**  
 P.O. BOX 1168  
 FRESNO, CA 93715

FIRST CHOICE® is a registered trademark of Western Farm Service, Inc.

Handwritten initials/signature.

30/3

1 of 3

# FIRST CHOICE DUSTING SULFUR

## An Agricultural Insecticide and Fungicide

ACTIVE INGREDIENT	BY WEIGHT
Sulfur .....	98%
INERT INGREDIENTS .....	2%
EPA Reg. No. 11056-3-ZA	EPA Est. No. 2935-CA-1
	TOTAL 100%

### READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT

#### KEEP OUT OF REACH OF CHILDREN CAUTION

FOR AGRICULTURAL USE ONLY OR FOR BULK TO USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN. SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

**STATEMENT OF PRACTICE/TREATMENT**

If an adverse reaction occurs after application: Promptly start the recommended procedures below. Seek medical attention if further treatment is required or call (703) 434-9700.

If symptoms: Give several glasses of water, induce vomiting by passing the victim with a finger placed on the back of the victim's tongue. Give fluids until vomiting is done. Do not induce vomiting or give anything by mouth to an unconscious or convulsing patient.

If inhaled: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

In case of contact: Flush all affected areas with plenty of water for several minutes.

For eye contact: Hold eyelids apart and flush with plenty of water for 15 minutes. Get medical attention if irritation occurs.

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Irritant if inhaled or absorbed through skin. Avoid breathing dust. Do not inhale directly. Avoid contact with skin, eyes and clothing.

#### PERSONAL PROTECTIVE EQUIPMENT:

- Application and other handlers must wear:
- Long-sleeved shirt and long pants
  - Waterproof gloves
  - Shoes with socks
  - Goggles or face shield
- Follow label/industry's instructions for clean/cleanability PPE. If no such instructions for washability, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATION

- User should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
  - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and get out on clean clothing.
  - Remove PPE immediately after handling the product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS:

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on label. Do not apply when weather conditions favor drift from treated areas.

**PHYSICAL OR CHEMICAL HAZARDS:**

Burning sulfur fumes highly irritant to eyes. Sulfur dust irritates in or irritates easily. Keep away from heat, sparks and fire. Avoid spraying dust in handling. Keep container closed. Use with adequate ventilation. Keep from contact with oxidizing materials.

In Case of Fire: Use water spray, fog, CO<sub>2</sub>, or sand. Wear AFFF approved self-contained breathing apparatus.

#### DIRECTIONS FOR USE

This is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or other, consult the agency responsible for pesticide registration. Use the higher dosage during periods of frequent and heavy rainfall. Dust often enough to keep susceptible growth covered. This may require weekly or bi-weekly application during periods of rapid growth and frequent rainfall.

#### GENERAL USE DIRECTIONS

Apply this product uniformly to the foliage where it is used, using ground or aerial. Timing and dosage may vary as a result of local weather conditions. Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information concerning timing, number and rate of applications needed, and with necessary interval.

#### CAUTIONS IN USE

Some fruit and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Sulfur may burn foliage and fruit, causing periods of high temperature and moist, humid climatic conditions. Sulfur is not to be used on plants unless the temperature is above 50°F. Do not use on Southern American type peaches, apricots, nectarines and certain peach varieties, blueberries, and other southern plants. Do not use on sulfur sensitive varieties. Do not use sulfur within 30 days of an all spray (30 days or more) if the crop to be treated will be used for drinking or feeding purposes, consult feed processor before applying sulfur.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. The Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural products. It contains requirements for testing, information, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the requirements on the label about personal protective equipment (PPE), notification to workers and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow another entity into treated areas during the restricted entry interval (REI) of 4 hours. PPE required for early entry, to limited areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: Chemical-resistant gloves, shoes and socks, goggles or face shield.

#### DOSEAGE RECOMMENDATION

**FRUIT CROPS**

**CITRUS:** Use against scale, citrus red scale, citrus false scale, Yucca scale, citrus white scale, citrus thrips. Apply 20-100 lbs. per acre. Follow recommendations of State Agricultural Experiment Stations.

**CAUTION:** Avoid applications during or preceding high temperatures. Do not apply within 2 weeks of an oil spray.

**GRAPE:** Powdery mildew Apply 5-20 lbs. per acre. Make first application when shoots are 6 in. to 8 in. long. Make a second application when shoots are 12 in. to 15 in. long and the third about 14 days later. Repeat applications at 10-14 day intervals if present leaf growth.

**CAUTION:** Avoid over application to hot water. Some workers are not permitted to enter the treated area to engage in activity requiring substantial contact with treated foliage within one (1) day after application.

**APPLIED:** Sulfur, powdery mildew Apply 20-30 lbs. per acre to the ground to kill leaf scale.

**CAUTION:** Use an sulfur tolerant varieties only.

**CUCUMBERS (BUCKLELEAF CURCUBIT):** Powdery mildew Apply 15-20 lbs. per acre. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commencing.

**STRAWBERRIES:** Powdery mildew Apply 10-20 lbs. per acre. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commencing.

**PEAS:** Do not apply 40 lbs. per acre which will cause leaf drop. Check the effectiveness of sulfur before in stored locations. State Agricultural Experiment Stations should be contacted as to the effectiveness before application.

**CHERRIES:** Spray hot blossoms and long light and brown rot of fruit, powdery mildew Apply 40-60 lbs. per acre. Make first application of pink bud stage, repeat at post bud and again at stages. Repeat as necessary through growing season.

**NECTARINES AND PEACHES:** Powdery mildew Spray hot blossoms and long light and brown rot of fruit, use one peach tree only. Apply 25-60 lbs. per acre. Dust in necessary during bloom and when rain threatens during growing season up to harvest for brown rot control. Apply when sulfur and sulfur leaf appear. Repeat as necessary.

**CAUTION:** Some varieties are not permitted to enter the treated area to engage in activity requiring substantial contact with treated foliage within one (1) day after application.

**PUMPS AND PUMPKINS:** Spray hot of fruit and make root. Apply 25-30 lbs. per acre starting 5 weeks before harvest and thereafter at weekly intervals if wet weather occurs.

**WATERMELON AND FIELD CROPS**

**MELONS (CUCURBITACEAE):** Powdery mildew Apply 15-20 lbs. per acre. Apply when sulfur mildew first appear & repeat if 14 day intervals or as necessary.

**CAUTION:** Sulfur may cause scald lower to some flowers.

**BANANA:** Substrate, powdery mildew Apply 15-20 lbs. per acre for early season white spread when populations first appear.

**CAUTION:** Sulfur may injury to some varieties of banana. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commencing.

**CARROTS, CHINESE CABBAGE:** Powdery mildew Apply 20-30 lbs. per acre during warm weather.

**COTTON:** Apply 20-30 lbs. per acre when sulfur first appear & repeat as necessary.

**CLOVER:** Powdery mildew Sulfur mildew Apply 15-20 lbs. per acre. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commencing.

**CUCUMBERS, SQUASH, MELONS (CUCURBITACEAE):** Powdery mildew Apply 15-20 lbs. per acre. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commencing.

**GRAPES:** Powdery mildew Apply 15-20 lbs. per acre. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commencing.

**SUGAR BEETS:** Sulfur mildew Apply 20-30 lbs. per acre. Powdery mildew For use in California only. Apply 10-20 lbs. per acre when first signs of powdery mildew appear on leaves. Apply second application 7 to 14 days after the initial application only if the disease is not adequately controlled by the initial application. Use the higher rate when favorable to severe disease pressure exists. Application made later than 2 weeks before harvest may not increase yields of sugar.

**TOMATOES:** Repeat sulfur late season sulfur Apply 15-20 lbs. per acre when sulfur first appear & repeat as necessary.

**TURNIPS AND RUBENGLAS:** Powdery mildew Apply 15-20 lbs. per acre. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commencing.

**ORNAMENTALS**

Make full coverage application on both sides of leaves. Keep new growth protected. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commencing.

**ORNAMENTALS:** Powdery mildew, rust. Do not apply when flowering. Apply 20-40 lbs. per acre.

**HOLLANDS:** Repeat Apply 20-40 lbs. per acre.

**ROSE:** Repeat sulfur, powdery mildew, rust Apply 10-20 lbs. per acre.

**CAUTION:** DO NOT use over 15 lbs. per acre when temperatures approach 90°F. Soil must be moist.

#### STORAGE AND DISPOSAL

1. Pesticides: Do not contaminate water, food or feed by storage or disposal.
2. Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.
3. Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary manner or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.
4. Recycle: Store in original container only and keep in stockpiling areas.

#### CONDITIONS OF SALE AND WARRANTY

WESTERN FARM SERVICE MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED concerning this product or its use when used beyond the use of the product under normal conditions in accord with the statements made on the label.

### NET WEIGHT: 50 LBS. (22.68 kgs.)

Manufactured for:  
**WESTERN FARM SERVICE, INC.**  
 P.O. BOX 1188  
 FRESNO, CA 93715

FIRST CHOICE® is a registered trademark of Western Farm Service, Inc.