

Jan. 16, 1976
1142-8

with comments

PRECAUTIONS IN USING

DANGER: KEEP CHILDREN AND ANIMALS OFF TREATED AREAS. CONCENTRATE CAUSES EYE AND SKIN DAMAGE. Do not get on skin, eyes or clothing. Use waterproof gloves and face shield or goggles when handling concentrate. May be fatal if swallowed or absorbed through skin. Avoid breathing spray mist. In case of contact, immediately remove contaminated clothing and flush skin or eyes with water; for eyes, get medical attention.

Avoid contamination of feed, foodstuffs and drinking water. Avoid storage near feed and food products. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Do not store diluted spray. **Make new dilution for each use. Wash equipment thoroughly after use.**

DO NOT SPRAY ON NECTARINES, ORNAMENTAL CHERRIES, LIQUIDAMBAR OR CHRYSANTHEMUMS AS INJURY MAY OCCUR.

This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crab may be killed at application rates recommended on this label. Do not use where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. When container is empty, immediately destroy. Containers should be disposed of by breaking and burying with wastes. Never re-use containers.

GENERAL DIRECTIONS

Use designated amount of product in the following gallons water per acre unless otherwise noted:

GROUND APPLICATION: Vegetable and Field Crops — 30 to 250 gals.; Cotton — 3 to 20 gals.; Deciduous Fruit Crops — 60 to 800 gals.; Citrus — 400 to 3000 gals. (On Young Fruit Plantings — Use a minimum of 15 gals. water per acre).

Where Recommended For **AIR APPLICATION:** Field Crops 3 to 10 gals.

Begin application at first sign of insects. Repeat as necessary. Do not apply within 4 days of harvest unless otherwise noted. Do not use with highly alkaline materials such as lime or bordeaux.

MIXING DIRECTIONS: Add to spray tank when ¼ to ½ full with agitator running and maintain agitation until spraying is completed.

USE AND APPLICATION DIRECTIONS

BEANS (Dry or Field); SOYBEANS, LIMA BEANS AND PEAS (For Processing): Alfalfa Looper, Aphids, Spider Mites, Lygus, Leaf Miner; **GREEN BEANS:** Aphids, Spider Mites, Loopers, Nitidulids, Leafhoppers — 1 pt. per acre in water to cover; 100 gals. per acre is preferable. Large Loopers, Stink Bugs — 1½ pt. per acre in water to cover.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KALE, COLLARDS: Imported Cabbageworm, Diamond-back Caterpillars, Aphids — 1 pt. per acre. Looper — 2 pts. per acre. May be used one day before harvest, except do not apply to kale and collards within four days of harvest.

CELERY: Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners — Use ground application only with fine mist spray — 1 pt. in 50 to 75 gals. water per acre on plants less than ¾ grown. On larger plants use 1½ pts. in 100 to 150 ga s. water per acre.

COTTON: Lygus — ½ to 1 pt. per acre by air or ground. Leaf Perforator, Fleahoppers, Armyworms, Bollworms, Loopers and for Suppression of Mites — 1 pt. per acre by air or ground. Apply at 7 to 10-day intervals while pests persist. For Heavy Populations of Listed Insects — Use ½ pt. per acre by air or ground with other insecticides registered for use on cotton.

CUCUMBERS, SUMMER SQUASH: Cucumber Beetles, Aphids, Spider Mites, Loopers, Thrips, Armyworms — 1 pt. per acre in water to cover. Pickleworms, Leaf Miners — 1½ pts. per acre in water to cover. By Air — Use minimum of 10 gals. water per acre. May be used up to harvest.

EGGPLANT, PEPPERS: Aphids, Spider Mites, Blister Beetles, Flea Beetles, Leaf Miners, Whitefly — 1 pt. per acre. May be used up to 1 day before harvest.

WALNUTS: Aphids — ½ to 1 pt. to 100 gals water. Use up to ten days of harvest.

GRAPES: Drosophila, Leafhoppers — ½ to ¾ pt. per 100 gals. water (100 to 130 gals. diluted spray per acre depending on size of vines). Use only ½ pt. per 100 gals. water on Muscat varieties. Do not use on Italia varieties as fruit injury may occur. Do not enter vineyards for 24 hours following treatment.

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, Cutworm — 1 pt. per acre.

SPINACH, CHARD, TURNIP GREENS: Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners — Use ground application only with fine mist spray — 1 pt. in 50 to 75 gals water per acre on plants less than ¾ grown. On larger plants, use 1½ pts. in 100 to 150 gals. water per acre. May be used 1 day before harvest.

MELONS (Cantaloupes, Honeydew, Muskmelons, Watermelons), PUMPKINS, WINTER SQUASH: Loopers, Armyworms, Leafhoppers, Spider Mites, Leaf Miners — 1 to 2 pts. per acre (200 gals. diluted spray per acre). May be used 1 day before harvest. Do not apply when temperature is over 90° F.



DIBROM® 8

EMULSIVE NALED INSECTICIDE
Contains 8 lbs. Technical Naled Per Gal.
Equivalent to 7.2 lbs. Naled

(FLAMMABLE LIQUID, N.O.S. — FLAMMABLE LIQUID)

ACTIVE INGREDIENTS	By Weight
Naled*	58%
Xylene	20%
INERT INGREDIENTS	22%
U. S. Pat No. 2,971,882	TOTAL 100%
*1,2-dibromo-2,2-dichloroethyl dimethyl phosphate	
EPA Reg. No. 11656-8 AA	NET CONTENTS GALS.

READ THE LABEL

DANGER:

KEEP OUT OF REACH OF CHILDREN

(SEE SIDE PANEL FOR ADDITIONAL DANGER STATEMENTS.)

Do not take internally. Do not get in eyes.

Do not get on skin.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENT, DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers.

STORE IN COOL DRY PLACE.

Not for use or storage in or around the home.

WESTERN FARM SERVICE, INC.

2401 Crow Canyon Road
San Ramon, California 94583

WFS 8-01 8-75

ORANGE
Soft Brown S
(Torrix) — 1
100 gals. wa
Arizona — Do
citrus groves

PEACHES
Mites, Peach
¾ pt. to 100
within 30 days
90° F. Do not

RICE: Gr
sign of insect
used 2 days b

STRAWB
ous Leaf Tier,
used 1 day be

SUGAR B
by air or grou
not apply with

TOBACCO
to 200 gals. di
in water to cov
sary.

TOMATO
less than 40 g
gals. by air. M
and continue t

Field and Gre
Miners, Flea
plants thorough

per acre. Do
used 1 day be

GREENHO
ORNAMENTAL
Mites, Adult W

per 10,000 cu
pipes (or in d
intermittent se

the pipes thro
Application to
heating pipes

at least 1 hour
house for at k
following treat

goggles and a
re-enter. Wear
tilate. Protect

covering with
house is vent
closer than 1

Non-Resistant
Repeat as nec
7-day interval

ACCEPTED WITH COMMENTS



DIBROM[®] 8

EMULSIVE NALED INSECTICIDE

Contains 8 lbs. Technical Naled Per Gal.
Equivalent to 7.2 lbs. Naled

(FLAMMABLE LIQUID, N.O.S. - FLAMMABLE LIQUID)

ACTIVE INGREDIENTS	By Weight
Naled*	58%
Xylene	20%
INERT INGREDIENTS	22%
U. S. Pat No. 2,971,882	TOTAL 100%

*1,2-dibromo-2,2-dichloroethyl dimethyl phosphite
EPA Reg. No. 11656-8 AA NET CONTENTS GALS.

READ THE LABEL

DANGER:

KEEP OUT OF REACH OF CHILDREN

(SEE SIDE PANEL FOR ADDITIONAL DANGER STATEMENTS.)

Do not take internally. Do not get in eyes.

Do not get on skin.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENT, DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers.

STORE IN COOL DRY PLACE.

Not for use or storage in or around the home.

WESTERN FARM SERVICE, INC.

2401 Crow Canyon Road
San Ramon, California 94583

WFS 8-01 8-75

ORANGES, LEMONS, GRAPEFRUIT, TANGERINES: Aphids, Soft Brown Scale Crawlers, Mites, Citrus Cutworms, Leaf Roller (Tortrix) — 1 pt. to 100 gals. water. Thrips (Bloom) — ½ pt. per 100 gals. water. Use up to 7 days of harvest. (California and Arizona — Do not apply within 1 day of harvest). Do not enter citrus groves for 24 hours following treatment.

PEACHES (Western States Only): Lygus, Suppression of Mites, Peach Twig Borer, Oriental Fruit Moth (California Only) — ¾ pt. to 100 gals. water. To avoid fruit spotting, do not use within 30 days of harvest. Do not apply when temperature is over 90° F. Do not enter orchards for 24 hours following treatment.

RICE: Grasshoppers — ½ to ¾ pt. per acre. Apply at first sign of insects. Do not make more than 3 applications. May be used 2 days before harvest.

STRAWBERRIES: Leafroller, Spider Mites, Aphids, Omnivorous Leaf Tier, Spittlebugs (Northwest) — 1 pt. per acre. May be used 1 day before harvest.

SUGAR BEETS: Spider Mites, Leafhoppers — 1 pt. per acre by air or ground. By Air — Use in 1 to 5 gals. water per acre. Do not apply within 5 days of harvest.

TOBACCO: Hornworm — 1 pt. to 100 gals. water, using up to 200 gals. diluted spray per acre. Flea Beetles — 1 pt. per acre in water to cover. Apply at first signs of insects. Repeat as necessary.

TOMATOES (Field): Fruit Flies (Drosophila) — 1 pt. in no less than 40 gals. water per acre with ground sprayer, 7 to 10 gals. by air. Make first application 5 to 7 days before first picking and continue at 5 to 7-day intervals until the day before harvest. Field and Greenhouse: Tomato Fruit Worm, Hornworms, Leaf Miners, Flea Beetles, Mites — 1 pt. in 100 gals. water. Spray plants thoroughly. For Field — Use up to 200 gals. diluted spray per acre. Do not apply when temperature is over 90° F. May be used 1 day before harvest.

GREENHOUSE, VAPOR TREATMENT OF ROSES; OTHER ORNAMENTAL PLANTS, TOMATOES, CUCUMBERS: Spider Mites, Adult Whiteflies, Aphids, Leafrollers, Mealybugs — 1 fl. oz. per 10,000 cu. ft. when plants are dry. Apply undiluted on heat pipes (or in open pans on hot plates). Apply continuously or to intermittent sections of pipe on each side, and on 1 or more of the pipes through center of each range depending upon width. Application to cold pipes should be followed by immediately heating pipes to 160° F. In steam-heated houses, close vents for at least 1 hour after pipes are hot. With hot water systems, close house for at least 3 hours. Houses may remain closed overnight following treatment. When application is made to hot pipes, wear goggles and approved respirator. Ventilate house before workers re-enter. Wear respirator when entering treated houses to ventilate. Protect hives of bees in cucumber and other houses by covering with plastic or rubber sheet during treatment and until house is ventilated. Do not apply to tomatoes or cucumbers closer than 1 day before harvest. Spider Mites (Resistant and Non-Resistant) — Make 3 to 4 treatments at 3 or 4-day intervals. Repeat as necessary. Adult Whiteflies — Make 2 applications at 7-day intervals. Repeat as necessary. Aphids — Apply when

aphids appear. Repeat as necessary. **WARNING:** Avoid overtreatment and direct application to plants as injury may result. White Butterfly and Golden Rapture Roses, Pink Champagne Chrysanthemums, Green Wandering Jew, Poinsettias and Dutchman's Pipe may be injured by DIBROM vapor.

FRUIT FLIES (Drosophila sp.) In and Around Food Processing Plants, Loading Docks, Cull Piles and Refuse Areas, Cider Mills: Use 2 ozs. in 2½ gals. water (5 pts. to 100 gals.) as a coarse spray to walls, floors, doorways, windows, refuse and cull piles where insects congregate. Apply every 5 to 7 days as necessary. Do not apply to cull fruit or refuse piles to be fed to livestock. Avoid contamination of feeds, foodstuffs and food processing machinery. Do not apply when plants are in operation or when foods are present or exposed. Do not spray surfaces which will come into contact with foods. Cover food containers during spraying.

ADULT MOSQUITO, GNAT, STABLEFLY (Dog Flies) AND HOUSEFLY CONTROL — Residential Areas, Crops Listed on Label, Tidal Marshes, Swamps, Woodland, Livestock Pastures, Feed Lots and Pastures Including Dairy Cattle: Consult your State Fish and Game Agency before applying this product. It is not necessary to avoid farm buildings and dairy barns. Make applications during peak of infestation and repeat as necessary. See crop recommendations for use limitations near harvest. This rate of application will kill shrimp. Do not apply to tidal or marsh waters which are important shrimp-producing areas. **AIRCRAFT:** East — Apply 0.8 to 4 fl. ozs. per acre diluted with water. Apply 2 to 8 qts. of diluted spray per acre. West — Apply 1.6 to 4 fl. ozs. per acre diluted with water. Apply 2 to 8 qts. of diluted spray per acre. **MIST OR COLD FOG:** Use 3 to 5 qts. per 100 gals. water. Calibrate equipment (rate of travel and output) to apply 0.1 to 0.25 lb. technical Naled per acre.

For outdoor adult mosquito control, treat shrubbery and vegetation where mosquitoes may rest. Shrubby and vegetation around stagnant pools, marshy areas, ponds and shore lines may be treated. Application of this product to any body of water is prohibited.

CORRALS, ADJACENT PASTURES, HOLDING PENS (Dairy and Beef Cattle, Hogs, Sheep, Horses): Houseflies, Mosquitoes (Airplane Application) — Use 1.6 to 4 fl. ozs. per acre. Dilute 1 to 2½ pts. in 10 gals. water. Apply 1 gal. diluted spray per acre. Apply over areas with animals present.

CLOVER MITES, ROACHES, EARWIGS AND LEAFHOPPERS Outside Dwellings and In Lawns: Use 1 fl. oz. in 3 gals. water. Apply to turf and to soil surfaces around flowers, shrubs and trees for general pest cleanup.

NOTICE OF WARRANTY

WESTERN FARM SERVICE, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

PRECAUTIONS IN USING

DANGER: KEEP CHILDREN AND ANIMALS OFF TREATED AREAS. CONCENTRATE CAUSES EYE AND SKIN DAMAGE. Do not get on skin, eyes or clothing. Use waterproof gloves and face shields or goggles when handling concentrate. May be fatal if swallowed or absorbed through skin. Avoid breathing spray mist. In case of contact immediately remove contaminated clothing and wash skin or eyes with water; for eyes, get medical attention.

Avoid contamination of feed, feedstuffs and drinking water. Avoid storage near feed and food products. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Do not store in a closed spray. **Make new dilution for each use. Wash equipment thoroughly after use.**

DO NOT SPRAY ON NECTARINES, ORNAMENTAL CHERRIES, LIQUIDAMBAR OR CHRYSANTHEMUMS AS INJURY MAY OCCUR.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Spraying and drift may be fatal at application rates recommended on this label. Do not use where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. When containers are empty, immediately destow. Clean area of spillage by wiping up by breaking and burying with wastes. Never reuse containers.

GENERAL DIRECTIONS

Use designated amount of product in the following gallons water per acre unless otherwise noted:

GROUND APPLICATION: Vegetable and Field Crops: 30 to 250 gals.; Cotton: 3 to 20 gals.; Deciduous Fruit Crops: 60 to 800 gals.; Citrus: 400 to 3000 gals. (On Young Fruit Plantings: Use a minimum of 15 gals. water per acre.)

Where Recommended For **AIR APPLICATION:** Field Crops: 3 to 10 gals.

Begin application at first sign of insects. Repeat as necessary. Do not apply within 4 days of harvest unless otherwise noted. Do not use with highly alkaline materials such as lime or bordeaux.

MIXING DIRECTIONS: Add to spray tank when 1/2 to 1 1/2 gal. of water running and maintain agitation until spraying is completed.

USE AND APPLICATION DIRECTIONS

BEANS (Dry or Field); SOYBEANS, LIMA BEANS AND PEAS (For Processing): Alfalfa Looper, Aphids, Spider Mite, Lygus, Leaf Miner, **GREEN BEANS:** Aphids, Spider Mites, Looper, Cutworms, Leafhoppers. 1 pt. per acre in water to cover. 100 gals. per acre is preferable. Large Looper, Stink Bugs: 1 1/2 pt. per acre in water to cover.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KALE, COLLARDS: Imported Cabbageworm, Diamond-back Caterpillars, Aphids: 1 pt. per acre. Looper: 2 pts. per acre. May be used one day before harvest, except do not apply to kale and collards within four days of harvest.

CELERY: Aphids, Looper, Armyworms, Salt Marsh Caterpillars, Leaf Miners. Use ground application only with the mist spray. 1 pt. in 50 to 75 gals. water per acre on plants less than 7 1/2 grown. On larger plants use 1 1/2 pts. in 100 to 150 gals. water per acre.

COTTON: Lygus: 1 1/2 to 1 pt. per acre by air or ground. Leaf Perforator, Fleahoppers, Armyworms, Bollworms, Loopers and for Suppression of Mites: 1 pt. per acre by air or ground. Apply at 7 to 10 day intervals while pests persist. For Heavy Populations of Listed Insects: Use 1 1/2 pt. per acre by air or ground with other insecticides registered for use on cotton.

CUCUMBERS, SUMMER SQUASH: Cucumber Beetles, Aphids, Spider Mites, Looper, Thrips, Armyworms: 1 pt. per acre in water to cover. Pickleworm, Leaf Miners: 1 1/2 pts. per acre in water to cover. By Air: Use minimum of 10 gals. water per acre. May be used up to harvest.

EGGPLANT, PEPPERS: Aphids, Spider Mites, Buster Beetles, Flea Beetles, Leaf Miners, Whitefly: 1 pt. per acre. May be used up to 1 day before harvest.

WALNUTS: Aphids: 1 1/2 to 1 pt. to 100 gals. water. Use up to 10 days before harvest.

GRAPES: Blackberry Leafhoppers: 1 1/2 to 2 1/2 pt. per 100 gals. water (100 to 150 gals. diluted spray per acre depending on size of vines). Use only 1 1/2 pt. per 100 gals. water on Muscat varieties. Do not use on other varieties as fruit injury may occur. Do not enter vineyards for 24 hours following treatment.

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, Cutworm: 1 pt. per acre.

SPINACH, CHARD, TURNIP GREENS: Aphids, Looper, Armyworms, Salt Marsh Caterpillars, Leaf Miners. Use ground application only with fine mist spray. 1 pt. in 50 to 75 gals. water per acre on plants less than 7 1/2 grown. On larger plants use 1 1/2 pts. in 100 to 150 gals. water per acre. May be used 1 day before harvest.

MELONS (Cantaloupes, Honeydew, Muskmelons, Watermelons), PUMPKINS, WINTER SQUASH: Looper, Armyworms, Leafhoppers, Spider Mites, Leaf Miners: 1 1/2 to 2 pts. per acre. 200 gals. diluted spray per acre. May be used 1 day before harvest. Do not apply when temperature is over 90° F.



DIBROM 8

EMULSIVE HALED INSECTICIDE
Contains 8 lbs. Technical Naled Per Gal.
Equivalent to 7.2 lbs. Naled

(FLAMMABLE LIQUID N O S - FLAMMABLE LIQUID)

ACTIVE INGREDIENTS
Naled[®]
Xylene
INERT INGREDIENTS
U.S. Pat. No. 2,971,882
EPA Reg. No. 11655-8 AA

NET CONTENTS

READ THE LABEL

DANGER:

KEEP OUT OF REACH OF CHILDREN
(SEE SIDE PANEL FOR ADDITIONAL DANGER STATE)
Do not take internally. Do not get in eyes.
Do not get on skin.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENT, DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.
Keep pesticide in original container. Do not put material into food or drink containers.
STORE IN COOL, DRY PLACE.
Not for use in or around the home.

WESTERN FARM SERVICE, INC.
2401 Crocker-Carlson Building
San Ramon, California 94583
WFS 8 01 8-75

80-9-8-75



DIBROM 8

EMULSIVE NALCO INSECTICIDE

Contains 8 lbs Technical Naled Per Gal
Equivalent to 7.2 lbs. Naled

(FLAMMABLE LIQUID, N.O.S. FLAMMABLE LIQUID)

ACTIVE INGREDIENTS	By Weight
Naled*	58%
Xylene	20%
INERT INGREDIENTS	22%
U. S. Pat. No. 2,971,882	TOTAL 100%
*1,2-dibromo-2,2-dichloroethyl dimethyl phosphate	

EPA Reg. No. 11656-8 AA NET CONTENTS GALS.

READ THE LABEL

DANGER:

KEEP OUT OF REACH OF CHILDREN

(SEE SIDE PANEL FOR ADDITIONAL DANGER STATEMENTS)

Do not take internally. Do not get in eyes.
Do not get on skin.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENT, DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers.

STORE IN COOL, DRY PLACE

Not for use or storage in or around the home.

WESTERN FARM SERVICE, INC.

2401 C.R.A. Canyon Road
San Ramon, California 94583

WFS 6 01 8 75

ORANGES, LEMONS, GRAPEFRUIT, TANGERINES: Aphids, Soft Brown Scale, Crawlers, Mites, Citrus Cutworms, Leaf Roller (Tortrix), 1 pt. to 100 gals. water; Trips (Blissus), 1/2 pt. per 100 gals. water. Use up to 7 days of harvest (California and Arizona). Do not apply within 1 day of harvest. Do not enter citrus groves for 24 hours following treatment.

PEACHES (Western States Only): Lygus. Suppression of Mites, Peach Twig Borer, Oriental Fruit Moth (California Only), 1/2 pt. to 100 gals. water. To avoid fruit splitting, do not use within 30 days of harvest. Do not apply when temperature is over 90° F. Do not enter orchards for 24 hours following treatment.

RICE: Grasshoppers, 1/2 to 1/4 pt. per acre. Apply at first sign of insects. Do not make more than 3 applications. May be used 2 days before harvest.

STRAWBERRIES: Leafroller, Spider Mites, Aphids, Omnivorous Leaf Tier, Spittlebugs (Northwest), 1 pt. per acre. May be used 1 day before harvest.

SUGAR BEETS: Spider Mites, Leafhoppers, 1 pt. per acre by air or ground. By Air: Use in 1 to 5 gals. water per acre. Do not apply within 5 days of harvest.

TOBACCO: Hornworm, 1 pt. to 100 gals. water, using up to 200 gals. diluted spray per acre; Flea Beetles, 1 pt. per acre in water to cover. Apply at first signs of insects. Repeat as necessary.

TOMATOES (Field): Fruit Flies (Drosophila), 1 pt. in no less than 40 gals. water per acre with ground sprayer; 7 to 10 gals. by air. Make first application 5 to 7 days before first picking and continue at 5 to 7 day intervals until the day before harvest. **GREENHOUSE, TOMATOES, CUCUMBERS:** Spider Mites, Adult Whiteflies, Aphids, Leafroller, Mealybugs, 1 fl. oz. per 10,000 cu. ft. when plants are dry. Apply undiluted on heat pipes for in open pans on hot plates. Apply continuously or to intermittent sections of pipe on each side, and on 1 or more of the pipes through center of each range depending upon width. Application to cold pipes should be followed by immediately heating pipes to 160° F. In steam heated houses, close vents for at least 1 hour after pipes are hot. With hot water systems, close house for at least 3 hours. Houses may remain closed overnight following treatment. When application is made to hot pipes, wear goggles and apply respirator. Ventilate house before workers re-enter. Wear respirator when entering treated houses to ventilate. Protect bees in cucumber and other houses by covering with plastic or rubber sheet during treatment and until house is ventilated. Do not apply to tomatoes or cucumbers closer than 1 day before harvest. Spider Mites (Resistant and Non-Resistant), Make 3 to 4 treatments at 3 or 4 day intervals. Repeat as necessary. Adult Whiteflies, Make 2 applications at 2 day intervals. Repeat as necessary. Aphids, Apply when

aphids appear. Repeat as necessary. **WARNING:** Avoid over-treatment and direct application to plants as injury may result. White Butterfly and Golden Rapture Roses, Pink Champagne Chrysanthemums, Green Wandering Jew, Poinsettias and Dutchman's Pipe may be injured by DIBROM vapor.

FRUIT FLIES (Drosophila sp.) In and Around Food Processing Plants, Loading Docks, Cull Piles and Refuse Areas, Cider Mills: Use 2 ozs. in 2 1/2 gals. water (5 pts. to 100 gals.) as a coarse spray to walkways, driveways, windows, refuse and cull piles where insects congregate. Apply every 5 to 7 days as necessary. Do not apply to cull fruit or refuse piles to be fed to livestock. Avoid contamination of feeds, foodstuffs, and food processing machinery. Do not apply when plants are in operation or when foods are present or exposed. Do not spray surfaces which will come into contact with foods. Cover food containers during spraying.

ADULT MOSQUITO, GNAT, STABLEFLY (Dog Flies) AND HOUSEFLY CONTROL — Residential Areas, Crops Listed on Label, Tidal Marshes, Swamps, Woodland, Livestock Pastures, Feed Lots and Pastures Including Dairy Cattle: Consult your State Fish and Game Agency before applying this product. It is not necessary to avoid farm buildings and dairy barns. Make applications during peak of infestation and repeat as necessary. See crop recommendations for use limitations near harvest. This rate of application will kill shrimp. Do not apply to tidal or marsh waters which are important shrimp producing areas. **AIRCRAFT:** East: Apply 0.8 to 4 fl. ozs. per acre diluted with water. Apply 2 to 8 qts. of diluted spray per acre. West: Apply 1.6 to 4 fl. ozs. per acre diluted with water. Apply 2 to 8 qts. of diluted spray per acre. **MIST OR COLD FOG:** Use 3 to 5 qts. per 100 gals. water. Calibrate equipment (rate of travel and output) to apply 0.1 to 0.25 lb. technical Naled per acre.

For outdoor adult mosquito control, treat shrubbery and vegetation where mosquitoes may rest. Shrubbery and vegetation around stagnant pools, marshy areas, ponds and shore lines may be treated. Application of this product to any body of water is prohibited.

CORRALS, ADJACENT PASTURES, HOLDING PENS (Dairy and Beef Cattle, Hogs, Sheep, Horses): Houseflies, Mosquitoes (Airplane Application): Use 1.6 to 4 fl. ozs. per acre. Dilute 1 to 2 1/2 pts. in 10 gals. water. Apply 1 gal. diluted spray per acre. Apply over areas with animals present.

CLOVER MITES, ROACHES, EARWIGS AND LEAFHOPPERS Outside Dwellings and in Lawns: Use 1 fl. oz. in 3 gals. water. Apply to turf and to soil surfaces around flowers, shrubs and trees for general pest cleanup.

NOTICE OF WARRANTY

WESTERN FARM SERVICE, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product. Its use will not extend beyond the use of the product under normal conditions in accord with the statements made on this label.