

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

DANGER!
Keep Children and Animals Off Treated Areas. Concentrate Causes Eye and Skin Damage.

Do not get on skin, eyes or clothing. Use water-proof gloves and face shield or goggles when handling concentrate. May be fatal if swallowed or absorbed through skin. Avoid breathing spray mist. In case of contact, immediately remove contaminated clothing and flush skin or eyes with water, for eyes get medical attention.

STATEMENT OF PRACTICAL TREATMENT
If Swallowed: Send for physician. Induce vomiting immediately by giving 2 glasses of water and sticking finger down throat.
If On Skin: Immediately remove all contaminated clothing including shoes. Wash all contaminated areas, including hair, with soap and plenty of water. Do not reuse contaminated clothing or shoes until thoroughly cleaned. Send for physician if symptoms of poisoning develop.
If In Eyes: Immediately flush eyes with plenty of water for at least 15 minutes. Send for physician.

ENVIRONMENTAL HAZARDS
This pesticide is toxic to fish, aquatic invertebrates, and wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water or wetlands. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use with highly alkaline materials. Shrimp and crabs may also be killed at application rates recommended. Do not apply to tidal or marsh waters which are important shrimp producing areas. Do not apply when weather conditions favor drift from treated areas.

DIRECTIONS FOR USE
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

REENTRY STATEMENT
Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The areas being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

ORAL WARNINGS shall inform workers of areas or fields that may not be entered, without protective equipment or clothing specified on this label, during treatment or until sprays have dried. Follow first aid treatment instructions shown on this label in case of accidental exposure.

When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

WRITTEN WARNINGS must include the following information.

DANGER-Area treated with Dibrom on (Date) .
Do not enter without appropriate protective clothing until sprays have dried. Follow first aid treatment instructions shown on this product label in case of accidental exposure.

STORAGE AND DISPOSAL

PROHIBITIONS
Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE
Keep pesticide in original container. Do not use, pour, spill or store near heat, or open flame.

Do not store diluted spray. Make new dilution for each use. Wash equipment thoroughly after use.

Do not store or transport near food or feed.

Store in cool dry place.

Not for use or storage in or around the home.

For help with any spill, leak, fire or exposure involving material, call day or night 1-800-424-9100.

PESTICIDE DISPOSAL
Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL
Metal: Triple rinse (or equivalent). Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
Glass: Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved state and local procedures.



DIBROM®

EMULSIVE NALED INSECTICIDE

Contains 8 lbs Technical Naled Per Gal
Equivalent to 7.2 lbs. Naled.

ACTIVE INGREDIENTS	By Wt
Naled.....	58%
Xylene.....	20%
INERT INGREDIENTS	22%
U.S. Pat No 2,971,887	TOTAL 100%
*1,2-dibromo-2,2-dichloroethyl dimethyl phosphate	

KEEP OUT OF REACH OF CHILDREN
DANGER — PELIGRO
FOR AGRICULTURAL USE ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN
PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le expliquen haya sido explicado ampliamente.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NOTICE OF WARRANTY
WESTERN FARM SERVICE, INC MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

WESTERN FARM SERVICE, INC.

3075 Citrus Circle
Walnut Creek, CA 94598
EPA Reg No. 11656-8 AA
EPA Est No. 11656-CA-2

See Packaged Label for Use Directions

NET CONTENTS: _____ GALLONS

ACCEPTED

DEC 17 1985

11656-8

BEST AVAILABLE COPY

GENERAL DIRECTIONS

Use designated amount of product in the following gallons water per acre unless otherwise noted
GROUND APPLICATION: Vegetable and Fruit Crops—30 to 250 gals. Cotton—3 to 20 gals. Deciduous Fruit Crops—50 to 300 gals. Citrus—400 to 3000 gals. (On Young Fruit Plantings—Use a minimum of 15 gals. water per acre)

Where Recommended for AIR APPLICATION: Fruit Crops—3 to 10 gallons
 Begin application at first sign of insects. Repeat as necessary. Do not apply within 4 days of harvest unless otherwise noted. Do not use with highly alkaline materials such as lime or bordeaux.

MIXING DIRECTIONS: Add to spray tank when 1/4 to 1/2 full with agitate turning and maintain agitation until spraying is completed.

DO NOT SPRAY ON NECTARINES, ORNAMENTAL, CHERRIES, LIQUIDAMBAR OR CHRYSANTHEMUMS AS INJURY MAY OCCUR.

USE AND APPLICATION DIRECTIONS

BEANS (Dry or Field), SOYBEANS, LIMA BEANS AND PEAS (For Processing): Aphid, Looper, Aonid, Spider Mite, Lygus, Leaf Miner, GREEN BEANS: Aonid, Spider Mite, Looper, THYRUS, Leaf-hoppers—1 pt. per acre in water to cover. 100 gals. per acre is preferable. Large Looper, Stem Bugs—1 1/2 pt. per acre in water to cover.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KALE, COLLARDS: Imported Cabbageworm, Diamondback Caterpillars, Aonid—1 pt. per acre. Looper—2 pts. per acre. May be used one day before harvest, except do not apply to kale and collards within four days of harvest.

CELERY: Aphid, Looper, Armyworm, Salt Marsh Caterpillar, Leaf Miner—Use ground application only with fine mist spray—1 pt. in 50 to 75 gals. water per acre on plants less than 1/2 grown. On larger plants use 1 1/2 pts. in 100 to 150 gals. water per acre.

COTTON: Lygus—1/2 to 1 pt. per acre by air or ground. Leaf Perler, Fleahopper, Armyworm, Bollworm, Looper and for Suppression of Mites—1 pt. per acre by air or ground. Apply at 7 to 10 days before harvest. Do not apply to cotton plants less than 1/2 grown. Do not apply by air or ground with other insecticides registered for use on cotton.

CUCUMBERS, SUMMER SQUASH: Cucumber Beetle, Aonid, Spider Mite, Looper, Thrip, Army worm—1 pt. per acre in water to cover. Pickleworm, Leaf Miner—1 1/2 pts. per acre in water to cover. By Air—Use minimum of 10 gals. water per acre. May be used up to harvest.

EGGPLANT, PEPPERS: Aonid, Spider Mite, Blister Beetle, Flea Beetle, Leaf Miner, Whitefly—1 pt. per acre. May be used up to 1 day before harvest.

WALNUTS: Aphid—1/2 to 1 pt. in 100 gals. water. Use up to ten days of harvest.

GRAPES: Drosophila, Leafhopper—1/2 to 1/4 pt. per 100 gals. water (100 to 130 gals. diluted spray per acre depending on size of vine). Use only 1/2 pt. per 100 gals. water on Muscat varieties. Do not use on other varieties as fruit injury may occur. Do not enter vineyards for 24 hours following treatment.

HOPS: Aonid, Spider Mite, Corn Earworm, Hop Looper, Caterpillar—1 pt. per acre.

SPINACH, CHARD, TURNIP GREENS: Aonid, Looper, Armyworm, Salt Marsh Caterpillar, Leaf Miner—Use ground application only with fine mist spray—1 pt. in 50 to 75 gals. water per acre on plants less than 1/2 grown. On larger plants, use 1 1/2 pts. in 100 to 150 gals. water per acre. May be used 1 day before harvest.

MELONS (Cantaloupe, Honeydew, Watermelon), PUMPKINS, WINTER SQUASH: Looper, Armyworm, Leafhopper, Spider Mite, Leaf Miner—1 to 2 pts. per acre (200 gals. diluted spray per acre). May be used 1 day before harvest. Do not apply when temperature is over 90°F.

ORANGES, LEMONS, GRAPEFRUIT, TANGERINES: Aphid, Soft Brown Scale, Citrus Greening, Citrus Leaf Miner (Tortrix)—1 pt. in 100 gals. water. Thrip (Blowfly)—1/2 pt. per 100 gals. water. Use 10 to 7 days of harvest. (California and Arizona—Do not apply within 7 days of harvest). Do not enter citrus groves for 24 hours following treatment.

PEACHES (Western States Only): Lygus, Suppression of Mites, Peach Twig Borer, Oriental Fruit Moth (California Only)—1/2 pt. in 100 gals. water. To avoid fruit spotting do not use within 30 days of harvest. Do not apply when temperature is over 90°F. Do not enter orchards for 24 hours following treatment.

RICE: Grasshopper—1/2 to 1/4 pt. per acre. Apply at first sign of insects. Do not make more than 3 applications. May be used 2 days before harvest.

STRAWBERRIES: Leafhopper, Spider Mite, Aonid, Omnivorous Leaf Tier, Spittlebug, Northwest—1 pt. per acre. May be used 1 day before harvest.

SUGAR BEETS: Spider Mite, Leafhopper—1 pt. per acre by air or ground. By Air—Use in 1 to 5 gals. water per acre. Do not apply within 5 days of harvest.

TOBACCO: Hornworm—1 pt. in 100 gals. water using up to 200 gals. diluted spray per acre. Flea Beetle—1 pt. per acre in water to cover. Apply at first signs of insects. Repeat as necessary.

TOMATOES (Field): Fruit Fly (Drosophila)—1 pt. in no less than 40 gals. water per acre with ground sprayer 7 to 10 gals. by air. Make first application 5 to 7 days before first picking and continue at 5 to 7 day intervals until the day before harvest. Field and Greenhouse Tomato Fruit Worm, Hornworm, Leaf Miner, Flea Beetle, Mite—1 pt. in 100 gals. water. Spray plants thoroughly. For Field—Use up to 200 gals. diluted spray per acre. Do not apply when temperatures is over 90°F. May be used 1 day before harvest.

GREENHOUSE, VAPOR TREATMENT OF BEES, OTHER ORNAMENTAL PLANTS, TOMATOES, CUCUMBERS: Spider Mite, Adult Whitefly, Aonid, Leafminer, Mealybug—1 R oz. per 10,000 cu. ft. when plants are dry. Apply undiluted on heat pipes (or in open parts on hot plates). Apply continuously or to intermittent sections of pipe on each side and on 1 or more of the pipes through center of each range depending upon width. Application to cold pipes should be followed by immediately heating pipes to 160°F. In steam-heated houses, close house for at least 1 hour after pipes are hot. With hot water systems close house for at least 3 hours. Houses may remain closed overnight following treatment. When ventilation is made to hot pipes wear goggles and approved respirator. Ventilate house before workers re-enter. Wear respirator when entering treated houses to ventilate. Protect hives of bees in cucumber and other houses by covering with plastic or rubber sheet. During treatment and until house is ventilated. Do not apply to tomatoes or cucumbers closer than 1 day before harvest. Spider Mite, Resistant and Non-Resistant—Make 3 to 4 treatments at 3 or 4 day intervals. Repeat as necessary. Adult Whiteflies—Make 2 applications at 7-day intervals. Repeat as necessary. Aphid—Apply when aphids appear. Repeat as necessary. **WARNING:** Avoid over-treatment and direct application to plants as injury may result. White Butterfly and Golden Rostrule Moth, Pine Chinch Bug, Chrysanthemum Green Wasp, Wandering Jew, Pansy Moth and Dutchman's Pipe may be injured by DIBROM. Water.

FRUIT FLIES (Drosophila sp.) in and Around Food Processing Plants, Landing Becks, Cell Floor and Balance Areas, Clear Mills: Use 2 pts. in 2 1/2 gals. water (5 pts. in 100 gals.) as a coarse spray to walls, floors, drains, windows, refuse and cell floor areas. Do not apply to food processing machinery. Do not apply to full top of vats or bins to be fed to livestock. Avoid contamination of feeds, feedstuffs and food processing machinery. Do not apply when plants are in operation or when feeds are present or exposed. Do not spray surfaces which will come into contact with feeds. Cover feed containers during spraying.

ADULT MOSQUITO, Gnat, Stablefly (Gnat Fly) AND HOUSEFLY CONTROL—Residential Areas, Crops Listed on Label, Total Wasteb, Swamps, Woodland, Unstock Pasture, Feed Lots and Pastures including Dairy Cattle: Consult your State Fish and Game Agency before applying this product. It is not necessary to avoid farm buildings and dairy barns. Make applications during peak of infestation and repeat as necessary. See crop recommendations for use limitations near harvest. This rate of application will kill shrimp. Do not apply to head or marsh waters which are important shrimp-producing areas. **AIRCRAFT:** East—Apply 0.8 to 4 R oz. per acre diluted with water. Apply 2 to 8 pts. of diluted spray per acre. West—Apply 1.6 to 4 R oz. per acre diluted with water. Apply 2 to 8 pts. of diluted spray per acre. **MIST OR COLD FOG:** Use 3 to 5 pts. per 100 gals. water. Calibrate equipment (rate of travel and output) to apply 0.1 to 0.25 lb. technical material per acre.

For outdoor adult mosquito control treat shrubbery and vegetation where mosquitoes may rest. Shrubby and vegetation around stagnant pools, marshy areas, ponds and spray areas, may be treated. Application of this product to any body of water is prohibited.

CORALS, ADJACENT PASTURES, HOLDING PENS (Bary salt Blue Crabs, Nept. Shrimp, Prawns): Mosquitoes, Houseflies (Aircraft Application)—Use 1.6 to 4 R oz. per acre. Dilute to 2 1/2 pts. in 10 gals. water. Apply 1 gal. diluted spray per acre. Apply over scrubby animals present.

CLOVER BITES, GOACHES, EARWIGS AND LEAFHOPPERS: Outside Buildings and in Livestock—Use 1 R oz. in 3 gals. water. Apply to turf and to soil surfaces around, beneath, shingles and eaves for general pest clean up.

NOTICE OF WARRANTY

WESTERN FARM SERVICE, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions, in accord with the statements made on this label.

WESTERN FARM SERVICE, INC.

3075 Citrus Circle
 Walnut Creek, CA 94598

EPA Reg. No. 11656-8 AA

EPA Est. No. 11656-CA-2

DEPT. OF AGRICULTURE