

PRECAUTIONARY STATEMENTS
 Hazards to Humans & Domestic Animals

DANGER:
 Keep Children and Animals Off Treated Areas. Concentrate causes eye and skin damage. May be fatal if swallowed or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Do not breathe spray mist. Use water-proof gloves and face shield or goggles when handling concentrate. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT
If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If On Skin: Immediately remove all contaminated clothing including shoes. Wash all contaminated areas, including hair, with soap and plenty of water. Do not reuse contaminated clothing or shoes until thoroughly cleaned. Seek medical attention if symptoms of poisoning develop.

If in Eyes: Immediately flush eyes with plenty of water for at least 15 minutes. Seek medical attention.

Note to Physician: Naled is a cholinesterase inhibitor, atropine is antidotal, 2-PAM may be used in conjunction with atropine in cases of severe poisoning. Mucosal damage may occur, and if gastric lavage is indicated, it should be performed cautiously.

ENVIRONMENTAL HAZARDS
 This pesticide is toxic to fish, aquatic invertebrates, and wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water or wetlands. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use with highly alkaline materials. Shrimp and crabs may also be killed at application rates recommended. Do not apply to tidal or marsh waters which are important shrimp producing areas. Do not apply when weather conditions favor drift from areas treated.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

REENTRY STATEMENT

Do not reapply this product in such a manner as to directly or through drift expose workers or other persons. The areas being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

ORAL WARNINGS shall inform workers of areas or fields that may not be entered, without protective equipment or clothing specified on this label, during treatment or until sprays have dried. Follow first aid treatment instructions shown on this label in case of accidental exposure.

When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

WRITTEN WARNINGS must include the following information:

DANGER: Area treated with Dibrom on (Date) _____.

Do not enter without appropriate protective clothing until sprays have dried. Follow first aid treatment instructions shown on this product label in case of accidental exposure.

STORAGE AND DISPOSAL

PROHIBITIONS:
 Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE:
 Keep pesticide in original container. Do not use, pour, spill or store near heat, or open flame.

Do not store diluted spray. Make new dilution for each use. Wash equipment thoroughly after use.

Do not store or transport near food or feed.


Store in cool dry place.

Not for use or storage in or around the home.

For help with any spill, leak, fire or exposure involving material, call day or night 1-800-474-9300.

PESTICIDE DISPOSAL:
 Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:
 Metal: Triple rinse (or equivalent). Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
 Glass: Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved state and local procedures.



DIBROM® 8

EMULSIVE NALED INSECTICIDE

Contains 8 lbs. Technical Naled Per Gal.
 Equivalent to 7.2 lbs. Naled.

ACTIVE INGREDIENTS	By Wt
Naled	58%
Xylene	20%
INERT INGREDIENTS	22%
TOTAL	100%

U.S. Pat. No. 2,971,887
 1,2-dibromo-2,2-dichloroethyl dimethyl phosphate

KEEP OUT OF REACH OF CHILDREN
DANGER—PELIGRO
 FOR AGRICULTURAL USE ONLY. DO NOT USE OR STORE IN OR AROUND THE HOME, OR IN OTHER AREAS ACCESSIBLE TO CHILDREN.
 PRECAUCION AL USUARIO. Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NOTICE OF WARRANTY

WESTERN FARM SERVICE, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

WESTERN FARM SERVICE, INC.
 3075 Citrus Circle
 Walnut Creek, CA 94598
 EPA Reg No. 11656-8 AA
 EPA Est. No. 11656 CA 2
 See Packaged Label for Use Directions
NET CONTENTS: _____ GALLONS

BEST AVAILABLE COPY

GENERAL DIRECTIONS

Use designated amount of product in the following gallons water per acre unless otherwise noted.
GROUND APPLICATION: Vegetables and Fruit Crops—30 to 250 gals. Corn—3 to 20 gals. Deciduous Fruit Crops—50 to 300 gals. Citrus—400 to 3000 gals. (On Young Fruit Plantings—Use a minimum of 15 gals. water per acre.)

Where Recommended for AIR APPLICATION: Fruit Crops—3 to 10 gallons.

Begin application at first sign of insects. Repeat as necessary. Do not apply within 4 days of harvest unless otherwise noted. Do not use with highly alkaline materials such as lime or bordeaux.

MIXING DIRECTIONS: Add to spray tank when 1/4 to 1/2 full with agitant running and maintain until spraying is completed.

DO NOT SPRAY ON NECTARINES, ORNAMENTAL CHERRIES, LIQUIDAMBAR OR CHRYSANTHEMUMS AS INJURY MAY OCCUR.

USE AND APPLICATION DIRECTIONS

BEANS (Dry or Field): SOYBEANS, Lima BEANS AND PEAS (For Processing): Aphids, Spider Mites, Leafhoppers, Leaf Miners. **GREEN BEANS:** Aphids, Spider Mites, Leafhoppers, Leaf Miners. 1 pt. per acre in water to cover 100 gals. per acre is preferable. Large Leafhoppers: Same as above. 1 1/2 pt. per acre in water to cover.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CANTIFLOWER, KALE, COLLARDS, Imported Cabbageworm, Cabbage Looper, Caterpillars, Aphids—1 pt. per acre. Looper—2 gts. per acre. May be used one day before harvest, except do not spray to head and collards within four days of harvest.

CELERY: Aphids, Looper, Armyworms, Salt Marsh Caterpillars, Leaf Miners—Use ground application only with fine mist spray—1 pt. in 50 to 75 gals. water per acre on plants less than 1/2 grown. On larger plants use 1 1/2 pts. in 100 to 150 gals. water per acre.

CITRUS: Lycaon—1/2 to 1 pt. per acre by air or ground. Leaf Perforator, Fleahopper, Armyworms, etc.—1 pt. per acre by air or ground.

CORN: European Cornborer, Corn Earworm, Fall Armyworm, etc.—1 pt. per acre by air or ground with other insecticides registered for use on corn.

CUCUMBERS, SUMMER SQUASH: Cucumber Beetles, Aphids, Spider Mites, Looper, Thrips, Armyworms—1 pt. per acre in water to cover. Pickworms, Leaf Miners—1 1/2 pts. per acre in water to cover. Do not use minimum of 10 gals. water per acre. May be used up to harvest.

EGGPLANT, PEPPERS, Aphids, Spider Mites, Beetle Beetles, Flea Beetles, Leaf Miners, Weevils—1 pt. per acre. May be used up to 1 day before harvest.

WALNUTS: Aphids—1/2 to 1 pt. in 100 gals. water. Use up to ten days of harvest.

GRAPES: Oriental Leafhoppers—1/2 to 1/4 pt. per 100 gals. water (100 to 130 gals. diluted spray per acre depending on size of vines). Use only 1/4 pt. per 100 gals. water on Muscad grapes. Do not use on table varieties as fruit injury may occur. Do not enter vineyards for 24 hours following treatment.

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, Cutworm—1 pt. per acre.

SPRUCH, CHARD, TURNIP GREENS: Aphids, Looper, Armyworms, Salt Marsh Caterpillars, Leaf Miners—Use ground application only with fine mist spray—1 pt. in 50 to 75 gal. water per acre on plants less than 1/2 grown. On larger plants, use 1 1/2 pts. in 100 to 150 gals. water per acre. May be used 1 day before harvest.

VEGETABLES (Cauliflower, Broccoli, Brussels Sprouts, Westland), PUMPKINS, WINTER SQUASH: Looper, Armyworms, Leafhoppers, etc. or Weevils, Leaf Miners—1 to 2 pts. per acre (200 gals. diluted spray per acre). May be used 1 day before harvest. Do not apply when temperature is over 90°F.

ORANGES, LEMONS, GRAPEFRUIT, TANGERINES, Aphids, Soft Brown Scale, Citrus Greening, Leaf Miner (Tortrix)—1 pt. in 100 gals. water. Thrips (Biont)—1/2 pt. per 100 gals. water. Use up to 7 days of harvest. California and Arizona—Do not apply within 1 day of harvest. Do not enter citrus groves or 24 hours following treatment.

PEACHES (Western States Only): Ligus, Suppression of Mites, Peach Twig Borer, Oriental Fruit Moth (California Only)—1/2 pt. in 100 gals. water. To avoid fruit spotting, do not use within 30 days of harvest. Do not apply when temperature is over 90°F. Do not enter orchards for 24 hours following treatment.

RICE: Grasshoppers—1/2 to 1/4 pt. per acre. Apply at first sign of insects. Do not make more than 3 applications. May be used 2 days before harvest.

STRAWBERRIES: Leafhopper, Spider Mites, Aphids. In various Leaf for Spottedbugs, Northwest—1 pt. per acre. May be used 1 day before harvest.

SUGAR BEETS: Spider mites, Leafhoppers—1 pt. per acre by air or ground. Do not use in 1 to 5 gals. water per acre. Do not apply within 5 days of harvest.

TOBACCO: Hornworm—1 pt. in 100 gals. water. Using up to 200 gals. diluted spray per acre. Flea Beetles—1 pt. per acre in water to cover. Apply at first sign of insects. Repeat as necessary.

TOMATOES (Field): Fruit Flies (Onionroot)—1 pt. in no less than 40 gals. water per acre with ground sprayer 7 to 10 gals. by air. Make first application 5 to 7 days before first picking and continue at 5 to 7 day intervals until the day before harvest. Field and Greenhouse Tomato Fruit Worm, Hornworms, Leaf Miners, Flea Beetles, Mites—1 pt. in 100 gals. water. Spray plants thoroughly. For Field—Use up to 200 gals. diluted spray per acre. Do not apply when temperature is over 90°F. May be used 1 day before harvest.

GREENHOUSE VAPOR TREATMENT OF ROSES, OTHER ORNAMENTAL PLANTS, TOMATOES, CUCUMBERS: Spider Mites, Adult Whiteflies, Aphids, Leafhoppers, Mealybugs—1 R. oz. per 10,000 cu. ft. when plants are dry. Apply undiluted on heat pipes (or in open pans on hot plates). Apply continuously at 20 intermittent sections of pipe on each side and on 1 or more at the pipes through center of each range depending upon width. Application to cold pipes should be followed by immediately heating pipes to 160°F. In steam-heated houses close vents for at least 1 hour after pipes are hot. With hot water systems close house for at least 3 hours. Houses may remain closed overnight following treatment. When application is made to hot pipes wear goggles and approved respirator. Ventilate house before workers re-enter. Wear respirator when entering treated houses to ventilate. Protect vines of bees in cucumber and other houses by covering with plastic or rubber sheet during treatment and until house is ventilated. Do not apply to tomatoes or cucumbers closer than 1 day before harvest. Spider Mites, Resistant and Non-Resistant—Make 3 to 4 treatments at 3 or 4 day intervals. Repeat as necessary. Adult Whiteflies—Make 2 applications at 7-day intervals. Repeat as necessary. Aphids—Apply when aphids appear. Repeat as necessary. **WARNING:** Avoid overtreatment and direct application to plants as injury may result. White Sulphur and San Jose Scale on Roses, Pink Champagne Chrysanthemums, Green Wandering Jew, Fuchsia and Dutchman's Pipe may be injured by DIBROM. Avoid.

FRUIT FLIES (Brownsie sp.) in and Around Food Processing Plants, Loading Berths, Cold Piles and Below Area: Cider Fly—Use 2 pts. in 250 gals. water. Diluted spray per acre.

Apply every 5 to 7 days as necessary. Do not apply to cut fruit or refuse piles to be fed to livestock. Avoid contamination of feeds, foodstuffs and food processing machinery. Do not apply when plants are in operation or when foods are present or exposed. Do not spray surfaces which will come in contact with foods. Cover food containers during spraying.

ADULT MOSQUITO, GNAT, STABLEFLY (Bog Fly) AND HOUSEFLY CONTROL—Residential Areas, Camps, Lanes, Lanes, Tidal Marshes, Swamps, Woodland, Livestock Pasture, Feed Lots and Pastures including Dairy Cattle. Consult your State Fish and Game Agency before applying this product. It is not necessary to avoid farm buildings and dairy barns. Make applications during peak of infestation and repeat as necessary. See also recommendations for use limitations near harvest.

As rate of application will kill shrimp. Do not apply to tidal or marsh waters which are important shrimp-producing areas. **AIRCRAFT:** Earth—Apply 0.8 to 4 R. ozs. per acre diluted with water. Apply 2 to 8 gts. of diluted spray per acre. Water—Apply 1.6 to 4 R. ozs. per acre diluted with water. Apply 2 to 8 gts. of diluted spray per acre. **MIST OR COLD FOG:** Use 3 to 5 gts. per 100 gals. water. Calibrate equipment (rate of travel and output) to apply 0.1 to 0.25 lb. technical base per acre.

For outdoor adult mosquito control trim shrubbery and vegetation where mosquitoes may rest. Shrubby and wet or around stagnant pools, marshy areas, ponds and shore lines may be treated. Application of this product to any body of water is prohibited.

CORN, ADJACENT PASTURES, HOUSING PENS (Dairy and Beef Cattle, Hogs, Sheep, Horses): Houseflies, Mosquitoes (Aerial Application)—Use 1.6 to 4 R. ozs. per acre. Dilute 1 to 2 1/2 pts. in 10 gals. water. Apply 1 gal. diluted spray per acre. Apply over areas with animals present.

CLOVER MITES, ROACHES, EARWIGS AND LEAFHOPPERS: Damp, Damp and Dry—Use 1 R. oz. in 3 gals. water. Apply to turf and to soil surfaces, among flowers, shrubs and trees for general pest control.

NOTICE OF WARRANTY

WESTERN FARM SERVICE, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

WESTERN FARM SERVICE, INC.

3075 Citrus Circle
Walnut Creek, CA 94598

EPA Reg. No. 11656-8 AA

EPA Est. No. 11656-CA-2